

**Is non-operative management the best first-line option for high-grade renal trauma?  
A systematic review.**

**AUTHORS / COLLABORATORS (Trauma EAU GUIDELINE PANEL):**

Arunan Sujenthiran, Department of Urology, St George's Healthcare NHS Trust, London, UK.

Elshout Pieter Jan, Department of Urology, Ghent University Hospital, Ghent, Belgium.

Erik Veskimae, Department of Urology, Tampere University Hospital, Tampere, Finland.

Steven MacLennan, Academic Urology Unit, University of Aberdeen, Scotland, United Kingdom.

Yuhong Yuan, Department of Medicine, Health Science Centre, McMaster University, Hamilton, Ontario, Canada.

Efraim Serafetinidis Department of Urology, Asklepieion General Hospital, Athens, Greece.

D.M.Sharma Department of Urology, St George's Healthcare NHS Trust, London, UK.

N.D. Kitrey Department of Urology, Chaim Sheba Medical Centre, Tel-Hashomer, Israel

N. Djakovic, Department of Urology, Muhldorf General Hospital, Muhldorf am Inn, Germany

N.Lumen, Department of Urology, Ghent University Hospital, Ghent, Belgium

FE.Kuehhas, London Andrology Institute, London, UK

D.J. Summerton, University Hospitals of Leicester NHS Trust, Leicester, UK

1 **Abstract**

2 **Context:**

3 The management of high-grade (grade IV-V) renal injuries remains controversial. There has been  
4 an increase in the use of non-operative management (NOM) but limited data exists comparing  
5 outcomes to open surgical exploration.

6 **Objective:** To conduct a systematic review to determine if NOM is the best first-line option for high-  
7 grade renal trauma in terms of safety and effectiveness.

8 **Evidence acquisition:**

9 Medline, Embase and Cochrane Library were searched for all relevant publications, without time or  
10 language limitations. The primary harm outcome was overall mortality and the primary benefit  
11 outcome was renal preservation rate. Secondary outcomes included length of hospital stay and  
12 complication rate. Single-arm studies were included as there were few comparative studies. Only  
13 studies with more than fifty patients were included. Data were narratively synthesised in light of  
14 methodological and clinical heterogeneity.

15 **Evidence synthesis:**

16 Seven non-randomized comparative and four single-arm studies were selected for data-extraction.  
17 787 patients were included from the comparative studies with 535 patients in the NOM group and  
18 252 in the open surgical exploration group. A further 825 patients were included from single-arm  
19 studies. Results from comparative studies: Overall mortality: NOM (0-3%), open surgical exploration  
20 (0-29%); renal preservation rate: NOM (84-100%), open surgical exploration (0-82%); complication  
21 rate: NOM (5-32%), open surgical exploration (10-76%). Overall mortality and renal preservation  
22 rate were significantly better in the NOM group whereas there was no statistical difference with  
23 regard to complication rate. Length of hospital stay was found be significantly reduced in the NOM  
24 group. Patients in the open surgical exploration group were more likely to have grade V injuries,  
25 have a lower systolic blood pressure and higher injury severity score on admission.

26 **Conclusion:**

27 No randomized controlled trials were identified and significant heterogeneity existed with regard to  
28 outcome reporting. However, NOM appeared to be safe and effective in a stable patient with a higher  
29 renal preservation rate, a shorter length of stay and a comparable complication rate to open surgical  
30 exploration. Overall mortality was higher in the open surgical exploration group though this was likely  
31 due to selection bias.

32 **Patient summary:**

33 The data of this systematic review suggest NOM continues to be favoured to surgical exploration in  
34 the management of high-grade renal trauma whenever possible. However, comparisons between  
35 both interventions are difficult as patients who have surgery are often more seriously injured than  
36 those managed non-operatively, and existing studies do not report on outcomes consistently.

37

38 **Keywords:**

39 High-grade renal injury, surgical exploration, conservative, non-operative management

40

41 **Total Word count:** 3956

42

43 **1. INTRODUCTION**

44

45 The kidney is the most commonly injured genito-urinary organ and occurs in approximately 1-5% of  
46 all trauma cases (1, 2). Renal injury can be classified as blunt or penetrating according to mechanism  
47 and by grade according to the American Association for the Surgery of Trauma (AAST) organ injury  
48 severity scale (Table 1) (3). Most cases of blunt renal trauma are low-grade injuries (grade I-III) and  
49 can be managed conservatively (4). There appears to be a trend towards the management of high-  
50 grade (IV-V) blunt renal trauma non-operatively, however strong comparative evidence is lacking in  
51 this cohort. Penetrating renal injuries have traditionally been managed with open surgical exploration  
52 though some studies have reported favourable outcomes with non-operative management (NOM),  
53 even in high-grade penetrating injuries (5, 6).

54 This shift towards NOM has been driven by rapid uptake of minimally-invasive techniques such as  
55 angioembolisation; improved clinical pathways; enhanced critical care treatment for trauma patients;  
56 readily accessible CT-imaging and a validated renal injury scoring system. Despite these advances,  
57 the optimal management of high-grade renal trauma still remains controversial with those supporting  
58 open surgical exploration reporting fewer complications (7-10) whereas advocates of NOM  
59 highlighting that conservative and minimally-invasive techniques reduce the inherent risk of  
60 nephrectomy and subsequent deterioration of renal function (11-16).

61 Current guidelines on management of high-grade renal trauma are based on retrospective  
62 comparative studies and single-arm case series' (17, 18). Existing reviews have not focused on high-  
63 grade injury and most were not conducted systematically (19) (4, 20). A systematic review of current  
64 evidence is required to establish whether the outcomes of open surgical exploration and NOM are  
65 comparable.

66 The objective of this systematic review was to compare NOM which encompasses  
67 angioembolisation, ureteric stenting and conservative management against open surgical  
68 exploration, in the management of high-grade renal injuries.

69

70 **2. EVIDENCE ACQUISITION**

71

72 The systematic review protocol was registered with PROSPERO.  
73 ([http://www.crd.york.ac.uk/PROSPERO/display\\_record.asp?ID=CRD42016035255](http://www.crd.york.ac.uk/PROSPERO/display_record.asp?ID=CRD42016035255))

74 **2.1 Search strategy and selection criteria**

75 The review was performed according to Preferred Reporting Items for Systematic Reviews and  
76 Meta-analysis (PRISMA)(21). Studies (January 1, 1946, to June 1, 2016) were identified by highly  
77 sensitive searches of electronic databases (Medline, Medline In-Process, Embase, Cochrane library  
78 databases). The initial literature search was performed in April 24 2015 and an updated search  
79 performed in June 03 2016. The search strategy is described in detail in Supplementary File 1.  
80 Animal studies, children, case reports and letters were excluded.

81 **2.2 Types of study design included**

82 There was no restriction on types of study design. Single-arm studies were included as there were  
83 only a small number of non-randomized comparative studies. All studies required a minimum of 50  
84 patients and there were no restrictions on language or date of publication.

85 **2.3 Types of participants**

86 The study population was adults ( $\geq 18$  years) with high-grade (grade IV to V according to AAST  
87 classification) CT-confirmed blunt and penetrating injuries.

88 **2.4 Types of Intervention**

89 The control group was open surgical exploration. The experimental group consisted of patients who  
90 received NOM which included conservative (supportive management only); minimally invasive  
91 intervention (angioembolisation, ureteric stent insertion, percutaneous drainage); "Package of care"  
92 involving step-wise approach (i.e. starting with conservative, followed by minimally invasive and/or  
93 surgical exploration if necessary).

94 **2.5 Types of outcome measures**

95 The primary harm outcome was mortality (overall and renal trauma-related). The primary benefit  
96 outcome was renal preservation (i.e. kidney removal or complete embolization versus preservation).  
97 Secondary outcomes included complications and length of hospital stay. Identified confounders  
98 included systolic blood pressure, injury severity score, renal function, blood loss, re-intervention rate  
99 and development of hypertension.

100 **2.6 Data collection and data extraction**

101 Following de-duplication of abstracts, two reviewers (A.S. and P.JE.) screened all abstracts and full-  
102 text articles independently. Disagreement was resolved by a third party (E.V.). References cited in  
103 all full-text articles were also assessed for additional relevant articles. A standardized data-extraction  
104 form was developed a priori to collect information on study design, renal injury details, patient  
105 characteristics and outcomes measures.

## 106 **2.7 Risk of bias in individual studies**

107 Two reviewers (A.S. and P.JE.) assessed the “risk of bias” of each included study independently.  
108 Any disagreements were resolved by discussion or by consulting a third review author.

109 Risk of bias in non-randomized comparative studies was evaluated using a modified version of a  
110 recommended tool used in the Cochrane Handbook for Systematic Reviews of Interventions. This  
111 was a pragmatic approach based on methodological literature (22, 23) and included an additional  
112 domain to assess the risk of confounding bias. A list of the 5 most important potential confounders  
113 for harm and benefit outcomes was developed a priori with clinical content experts (European  
114 Association Urology (EAU) Trauma guideline panel). The confounding factors were: Type of injury  
115 (blunt/penetrating), associated injuries, haemodynamic stability of patient, patient fitness and  
116 available interventions. This approach is detailed in our study protocol(24)

117 For single-arm studies, risk of attrition bias, whether an a priori protocol was available (indicating  
118 prospective design) and selective outcome reporting were assessed. External validity was also  
119 addressed by assessing whether study participants were selected consecutively or representative  
120 of a wider patient population. This too is a pragmatic approach informed by methodological literature  
121 (25, 26).

## 122 **2.8 Statistical analysis**

123 Meta-analysis could not be performed due to methodological and clinical heterogeneity of the  
124 included studies. Therefore a narrative synthesis was performed instead  
125 (<https://www.york.ac.uk/crd/guidance/>). Forest plots of risk difference were constructed for  
126 comparative studies for three outcome measures (mortality, complications and renal preservation).  
127 This was not done for length of stay since standard deviations were not consistently reported in the  
128 included studies. Statistical methods of assessing heterogeneity were not feasible therefore potential  
129 reasons for heterogeneity were explored in relation to population differences between, outcome  
130 definitions as well as the methods used to report outcomes. Planned formal subgroup analyses were  
131 not possible due to inclusion of non-randomized controlled studies. Therefore, any subgroup  
132 differences were discussed narratively to explore potential effect size differences. The planned  
133 sensitivity analysis to assess the robustness of our review results, by repeating the analysis only  
134 including studies with an overall medium to low risk of bias, was also not performed due to the  
135 inclusion of non-randomized comparative studies.

136 **3. EVIDENCE SYNTHESIS**

137

138 **3.1. Quality of the studies**

139 A total of 1,375 studies were identified by the literature search and two reviewers screened all study  
140 abstracts independently. Of these, 54 articles were selected for full-text screening and 11 studies (7  
141 non-randomized comparative studies, 4 single-arm studies) were eligible for inclusion (Figure 1).  
142 The quality of studies was assessed as described above. Risk of bias is summarized for comparative  
143 studies in Figure 2 and for single-arm studies in Figure 3. Overall there was a high risk of bias across  
144 both comparative and single-arm studies. Study design was retrospective for all studies. Although  
145 some studies prospectively inputted data into database, they were still retrospective in study design  
146 (27-31).

147 **3.2 Study details**

148 Three of the comparative studies included penetrating and blunt injuries and four only reported on  
149 blunt injuries. All single-arm studies reported on patients who had received NOM for blunt injuries.  
150 The recruitment period ranged from 1981-2015 and studies were published from 2006-2015. Most  
151 studies were performed at trauma centres although three were from a general hospital (29, 32, 33).  
152 Most studies were performed in a single-centre. One study was performed across two centres,  
153 another across 12 and a multi-centre study used data from 331 units (National Trauma Database  
154 Bank).

155 **3.2.1 Participants**

156 In total, 787 patients were included from the comparative studies with 535 patients in the NOM group  
157 and 252 in the open surgical exploration group. Four studies included both grade IV and V injuries  
158 (28, 30, 32, 34) and two studies only included grade IV injuries (29, 35). Sarani et al. classified grade  
159 III to V as high-grade injuries with a mean grade injury of 4 and 3.9 in the NOM and open surgical  
160 exploration group, respectively. Apart from grade, there was no strict exclusion criteria stated in most  
161 selected studies. One study excluded patients below fifteen years old and those who died before  
162 arrival to the hospital. Sarani et al. excluded patients who had a laparotomy without pre-operative  
163 CT.

164 Allocation to the different treatment groups was not randomized in any of the studies. Six studies  
165 opted for open surgical exploration if the patient was haemodynamically unstable at presentation  
166 and/or was not responding to resuscitation (28-30, 32, 34, 35). Other indications for open surgical  
167 exploration in these studies included peritonitis, failed embolization, persistent bleeding, an  
168 expanding or pulsatile haematoma, and polytrauma patients in haemorrhagic shock. One study did  
169 not specify indications for open surgical exploration (36). Three studies followed an institutional first-

170 line NOM protocol (28, 30, 31) with one study explicitly stating that even unstable patients should  
171 receive angioembolisation as first line therapy (30).

172 825 patients were included from single-arm studies with blunt injuries and received NOM. Three  
173 studies included only grade IV injuries (31, 37, 38) and one study included grade III-V studies (33).  
174 Of these studies only Long et al. stated the use of a first line non-operative protocol whereby NOM,  
175 including angioembolisation in haemodynamically unstable patients, was preferred and open  
176 surgical exploration was only performed if immediate resuscitation failed (31). There was a lack of  
177 consistency with regard to which outcomes were reported and how they were measured in  
178 comparative and single-arm studies. Only three of the comparative studies reported on all four study  
179 outcome measures (29, 32, 34)

### 180 **3.3. Outcomes**

#### 181 **3.3.1. Mortality**

182 Five comparative studies reported on overall mortality (29, 30, 32, 34, 36). A significant difference in  
183 overall mortality existed in favour of NOM in two studies (34, 36) (Table 4). Van der Wilden et al.  
184 reported 3 (2%) patients with renal-related deaths but did not compare rates between NOM and  
185 open surgical exploration. Buckley and Shoobridge both reported that both deaths in the NOM group  
186 were not renal trauma-related therefore there was no difference found between groups in these two  
187 studies with regard to renal-trauma related mortality. Only one case series reported overall mortality  
188 and it was 21% in the NOM group (37). No included studies reported the specific time-to-death  
189 following renal injury. 4 out of the 5 studies that reported on overall mortality, used in-hospital  
190 mortality (30, 32, 34, 36).

#### 191 **3.3.2. Renal Preservation**

192 Four comparative studies provided data on renal preservation (28, 29, 32, 34). In all four studies,  
193 renal preservation rate was higher in NOM (range 84%- 100%) compared to open surgical  
194 exploration (range 0%-82%) and in three of these studies there was a significant risk difference in  
195 favour of NOM (Table 4).

#### 196 **3.3.3 Complications**

197 Six comparative studies provided data on complications. In terms of absolute rates, four studies  
198 found a higher complication rate in NOM groups and two studies found patients who underwent open  
199 surgical exploration had a higher complication rate. However, only two studies reported a significant  
200 difference between groups and showed a lower rate in NOM (30) (Table 4). The 3 studies that  
201 reported lower complication rates in the open surgical exploration cohort showed no statistical  
202 difference compared to NOM.

203 Although all studies specifically reported on renal-related complications, there was a large amount  
204 of heterogeneity in their classification and reporting. Only one study used a recognized grading  
205 system (Clavien-Dindo) (30). Common complications in the NOM group included fever, haematuria,  
206 acute kidney injury and non-resolving urinomas requiring either ureteric stenting or percutaneous  
207 drainage. In the open surgical exploration group, complications included wound infection, urinary  
208 tract infection and perinephric abscess requiring drainage.

209 No included studies reported on exact time-to-event for complications, though 4 out of 5 comparative  
210 studies (30, 32, 34-36) and 3 out of 4 single-arm studies (31, 33, 38) which reported on complications  
211 used short-term in-hospital complications.

#### 212 **3.3.4 Length of Stay**

213 Six studies reported on length of stay and across these studies it was longer in open surgical  
214 exploration group (24 days) compared to NOM group (17 days). This was the trend in all the studies  
215 and two studies found there to be a statistically significant difference between the two interventions  
216 (32, 35).

#### 217 **Confounders**

218 Some confounders developed a priori including patient fitness and available interventions were not  
219 consistently reported in studies. Data was available on grade of injury, systolic blood pressure on  
220 admission and ISS in two or more studies (Table 5). There was a higher proportion of grade IV  
221 injuries in the NOM group and a higher proportion of grade V injuries in the open surgical exploration  
222 group. Two studies both found the mean systolic blood pressure to be significantly lower in the open  
223 surgical exploration group than NOM group. ISS was available in two studies and was also found to  
224 be significantly higher in the open surgical exploration group than NOM group.

#### 225 **Subgroup analysis:**

##### 226 ***Blunt versus penetrating***

227 Three studies included penetrating high-grade injuries in their population cohort. One study found  
228 that three injuries were managed successfully using conservative measures and the one patient who  
229 underwent open surgical exploration survived but required a nephrectomy (30). Two studies further  
230 divided penetrating injuries into stab and gun-shot injuries (29, 35). Both studies found that patients  
231 with gunshot injuries were the most likely to undergo surgical exploration and subsequent  
232 nephrectomy compared to stab and blunt injuries.

##### 233 ***Isolated Renal Injuries***

234 One study (29) reported on the outcomes of 43 patients who sustained isolated grade IV renal  
235 injuries. Surgical exploration was performed in 18 of 43 patients with a renal salvage rate of 83%.  
236 The remaining 25 patients were managed non-operatively with a renal salvage rate of 88%. Average

237 hospital stay was similar in both groups and transfusion rates were higher in the surgical exploration  
238 group.

## 239 240 **3.4. Discussion**

241 This is the first systematic review to use transparent and rigorous methodology to compare NOM  
242 and open surgical exploration in the management of high-grade renal trauma. In many units, first-  
243 line non-operative protocols have been implemented ahead of acquiring objective evidence due to  
244 the difficulty in conducting adequately powered randomized controlled trials. Nonetheless, this study  
245 focuses on the best available studies with population sizes greater than fifty patients, and appraises  
246 the risk of bias in a transparent way, to assess important outcomes that may not be apparent when  
247 reviewed in isolation.

### 248 **3.4.1 Principal Findings**

#### 249 *Mortality*

250 Overall mortality was found to be worse in the open surgical exploration group compared to NOM  
251 group albeit in three out of 5 comparative studies with small sample sizes and low event rates.  
252 Patients in the open surgical exploration group had higher rate of grade V injuries, higher ISS scores  
253 and lower systolic blood-pressure values on admission. Both ISS scores and lower systolic blood  
254 pressure values on admission have been shown to be predictors of increased mortality following  
255 trauma (39, 40). Therefore, this finding, together with selection bias present in most included studies  
256 whereby the most 'unstable' patients underwent open surgical exploration, could explain the  
257 difference in overall mortality between both groups. There was no evidence of a difference in renal-  
258 trauma related mortality between the two interventions in two studies (29, 30).

#### 259 *Complications*

260 Included studies rarely defined and reported complications in a consistent manner. Comparisons  
261 can still be made between interventions in the same study. Although three studies reported  
262 increasing complication rates in the NOM group, these were not statistically different. Only one study  
263 showed a statistical difference and graded complications according to the Clavien-Dindo  
264 classification (30). Given the substantial heterogeneity it is difficult to conclude that a higher  
265 complication rate exists. This is contrary to many other studies that reported a weakness of NOM to  
266 be the high frequency of short-term complications (7-9).

#### 267 *Renal Preservation*

268 Previous studies have shown that open surgical exploration can lead to higher nephrectomy rates  
269 (5, 41-43). Our data showed 84-100% of patients had preserved renal units following NOM compared

270 to a 0-82% renal-preservation rate following open surgical exploration. This finding confirms the  
271 greater risk of nephrectomy once a decision for open surgical exploration is undertaken.

272 A weakness of many studies related to renal trauma is a lack of long-term follow-up to measure  
273 residual renal function. Only one study (28) reported on relative post-operative renal function six  
274 months post-trauma using dimercapto-succinic acid renal scintigraphy (DMSA) and found poorer  
275 long-term renal function was related to percentage of devitalized parenchyma and associated  
276 visceral lesions. Studies comparing radical nephrectomy versus partial nephrectomy, although  
277 performed on a different population, provide an insight into the potential long-term negative impact  
278 of trauma nephrectomy. In selected patients, radical nephrectomy was shown to be associated with  
279 poorer survival and the development of chronic kidney disease compared to partial nephrectomy  
280 (44-46).

### 281 *Comparison with Current Guidelines*

282 Current guidelines recommend immediate intervention (open surgical exploration or  
283 angioembolisation) for haemodynamically unstable patients (18, 27). The AUA guidelines state that  
284 angioembolisation is an option only in experienced centres and surgical exploration should be used  
285 in other units. The EAU guidelines state angioembolisation is a first-line option in patients with active  
286 bleeding and no other indications for immediate open surgery. For those who do not meet the criteria  
287 for immediate intervention, AUA guidelines state that injury grade should not influence whether a  
288 patient receives surgical exploration or NOM and the EAU recommends surgical exploration only for  
289 grade V vascular injuries. These guidelines highlight the importance of clinical as well as institutional  
290 factors (angioembolisation facilities, availability of minimally invasive techniques, and level of critical  
291 care support) in deciding on the appropriate management. The current study classified  
292 angioembolisation as a non-operative intervention therefore directs comparisons to the guidelines  
293 are difficult. However, the benefits of a conservative approach to high-grade renal injury are evident.

### 294 **3.4.2 Clinical Implications**

295 The ultimate goal of conservative or minimally-invasive management is to minimize unnecessary  
296 explorations and reduce iatrogenic nephrectomy rates without increasing morbidity or mortality. This  
297 study has shown that outcomes following NOM are at the very least non-inferior to those following  
298 open surgical exploration, all while avoiding the morbidity associated with surgery. The findings from  
299 our study help to strengthen the argument for conservative management taking into account some  
300 of the absolute indications for surgical exploration that have been discussed.

301 The NOM of trauma can be viewed as a “package of care”; a step-wise approach starting with  
302 conservative, followed by minimally invasive and/or surgical exploration if necessary. It should be  
303 noted that an algorithm for “package of care” will vary in different centres according to available  
304 interventions however, the importance of escalation in treatment interventions should be  
305 emphasized.

306 **3.4.3 Limitations**

307 High-powered studies on trauma are difficult to conduct due to relatively low incidence and concerns  
308 about studies in life-threatening situations. Using retrospective comparative studies is the next best  
309 approach but remains a challenge as management has already shifted to NOM in many units. It is  
310 our belief that this review provides the first rigorously conducted systematic review on high-grade  
311 renal injury and therefore represents a review of current available best evidence.

312 There was high risk of bias in the included studies predominantly due to the retrospective study  
313 design and selection bias. Analysis of study confounders showed that patients in the open surgical  
314 exploration group were more likely to have grade V injuries, be more clinically unstable on admission  
315 and have a higher ISS compared to those in the NOM group. It is important therefore that certain  
316 outcomes heavily influenced by such confounders such as overall mortality are interpreted with  
317 caution. Mortality and complication rates were not reported on a time-to-event basis in included  
318 studies which together with small sample sizes and low event rates mean findings should be also  
319 interpreted cautiously. Although most studies reported mortality and complications that occurred “in-  
320 hospital”, the lack of defined time-periods is a key limitation. Included studies which reported on  
321 complications did not provide separate data for men in the open surgical exploration group who did  
322 not require nephrectomy. Subsequently some of the complications incurred in this group could be  
323 related specifically to the nephrectomy. However, given that most patients who underwent  
324 exploration did not require nephrectomy and that the spectra of complications with or without  
325 nephrectomy will be similar, the degree of over-estimation of complications in the exploration group  
326 will be low.

327 High grade renal injury conventionally encompasses grade IV and V renal injuries according to the  
328 AAST classification. Variation may exist across institutions on whether injuries are classified as  
329 grade IV or V dependent on reporting radiologists. Caution must be exercised when allocating a  
330 defined protocol for high-grade renal injuries when grade IV and V injuries are grouped.

331 Well-designed trials comparing these two modalities are lacking and the mainstay of reports in the  
332 literature remain retrospective case-series. The comparative observational studies identified are  
333 limited by selection bias that occurs between interventions and therefore any statistical pooling of  
334 data is misleading. Furthermore, consensus is needed regarding which outcomes are reported, how  
335 they are defined, as well as how and when they are measured. This will enable more meaningful  
336 comparisons in the evidence base in future,

337 **3.4.4 Conclusion**

338 This systematic review has provided evidence that NOM is the most appropriate first-line  
339 management option in high-grade renal trauma resulting in a renal preservation rate of  
340 approximately 84 - 100%. This systematic review has highlighted the difficulty in comparing NOM  
341 and open surgical exploration due to inherent selection bias that will remain an issue unless

342 consensus on outcome definition, measurement and reporting is achieved and adopted for future  
343 studies. The use of functional tests such as DMSA or blood parameters such as serum creatinine  
344 should be more often reported in comparative studies, if possible beyond six months. We  
345 recommend the development of prospective multi-centre trauma registers as well as standardized  
346 reporting of outcome measures to assist in making fair comparisons between studies.

347

*Supplementary File 1*

Database: EBM Reviews - Cochrane Central Register of Controlled Trials <May 2016>, EBM Reviews - Cochrane Database of Systematic Reviews <2005 to June 02, 2016>, Embase <1974 to 2016 June 03>, Epub Ahead of Print, In-Process & Other Non-Indexed Citations, Ovid MEDLINE(R) Daily and Ovid MEDLINE(R) <1946 to Present>

Search Strategy:

- 
- 1 exp kidney injury/
  
  - 2 exp Acute Kidney Injury/
  
  - 3 exp kidney/ and (exp blunt trauma/ or exp penetrating trauma/ or exp laceration/)
  
  - 4 exp Kidney/ and (exp "Wounds and Injuries"/ or exp Lacerations/)
  
  - 5 ((kidney or kidneys or renal) adj5 (trauma\* or injur\* or lesion\* or rupture\* or laceration\* or avulsion\* or contusion\* or damage\*)).tw,kw.
  
  - 6 or/1-5
  
  - 7 ((grade or grades or grading) adj5 ("4" or "5" or four or five or IV or V or "4-5" or "IV-V")).tw.
  
  - 8 ((high or higher or advance\*) adj5 (grade or grades or grading)).tw.
  
  - 9 (severe adj2 (trauma\* or injur\* or lesion\* or rupture\* or laceration\* or avulsion\* or contusion\* or damage\*)).tw. (

- 10 ((subgroup\* or sub-group or sub-analysis or sub-analyses or different or groups or categories) adj5 (grade or grading)).tw.
- 11 exp kidney pelvis/
- 12 ((renal or kidney\*) adj5 (pelvis or pelvic or ureteropelvic or hilar or hilum or collecting system)).tw.
- 13 ((urinary or urine) adj5 (extravasation or extra-vasation)).tw.
- 14 exp kidney artery/ or exp renal artery/ (21039)
- 15 exp Renal Veins/
- 16 ((renal or kidney\*) and (artery or arteries or vein or veins or vascular)).tw.
- 17 ((segmental adj2 infarction\*) or (subcapsular adj2 hematomas) or (ureteropelvic adj2 avulsion\*)).tw.
- 18 ((shattered or devasculariz\* or devascularis\*) adj5 kidney).tw.
- 19 ((major or penetrating or blunt) adj2 (trauma\* or injur\* or lesion\* or rupture\* or laceration\* or avulsion\* or contusion\* or damage\*)).tw.
- 20 or/7-19
- 21 6 and 20
- 22 exp conservative treatment/

- 23 exp minimally invasive surgery/
- 24 exp Minimally Invasive Surgical Procedures/
- 25 exp ureter stent/
- 26 exp percutaneous drainage/
- 27 (minimal\* adj5 invasive).tw.
- 28 ((ureter\* adj2 stent\*) or (percutaneous adj2 drainage)).tw.
- 29 (((angiograph\* or blood vessel or vasculograph\*) adj5 (embolization or embolisation or embolism or embolus or occlusion\*)) or embolotherap\*).tw.
- 30 (conservative or supportive or less aggressive or "not aggressive" or "non aggressive").tw.
- 31 (nonopera\* or non-opera\* or non-surgical or nonsurgical or organ sparing or without operation\* or nonresect\* or non resect\*).tw.
- 32 (package of care or step wise).tw.
- 33 or/22-32
- 34 21 and 33
- 35 ((exp animals/ or exp animal/ or exp nonhuman/ or exp animal experiment/ or animal model/ or animal tissue/ or non human/) not (humans/ or human/)) or ((rats or mice or mouse or cats or dogs or animal\* or in vitro or cell lines) not (human\* or men or women)).ti. (

36 34 not 35

37 ((child/ or Pediatrics/ or Adolescent/ or Infant/ or adolescence/ or newborn/) not adult/) or ((child or children or pediatric\* or paediatric\* or peadiatric\* or infant\* or new born or adolescent or preschool or pre-school) not (aged or adult\* or senior or men or women)).ti.

38 36 not 37

39 (case report/ or case reports/ or case report.ti.) not (cases or case series).tw.

40 38 not 39

41 remove duplicates from 40

*List of abbreviations:*

AAST: The American Association for the Surgery of Trauma

AE: Angioembolisation

CT: Computerised tomography

EAU: European Association of Urology

NOM: non-operative management

OSE: Open surgical exploration

## References

1. Meng MV, Brandes SB, McAninch JW. Renal trauma: indications and techniques for surgical exploration. *World J Urol.* 1999;17(2):71-7.
2. McAninch JW. Genitourinary trauma. *World J Urol.* 1999;17(2):65.
3. Moore EE, Shackford SR, Pachter HL, McAninch JW, Browner BD, Champion HR, et al. Organ injury scaling: spleen, liver, and kidney. *The Journal of trauma.* 1989;29(12):1664-6.
4. Broghammer JA, Fisher MB, Santucci RA. Conservative management of renal trauma: a review. *Urology.* 2007;70(4):623-9.
5. Bjurlin MA, Jeng EI, Goble SM, Doherty JC, Merlotti GJ. Comparison of nonoperative management with renorrhaphy and nephrectomy in penetrating renal injuries. *The Journal of trauma.* 2011;71(3):554-8.
6. Moolman C, Navsaria PH, Lazarus J, Pontin A, Nicol AJ. Nonoperative management of penetrating kidney injuries: a prospective audit. *The Journal of urology.* 2012;188(1):169-73.
7. Kristjánsson A, Pedersen J. Management of blunt renal trauma. *British journal of urology.* 1993;72(5 Pt 2):692-6.
8. Wilson RF, Ziegler DW. Diagnostic and treatment problems in renal injuries. *The American surgeon.* 1987;53(7):399-402.
9. Husmann DA, Gilling PJ, Perry MO, Morris JS, Boone TB. Major renal lacerations with a devitalized fragment following blunt abdominal trauma: a comparison between nonoperative (expectant) versus surgical management. *The Journal of urology.* 1993;150(6):1774-7.
10. Cass AS, Luxenberg M, Gleich P, Smith C. Long-term results of conservative and surgical management of blunt renal lacerations. *British journal of urology.* 1987;59(1):17-20.
11. Altman AL, Haas C, Dinchman KH, Spirnak JP. Selective nonoperative management of blunt grade 5 renal injury. *The Journal of urology.* 2000;164(1):27-30; discussion -1.
12. Matthews LA, Smith EM, Spirnak JP. Nonoperative treatment of major blunt renal lacerations with urinary extravasation. *J Urol.* 1997;157(6):2056-8.
13. Roberts RA, Belitsky P, Lannon SG, Mack FG, Awad SA. Conservative management of renal lacerations in blunt trauma. *Can J Surg.* 1987;30(4):253-5.
14. Danuser H, Wille S, Zoscher G, Studer U. How to treat blunt kidney ruptures: primary open surgery or conservative treatment with deferred surgery when necessary? *European urology.* 2001;39(1):9-14.
15. Moudouni SM, Patard JJ, Manunta A, Guiraud P, Guille F, Lobel B. A conservative approach to major blunt renal lacerations with urinary extravasation and devitalized renal segments. *BJU international.* 2001;87(4):290-4.
16. Robert M, Drianno N, Muir G, Delbos O, Guiter J. Management of major blunt renal lacerations: surgical or nonoperative approach? *European urology.* 1996;30(3):335-9.
17. Bryk DJ, Zhao LC. Guideline of guidelines: a review of urological trauma guidelines. *BJU international.* 2015.
18. Summerton DJ DN, Kitrey ND et al. . Guidelines on Urological Trauma European Association of Urology Guidelines 2016.
19. Santucci RA, Fisher MB. The literature increasingly supports expectant (conservative) management of renal trauma--a systematic review. *The Journal of trauma.* 2005;59(2):493-503.
20. Voelzke BB, Leddy L. The epidemiology of renal trauma. *Translational Andrology and Urology.* 2014;3(2):143-9.
21. Moher D, Liberati A, Tetzlaff J, Altman DG, Group P. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med.* 2009;6(7):e1000097.
22. Deeks JJ, Dinnes J, D'Amico R, Sowden AJ, Sakarovitch C, Song F, et al. Evaluating non-randomised intervention studies. *Health Technol Assess.* 2003;7(27):iii-x, 1-173.
23. Reeves B, Deeks JD, Higgins JPT, Wells GA. Chapter 13: Including non-randomized studies.
24. (WHO) WHO. *International Classification of Diseases* (10th revised edn.). WHO

:Geneva, 1994.

25. Viswanathan M NP, Dalberth B, Voisin C, Lohr KN, Tant E, et al. . Assessing the impact of systematic reviews on future research: two case studies. *J Comp Eff Res*. 2012;1((4):329-46.
26. Dalziel K, Round A, Stein K, Garside R, Castelnuovo E, Payne L. Do the findings of case series studies vary significantly according to methodological characteristics? *Health Technol Assess*. 2005;9(2):iii-iv, 1-146.
27. Morey AF, Brandes S, Dugi DD, Armstrong JH, Breyer BN, Broghammer JA, et al. UROTRAUMA: AUA GUIDELINE. *The Journal of urology*. 2014;192(2):327-35.
28. Lanchon C, Fiard G, Arnoux V, Descotes JL, Rambeaud JJ, Terrier N, et al. High Grade Blunt Renal Trauma: Predictors of Surgery and Long-Term Outcomes of Conservative Management. A Prospective Single Center Study. *The Journal of urology*. 2015.
29. Buckley JC, McAninch JW. Selective management of isolated and nonisolated grade IV renal injuries. *The Journal of urology*. 2006;176(6 Pt 1):2498-502; discussion 502.
30. Shoobridge JJ, Bultitude MF, Koukounaras J, Martin KE, Royce PL, Corcoran NM. A 9-year experience of renal injury at an Australian level 1 trauma centre. *BJU international*. 2013;112 Suppl 2:53-60.
31. Long JA, Fiard G, Descotes JL, Arnoux V, Arvin-Berod A, Terrier N, et al. High-grade renal injury: non-operative management of urinary extravasation and prediction of long-term outcomes. *BJU international*. 2013;111(4 Pt B):E249-55.
32. Elashry OM, Dessouky BA. Conservative Management of Major Blunt Renal Trauma with Extravasation: A Viable Option? *Eur J Trauma Emerg Surg*. 2009;35(2):115.
33. Maarouf AM, Ahmed AF, Shalaby E, Badran Y, Salem E, Zaiton F. Factors predicting the outcome of non-operative management of high-grade blunt renal trauma. *African Journal of Urology*. 2015;21(1):44-51.
34. van der Wilden GM, Velmahos GC, Joseph DK, Jacobs L, Debusk MG, Adams CA, et al. Successful nonoperative management of the most severe blunt renal injuries: a multicenter study of the research consortium of New England Centers for Trauma. *JAMA surgery*. 2013;148(10):924-31.
35. Shariat SF, Jenkins A, Roehrborn CG, Karam JA, Stage KH, Karakiewicz PI. Features and outcomes of patients with grade IV renal injury. *BJU international*. 2008;102(6):728-33.
36. Sarani B, Powell E, Taddeo J, Carr B, Patel A, Seamon M, et al. Contemporary comparison of surgical and interventional arteriography management of blunt renal injury. *J Vasc Interv Radiol*. 2011;22(5):723-8.
37. Sangthong B, Demetriades D, Martin M, Salim A, Brown C, Inaba K, et al. Management and hospital outcomes of blunt renal artery injuries: analysis of 517 patients from the National Trauma Data Bank. *J Am Coll Surg*. 2006;203(5):612-7.
38. Malaeb B, Figler B, Wessells H, Voelzke BB. Should blunt segmental vascular renal injuries be considered an AAST grade 4 renal injury? *The journal of trauma and acute care surgery*. 2014;76(2):484-7.
39. Baker SP, O'Neill B. The injury severity score: an update. *The Journal of trauma*. 1976;16(11):882-5.
40. Perel P, Prieto-Merino D, Shakur H, Clayton T, Lecky F, Bouamra O, et al. Predicting early death in patients with traumatic bleeding: development and validation of prognostic model. *BMJ (Clinical research ed)*. 2012;345:e5166.
41. McGuire J, Bultitude MF, Davis P, Koukounaras J, Royce PL, Corcoran NM. Predictors of outcome for blunt high grade renal injury treated with conservative intent. *The Journal of urology*. 2011;185(1):187-91.
42. McClung CD, Hotaling JM, Wang J, Wessells H, Voelzke BB. Contemporary trends in the immediate surgical management of renal trauma using a national database. *The journal of trauma and acute care surgery*. 2013;75(4):602-6.
43. Wright JL, Nathens AB, Rivara FP, Wessells H. Renal and extrarenal predictors of nephrectomy from the national trauma data bank. *The Journal of urology*. 2006;175(3 Pt 1):970-5; discussion 5.
44. Huang WC, Elkin EB, Levey AS, Jang TL, Russo P. Partial nephrectomy versus radical nephrectomy in patients with small renal tumors--is there a difference in mortality and cardiovascular outcomes? *The Journal of urology*. 2009;181(1):55-61; discussion -2.

45. Huang WC, Levey AS, Serio AM, Snyder M, Vickers AJ, Raj GV, et al. Chronic kidney disease after nephrectomy in patients with renal cortical tumours: a retrospective cohort study. *The lancet oncology*. 2006;7(9):735-40.
46. Thompson RH, Boorjian SA, Lohse CM, Leibovich BC, Kwon ED, Cheville JC, et al. Radical nephrectomy for pT1a renal masses may be associated with decreased overall survival compared with partial nephrectomy. *The Journal of urology*. 2008;179(2):468-71; discussion 72-3.

Figure 1: Preferred Reporting Items for Systematic Reviews and Meta-analysis flow diagram: search and study selection process for this review




Figure 2: Risk of Bias Table for non-randomised comparative studies

	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Blinding of participants and personnel (performance bias): Renal preservation	Blinding of participants and personnel (performance bias): Complications	Blinding of participants and personnel (performance bias): Mortality	Blinding of outcome assessment (detection bias): Renal preservation	Blinding of outcome assessment (detection bias): Complications	Blinding of outcome assessment (detection bias): Mortality	Incomplete outcome data (attrition bias): Renal preservation	Incomplete outcome data (attrition bias): Complications	Incomplete outcome data (attrition bias): Mortality	Selective reporting (reporting bias)	Other bias	Confounder 1: Blunt versus penetrating trauma	Confounder 2: Associated injuries	Confounder 3: Haemodynamic status	Confounder 4: Patient fitness	Confounder 5: Available interventions
Buckley 2006	-	-	-	?	+	-	-	+	+	+	+	?	-	+	+	-	-	?
Elashry 2008	-	-	-	?	+	-	?	+	+	?	+	?	-	+	?	?	-	?
Lancho 2015	-	-	?	?	?	-	?	?	-	?	?	?	-	+	+	+	-	?
Sarani 2011	-	-	?	?	+	?	?	?	?	?	+	?	-	+	+	+	-	+
Shariat 2008	-	-	-	-	+	-	-	+	?	+	?	?	-	+	+	+	-	?
Shoobridge 2012	-	+	?	?	+	?	?	+	?	+	+	?	-	+	?	?	-	?
van der Wilden 2013	-	-	-	-	+	-	-	+	+	+	+	?	-	+	+	+	-	?

Figure 3: Risk of Bias Table for single-centre studies

	A priori protocol?	Total population or consecutive patients?	Incomplete outcome data (attrition bias): Renal preservation	Incomplete outcome data (attrition bias): Complications	Incomplete outcome data (attrition bias): Mortality	Selective reporting (reporting bias)	Outcome appropriately measured (outcome measurement bias)? Renal preservation	Outcome appropriately measured (outcome measurement bias)? Complications	Outcome appropriately measured (outcome measurement bias)? Mortality
Long 2012	+	-	+	?	?	?	+	?	?
Maarouf 2014	-	-	+	?	?	?	+	?	?
Malaeb 2014	-	-	+	+	?	?	+	+	?
Sangthong 2006	-	-	?	?	+	?	?	?	+



*Table 1: The American Association for the Surgery of Trauma (AAST) kidney injury severity scale.*

<i>Grade*</i>	<i>Description of Injury</i>
<i>1</i>	<i>Contusion or non-expanding subcapsular haematoma</i>  <i>No laceration</i>
<i>2</i>	<i>Non-expanding peri-renal haematoma</i>  <i>Cortical laceration &lt;1cm deep without extravasation</i>
<i>3</i>	<i>Cortical laceration &gt;1cm without urinary extravasation</i>
<i>4</i>	<i>Laceration: through corticomedullary junction into collecting system</i>  <i>or</i>  <i>Vascular: segmental renal artery or vein injury with contained haematoma,</i>  <i>or partial vessel laceration, or vessel thrombosis</i>
<i>5</i>	<i>Laceration: shattered kidney</i>  <i>or</i>  <i>Vascular: renal pedicle or avulsion</i>

*\*Advance one grade for bilateral injuries up to grade III*

Table 2: Characteristics of Included Studies

Author	Year	Study Design	Country	Number of centres	Type of centre(s)	Recruitment period	No. of patients NOM	No. of patients open surgical exploration	Blunt / Penetrating	Outcomes reported
<i>Comparative</i>										
Buckley (29)	2006	retrospective	USA	1	General Hospital	25 y (1981-2006)	50	103	Both	Mortality, Comps, RP, LOS
Elashry (32)	2009	retrospective	Saudi Arabia	1	General Hospital	10 y (1999-2008)	51	21	Blunt only	Mortality, Comps, LOS, RP
Lanchon (28)	2015	retrospective	France	1	Trauma centre	11y (2004-2015)	148	3	Blunt only	RP
Sarani (36)	2011	retrospective	USA	2	Trauma centre	10 y (1998-2008)	20	17	Blunt only	Mortality, Comps, LOS
Shariat (35)	2008	retrospective	USA	1	Trauma centre	9 y (1997-2006)	45	32	Both	Comps, LOS
Shoobridge (30)	2013	retrospective	Australia	1	Trauma centre	9 y (2001-2010)	67	24	Both	Mortality, Comps, LOS
VanderWilden (34)	2013	retrospective	USA	12	Trauma centres	11 y (2000-2011)	154	52	Blunt only	Mortality, Comps, RP, LOS
Total							535	252		
<i>Case series</i>										
Long (31)	2012	retrospective	France	1	Trauma centre	7 y (2004-2011)	99	NA	Blunt only	Mortality, RP, LOS
Maarouf (33)	2015	retrospective	Saudi Arabia	3	General Hospitals	7y (2007-2014)	206	NA	Blunt only	RP
Malaeb (38)	2014	retrospective	USA	1	Trauma centre	7y (2003-2010)	144	NA	Blunt only	Comps, RP
Sangthong (37)	2006	retrospective	USA	331	Trauma centres	13 y (1991-2003)	376	NA	Blunt only	Mortality
Total							825			

RP: Renal preservation; Comps: Complications; LOS: Length of Stay

Table 3: Outcomes


Author	Overall Mortality N (%)				Complications N (%)				Renal Preservation N (%)			Length of Stay Days		
	NOM	OSE	Time period	p-value	NOM	OSE	p-value	Time period	NOM	OSE	p-value	NOM	OSE	p-value
<i>Comparative Studies</i>														
<b>Buckley (29)</b>	1/50 <b>(2%)</b>	0/103 <b>(0%)</b>	Not specified	-	3 /50 <b>(6%)</b>	10 /103 <b>(10%)</b>	-	Not specified	44/50 <b>(88%)</b>	84 /103 <b>(82%)</b>	-	12 <sup>^</sup>	12	-
<b>Elashry (32)</b>	0 /51 <b>(0%)</b>	3/21 <b>(14%)</b>	In-hospital	-	11/51 <b>(22%)</b>	16 /21 <b>(76%)</b>	<0.001	In-hospital	51 /51 <b>(100%)</b>	6 /21 <b>(29%)</b>	-	12 <sup>^</sup>	16	0.003
<b>Lanchon (28)</b>	NR	NR	n/a	-	NR	NR	-	n/a	124/148 <b>(84%)</b>	0 /3 <b>(0%)</b>	-	NR	NR	-
<b>Sarani (36)</b>	0/20 <b>(0%)</b>	5/17 <b>(29%)</b>	In-hospital	0.01	4 /20 <b>(20%)</b>	2 /17 <b>(12%)</b>	0.51	In-hospital	NR	NR	-	17 <sup>^</sup>	24	-
<b>Shariat (35)</b>	NR	NR	n/a	-	13 /45 <b>(28%)</b>	4/32 <b>(13%)</b>	0.2	In-hospital	NR	NR	-	7 <sup>*</sup>	12	0.001
<b>Shoobridge (30)</b>	1/67 <b>(2%)</b>	0/24 <b>(0%)</b>	In hospital	-	3 /67 <b>(5%)</b>	2 /24 <b>(8%)</b>	0.004	In-hospital	NR	NR	-	13*(AE), 11* (Cons)	20	-
<b>Van der Wilden (34)</b>	5/154 <b>(3%)</b>	12/52 <b>(23%)</b>	In-hospital	<0.01	49 /154 <b>(32%)</b>	12/52 <b>(23)</b>	0.23	In-hospital	139 /154 <b>(90%)</b>	18 /52 <b>(35%)</b>	-	13.1 <sup>^</sup>	23	-
<i>Single-arm Studies</i>														
<b>Long (31)</b>	NR	n/a	n/a	-	27/99 <b>(27%)</b>	n/a	-	In-hospital	87/99 <b>(88%)</b>	n/a	-	7 <sup>^</sup>	n/a	-
<b>Maarouf (33)</b>	NR	n/a	n/a	-	12/206 <b>(56%)</b>	n/a	-	In-hospital	189/206 <b>(92%)</b>	n/a	-	NR	n/a	-
<b>Malaeb (38)</b>	NR	n/a	n/a	-	44 /144 <b>(31%)</b>	n/a	-	In-hospital	141/144 <b>(98%)</b>	n/a	-	NR	n/a	-
<b>Sangthong (37)</b>	79/376 <b>(21%)</b>	n/a	Not specified	-	NR	n/a	-	n/a	NR	n/a	-	NR	n/a	-

\*Median, <sup>^</sup>Mean


AE: Angioembolisation; Cons: Conservative management; OSE: Open surgical exploration

Table 4 –Forest plots of risk difference between open surgical exploration (OSE) and NOM in comparative studies:

Overall Mortality



Complications



Renal Preservation




Table 5: Confounders

Author	Grade of Injury (IV/V)		Admission Systolic Blood Pressure Mean		Injury Severity Score (ISS)	
	NOM	open surgical exploration	NOM	open surgical exploration	NOM	open surgical exploration
<i>Comparative Studies</i>						
<b>Buckley (29)</b>	All Grade IV	All Grade IV	NR	NR	NR	NR
<b>Elashry (32)</b>	48 (94%) / 3 (6%)	9 (43%) / 12 (57%)	NR	NR	NR	NR
<b>Lanchon (28)</b>	124 (82%) / 27 (18%)	0 (0%) / 3 (100%)	NR	NR	NR	NR
<b>Sarani (36)</b>	4.0 (mean grade)	3.9 (mean grade)	121	100*	24	40*
<b>Shariat (35)</b>	All Grade IV	All Grade IV	NR	NR	NR	NR
<b>Shoobridge (30)</b>	53 (79%) / 14 (21%)	1 (4%) / 23 (96%)	NR	NR	25 (AE), 25(Cons)	38
<b>Van der Wilden (34)</b>	128 (83%) / 26 (17%)	26 (50%) / 26 (50%)	121	105*	23	34*
<i>Single-arm Studies</i>						
<b>Long (31)</b>	All Grade IV	NA	NR	NA	NR	NA
<b>Maarouf (33)</b>	Grade III-V	NA	NR	NA	NR	NA
<b>Malaeb (38)</b>	All Grade IV	NA	NR	NR	28(AE), 30 (Cons)	NA
<b>Sangthong (37)</b>	All Grade IV	NA	NR	NA	30	NA

AE: Angioembolisation; Cons: Conservative management

*Supplementary File 1*

Database: EBM Reviews - Cochrane Central Register of Controlled Trials <May 2016>, EBM Reviews - Cochrane Database of Systematic Reviews <2005 to June 02, 2016>, Embase <1974 to 2016 June 03>, Epub Ahead of Print, In-Process & Other Non-Indexed Citations, Ovid MEDLINE(R) Daily and Ovid MEDLINE(R) <1946 to Present>

Search Strategy:

- 
- 1 exp kidney injury/
  - 2 exp Acute Kidney Injury/
  - 3 exp kidney/ and (exp blunt trauma/ or exp penetrating trauma/ or exp laceration/) (
  - 4 exp Kidney/ and (exp "Wounds and Injuries"/ or exp Lacerations/)
  - 5 ((kidney or kidneys or renal) adj5 (trauma\* or injur\* or lesion\* or rupture\* or laceration\* or avulsion\* or contusion\* or damage\*)).tw,kw.
  - 6 or/1-5
  - 7 ((grade or grades or grading) adj5 ("4" or "5" or four or five or IV or V or "4-5" or "IV-V")).tw.
  - 8 ((high or higher or advance\*) adj5 (grade or grades or grading)).tw.
  - 9 (severe adj2 (trauma\* or injur\* or lesion\* or rupture\* or laceration\* or avulsion\* or contusion\* or damage\*)).tw. (
  - 10 ((subgroup\* or sub-group or sub-analysis or sub-analyses or different or groups or categories) adj5 (grade or grading)).tw.

- 11 exp kidney pelvis/
- 12 ((renal or kidney\*) adj5 (pelvis or pelvic or ureteropelvic or hilar or hilum or collecting system)).tw.
- 13 ((urinary or urine) adj5 (extravasation or extra-vasation)).tw.
- 14 exp kidney artery/ or exp renal artery/ (21039)
- 15 exp Renal Veins/
- 16 ((renal or kidney\*) and (artery or arteries or vein or veins or vascular)).tw.
- 17 ((segmental adj2 infarction\*) or (subcapsular adj2 hematomas) or (ureteropelvic adj2 avulsion\*)).tw.
- 18 ((shattered or devasculariz\* or devascularis\*) adj5 kidney).tw.
- 19 ((major or penetrating or blunt) adj2 (trauma\* or injur\* or lesion\* or rupture\* or laceration\* or avulsion\* or contusion\* or damage\*)).tw.
- 20 or/7-19
- 21 6 and 20
- 22 exp conservative treatment/
- 23 exp minimally invasive surgery/

24 exp Minimally Invasive Surgical Procedures/

25 exp ureter stent/

26 exp percutaneous drainage/

27 (minimal\* adj5 invasive).tw.

28 ((ureter\* adj2 stent\*) or (percutaneous adj2 drainage)).tw.

29 (((angiograph\* or blood vessel or vasculograph\*) adj5 (embolization or embolisation or embolism or embolus or occlusion\*)) or embolotherap\*).tw.

30 (conservative or supportive or less aggressive or "not aggressive" or "non aggressive").tw.

31 (nonopera\* or non-opera\* or non-surgical or nonsurgical or organ sparing or without operation\* or nonresect\* or non resect\*).tw.

32 (package of care or step wise).tw.

33 or/22-32

34 21 and 33

35 ((exp animals/ or exp animal/ or exp nonhuman/ or exp animal experiment/ or animal model/ or animal tissue/ or non human/) not (humans/ or human/)) or ((rats or mice or mouse or cats or dogs or animal\* or in vitro or cell lines) not (human\* or men or women)).ti. (

36 34 not 35

37 ((child/ or Pediatrics/ or Adolescent/ or Infant/ or adolescence/ or newborn/) not adult/) or ((child or children or pediatric\* or paediatric\* or peadiatric\* or infant\* or new born or adolescent or preschool or pre-school) not (aged or adult\* or senior or men or women)).ti.

38 36 not 37

39 (case report/ or case reports/ or case report.ti.) not (cases or case series).tw.

40 38 not 39

41 remove duplicates from 40

*List of abbreviations:*

AAST: The American Association for the Surgery of Trauma

AE: Angioembolisation

CT: Computerised tomography

EAU: European Association of Urology

NOM: non-operative management

OSE: Open surgical exploration

## *References*