

DRINFELD CENTERS FOR BICATEGORIES

EHUD MEIR AND MARKUS SZYMIK

Received: February 3, 2015

Revised: May 20, 2015

Communicated by Joachim Cuntz

ABSTRACT. We generalize Drinfeld's notion of the center of a tensor category to bicategories. In this generality, we present a spectral sequence to compute the basic invariants of Drinfeld centers: the abelian monoid of isomorphism classes of objects, and the abelian automorphism group of its identity object. There is an associated obstruction theory that explains the difference between the Drinfeld center and the center of the classifying category. For examples, we discuss bicategories of groups and bands, rings and bimodules, as well as fusion categories.

2010 Mathematics Subject Classification: Primary 18D05; Secondary 55T99.

Keywords and Phrases: Drinfeld centers, bicategories, spectral sequences, obstruction theory, bands, bimodules, fusion categories.

INTRODUCTION

If M is a monoid, then its center $Z(M)$ is the abelian submonoid of elements that commute with all elements. Monoids are just the (small) categories with only one object. It has therefore been natural to ask for a generalization of the center construction to categories \mathbf{C} . The resulting notion is often referred to as the *Bernstein center* $Z(\mathbf{C})$ of \mathbf{C} , see [Bas68, II §2], [Mac71, II.5, Exercise 8] as well as [Ber84, 1.9], for example. It is the abelian monoid of natural transformations $\text{Id}(\mathbf{C}) \rightarrow \text{Id}(\mathbf{C})$, so that its elements are the families $(p(x): x \rightarrow x \mid x \in \mathbf{C})$ of self-maps that commute with all morphisms in \mathbf{C} . Centers in this generality have applications far beyond those provided by monoids alone. As an example which is at least as important as elementary, for every prime number p the center of the category of commutative rings in characteristic p is freely generated

by Frobenius, exhibiting Frobenius as a universal symmetry of commutative algebra in prime characteristic.

Drinfeld, Joyal, Majid, Street, and possibly others have generalized the notion of center in a different direction, from monoids to (small) monoidal (or tensor) categories, see [JS91a, Definition 3] and [Maj91, Example 3.4]. The resulting notion is often referred to as the *Drinfeld center*. Since tensor categories are just the bicategories with only one object, it is therefore natural to ask for a generalization of the Drinfeld center construction to bicategories. This will be the first achievement in the present paper.

After we have set up our conventions and notation for bicategories in Section 1, Section 2 contains our definition and the main properties of the Drinfeld center of bicategories: The center is a braided tensor category that is invariant under equivalences. In the central Section 3, we will explain a systematic method (a spectral sequence) to compute the two primary invariants of the Drinfeld center of every bicategory as a braided tensor category: the abelian monoid of isomorphism classes of objects, and the abelian group of automorphisms of its unit object.

We will also explain the relation of the Drinfeld center with a more primitive construction: the center of the classifying category. These are connected by a characteristic homomorphism (3.1) that, in general, need not be either injective or surjective. As an explanation of this phenomenon, we will see that the characteristic homomorphism can be interpreted as a fringe homomorphism of our spectral sequence. The word ‘fringe’ here refers to the fact that spectral sequences in non-linear contexts only rarely have a well-defined edge. As in [Bou89], this will lead us into an associated obstruction theory that will also be explained in detail.

The final Sections 4, 5, and 6 discuss important examples where our spectral sequence can be computed and where it sheds light on less systematic approaches to computations of centers: the 2-category of groups, where the 2-morphisms are given by conjugations, fusion categories in the sense of [ENO05], where Drinfeld centers have been in the focus from the beginning of the theory on, and the bicategory that underlies Morita theory: the bicategory of rings and bimodules .

1 A REVIEW OF BICATEGORIES

To fix notation, we review the definitions and some basic examples of 2-categories and bicategories in this section. Some useful references for this material are [Ben67], [KS74], [KV94], [Str96], [Lac10], and of course [Mac71, Chapter XII]. For clarity of exposition, we will use different font faces for ordinary categories and 2-categories/bicategories.

Ordinary categories will be denoted by boldface letters such as $\mathbf{C}, \mathbf{D}, \dots$ and their objects will be denoted by x, y, \dots . The set of morphisms in \mathbf{C} from x to y will be denoted by $\text{Mor}_{\mathbf{C}}(x, y)$, or sometimes $\mathbf{C}(x, y)$ for short. We will write $x \in \mathbf{C}$ to indicate that x is an object of \mathbf{C} . If \mathbf{C} is a small ordinary

category, then $\text{Iso}(\mathbf{C})$ will be the set of isomorphism classes of objects. If $x \in \mathbf{C}$ is any object, then $\text{Aut}_{\mathbf{C}}(x)$ will be its automorphism group in \mathbf{C} .

DEFINITION 1.1. A *2-category* (or sometimes: *strict 2-category*) is a category enriched in small categories. This is a category \mathbb{B} in which for every two objects (or 0-morphisms) x, y in \mathbb{B} the morphism set is the underlying object set of a given small category $\mathbf{Mor}_{\mathbb{B}}(x, y)$; there are identity objects, and a composition functor that satisfy the evident axioms.

EXAMPLE 1.2. A basic example of a (large) 2-category can be described as follows: It has as objects the (small) categories, and the categories of morphisms are the functor categories $\mathbf{Fun}(\mathbf{C}, \mathbf{D})$ with natural transformations as morphisms. We will write $\mathbf{End}(\mathbf{C}) = \mathbf{Fun}(\mathbf{C}, \mathbf{C})$ for short.

Bicategories (or sometimes: lax/weak 2-categories) are similar to 2-categories, except that the associativity and identity properties are not given by equalities, but by natural isomorphisms. More precisely, we have the following definition.

DEFINITION 1.3. A *bicategory* \mathbb{B} consists of

- objects (the 0-morphisms) x, y, \dots ,
- a category $\mathbf{Mor}_{\mathbb{B}}(x, y)$ of morphisms (the 1-morphisms) for every ordered pair of objects x, y of \mathbb{B} ,
- a functor, the *horizontal composition*,

$$\begin{aligned} \mathbf{Mor}_{\mathbb{B}}(y, z) \times \mathbf{Mor}_{\mathbb{B}}(x, y) &\longrightarrow \mathbf{Mor}_{\mathbb{B}}(x, z), \\ (M, N) &\longmapsto M \otimes N, \end{aligned}$$

- and an identity object $\text{Id}(x) \in \mathbf{Mor}_{\mathbb{B}}(x, x)$ for every object x .

We also require natural transformations α , λ and ρ of functors that make the canonical associativity and identity diagrams commute.

EXAMPLE 1.4. A basic example of a (large) bicategory has as objects the (small) categories, and the categories of morphisms are the bimodules (or pro-functors or distributors), see [Bor94, Proposition 7.8.2].

Bicategories will be denoted by blackboard bold letters such as $\mathbb{B}, \mathbb{C}, \dots$. The objects (that is, the 0-morphisms) of a bicategory will be denoted by small letters x, y, \dots . Objects in $\mathbf{Mor}_{\mathbb{B}}(x, y)$ (that is, the 1-morphisms in \mathbb{B}) will be denoted by capital letters M, N, \dots , and we will sometimes write $\mathbb{B}(x, y) = \mathbf{Mor}_{\mathbb{B}}(x, y)$ for short.

Every category of the form $\mathbf{Mor}_{\mathbb{B}}(x, x)$ is a tensor category with respect to \otimes and $\text{Id}(x)$:

EXAMPLE 1.5. We refer to [Mac71, Chapter VII] and [JS91b] for introductions to tensor (or monoidal) categories. Every tensor category \mathbf{M} defines a bicategory $\mathbb{B}(\mathbf{M})$ with one just object, which will be denoted by \star :

$$\mathbf{Mor}_{\mathbb{B}(\mathbf{M})}(\star, \star) = \mathbf{M}.$$

Conversely, every bicategory with precisely one object is of this form. Examples of the form $\mathbf{End}(\mathbf{C}) = \mathbf{Fun}(\mathbf{C}, \mathbf{C})$ for small categories \mathbf{C} should be thought of as typical in the sense that the product need not be symmetric. Weaker notions, such as braidings [JS93], will be described later when needed.

EXAMPLE 1.6. On the one hand, every category \mathbf{C} defines a ('discrete') bicategory $\mathbb{D}(\mathbf{C})$, where the morphism sets $\mathbf{Mor}_{\mathbf{C}}(x, y)$ from \mathbf{C} are interpreted as categories $\mathbf{Mor}_{\mathbb{D}(\mathbf{C})}(x, y)$ with only identity arrows. These examples are in fact always 2-categories.

DEFINITION 1.7. Every bicategory \mathbb{B} determines an ordinary category $\mathbf{Ho}(\mathbb{B})$, its *classifying category*, as follows [Ben67, Section 7]: The objects of $\mathbf{Ho}(\mathbb{B})$ and \mathbb{B} are the same; the morphism set from x to y in the classifying category $\mathbf{Ho}(\mathbb{B})$ is the set of isomorphism classes of objects in the morphism category $\mathbf{Mor}_{\mathbb{B}}(x, y)$. In the notation introduced before,

$$\mathbf{Mor}_{\mathbf{Ho}(\mathbb{B})}(x, y) = \mathbf{Iso}(\mathbf{Mor}_{\mathbb{B}}(x, y)).$$

The associativity and identity constraints for \mathbb{B} prove that horizontal composition provides $\mathbf{Ho}(\mathbb{B})$ with the structure of an ordinary category such that the isomorphism classes of the $\mathbf{Id}(x)$ become the identities.

REMARK 1.8. The classifying category of a bicategory is not to be confused with the *Poincaré category* of a bicategory, see [Ben67, Section 7] again.

EXAMPLE 1.9. If \mathbf{M} is a tensor category, and $\mathbb{B} = \mathbb{B}(\mathbf{M})$ is the associated bicategory with one object, then $\mathbf{Ho}(\mathbb{B})$ is the monoid $\mathbf{Iso}(\mathbf{M})$ of isomorphism classes of objects in \mathbf{M} , thought of as an ordinary category with one object.

EXAMPLE 1.10. If $\mathbb{B} = \mathbb{D}(\mathbf{C})$ is a discrete bicategory defined by an ordinary category \mathbf{C} , then the classifying category $\mathbf{Ho}(\mathbb{B}) = \mathbf{C}$ gives back the ordinary category \mathbf{C} .

EXAMPLE 1.11. There is a bicategory \mathbb{T} that has as objects the topological spaces, as 1-morphisms the continuous maps, and as 2-morphisms the homotopy classes of homotopies between them. In other words, one may think of $\mathbf{Mor}_{\mathbb{T}}(X, Y)$ as the fundamental groupoid of the space of maps $X \rightarrow Y$ (with respect to a suitable topology). The bicategory \mathbb{T} is actually a 2-category. Its classifying category $\mathbf{Ho}(\mathbb{T})$ is the homotopy category of topological spaces with respect to the (strong) homotopy equivalences.

The preceding example explains our choice of notation $\mathbf{Ho}(\mathbb{B})$ for general bicategories \mathbb{B} .

We will not recall the appropriate notions of functors and natural transformations for bicategories here, but we note that every bicategory is equivalent to a 2-category, see [MP85].

2 DRINFELD CENTERS FOR BICATEGORIES

Drinfeld centers for tensor categories were introduced independently by Drinfeld, Majid [Maj91, Example 3.4], and Joyal-Street [JS91a, Definition 3]. In this section, we extend their definition and basic properties to bicategories.

2.1 DEFINITION

Drinfeld centers for bicategories are defined as follows.

DEFINITION 2.1. Let \mathbb{B} be a (small) bicategory. Its *Drinfeld center* $\mathbf{Z}(\mathbb{B})$ is the following ordinary category. The objects in $\mathbf{Z}(\mathbb{B})$ are pairs (P, p) where

$$P = (P(x) \in \mathbf{Mor}_{\mathbb{B}}(x, x) \mid x \in \mathbb{B})$$

is a family of objects in the endomorphism (tensor) categories $\mathbf{Mor}_{\mathbb{B}}(x, x)$, one for each object $x \in \mathbb{B}$, and

$$p = (p(M) : P(y) \otimes M \xrightarrow{\cong} M \otimes P(x) \mid x, y \in \mathbb{B}, M \in \mathbf{Mor}_{\mathbb{B}}(x, y))$$

is a family of natural isomorphisms in the category $\mathbf{Mor}_{\mathbb{B}}(x, y)$, one for each object $M \in \mathbf{Mor}_{\mathbb{B}}(x, y)$. These pairs of families have to satisfy two conditions: Firstly, the isomorphism

$$p(\text{Id}(x)) : P(x) \otimes \text{Id}(x) \rightarrow \text{Id}(x) \otimes P(x)$$

is the composition of the identity constraints λ and ρ . Secondly, ignoring the obvious associativity constraints, there is an equality

$$p(M \otimes N) = (\text{id}(M) \otimes p(N))(p(M) \otimes \text{id}(N)) \tag{2.1}$$

that has to hold between morphisms in the category $\mathbf{Mor}_{\mathbb{B}}(x, z)$ for all objects $M \in \mathbf{Mor}_{\mathbb{B}}(y, z)$ and $N \in \mathbf{Mor}_{\mathbb{B}}(x, y)$ that are horizontally composable. A *morphism* from (P, p) to (Q, q) in the category $\mathbf{Z}(\mathbb{B})$ is a family of morphisms

$$f(x) : P(x) \rightarrow Q(x)$$

in the categories $\mathbf{Mor}_{\mathbb{B}}(x, x)$ such that the diagram

$$\begin{array}{ccc} P(y) \otimes M & \xrightarrow{p(M)} & M \otimes P(x) \\ f(y) \otimes \text{id}(M) \downarrow & & \downarrow \text{id}(M) \otimes f(x) \\ Q(y) \otimes M & \xrightarrow{q(M)} & M \otimes Q(x) \end{array}$$

commutes for all objects $M \in \mathbf{Mor}_{\mathbb{B}}(x, y)$. Identities and composition in the category $\mathbf{Z}(\mathbb{B})$ are defined so that there is a faithful functor

$$\mathbf{Z}(\mathbb{B}) \longrightarrow \prod_{x \in \mathbb{B}} \mathbf{Mor}_{\mathbb{B}}(x, x)$$

to the product of the endomorphism categories which is forgetful on objects.

REMARK 2.2. Equation (2.1) means that the triangles

$$\begin{array}{ccc}
 & M \otimes P(y) \otimes N & \\
 p(M) \otimes \text{id}(N) \nearrow & & \searrow \text{id}(M) \otimes p(N) \\
 P(z) \otimes M \otimes N & \xrightarrow{p(M \otimes N)} & M \otimes N \otimes P(x)
 \end{array}$$

in the category $\mathbf{Mor}_{\mathbb{B}}(x, z)$ commute on the nose, again ignoring the given associativity constraints. This means that the bottom arrow is *equal* to the composition of the other two. For bicategories as defined here, there is no other notion of *equivalence* between morphisms in $\mathbf{Mor}_{\mathbb{B}}(x, z)$, so this is the appropriate definition in our situation. Further generalization—with even higher order structure (P^0, P^1, P^2, \dots) instead of just (P, p) —is called for in higher categories. This has been developed for the context of simplicial categories in [Szy]. A detailed comparison, while clearly desirable, is not within the scope of the present text, and apart from some occasional hints such as in Remark 3.2, we will focus entirely on the categorical situation here.

EXAMPLE 2.3. By inspection, if $\mathbb{B} = \mathbb{B}(\mathbf{M})$ is a tensor category, thought of as a bicategory with one object as in Example 1.5, then our definition recovers the Drinfeld center as defined in [Maj91, Example 3.4] and [JS91a, Definition 3].

Other examples related to categories of groups, bands, and fusion categories will be discussed later, see Section 4 and Section 5, respectively.

2.2 BASIC PROPERTIES

We now list the most basic properties of Drinfeld centers for bicategories: They are invariant under equivalences, and carry canonical structures of braided tensor categories. All of these are straightforward generalizations from the one-object case of tensor categories.

PROPOSITION 2.4. *For every (small) bicategory \mathbb{B} , its Drinfeld center $\mathbf{Z}(\mathbb{B})$ has a canonical structure of a tensor category. The tensor product*

$$(P, p) \otimes (Q, q) = (P \otimes Q, p \otimes q)$$

is defined by

$$(P \otimes Q)(x) = P(x) \otimes Q(x)$$

and

$$(p \otimes q)(M) = (p(M) \otimes \text{id}(Q(x)))(\text{id}(P(y)) \otimes q(M))$$

as morphisms

$$P(y) \otimes Q(y) \otimes M \longrightarrow P(y) \otimes M \otimes Q(x) \longrightarrow M \otimes P(x) \otimes Q(x)$$

for $M \in \mathbf{Mor}_{\mathbb{B}}(x, y)$. The tensor unit is (E, e) with

$$E(x) = \text{Id}(x)$$

and

$$e(M): \text{Id}(y) \otimes M \cong M \cong M \otimes \text{Id}(x)$$

is given by the constraints of the tensor structure.

Proof. See [Kas95, XIII.4.2] for the case of tensor categories. \square

COROLLARY 2.5. *The group $\text{Aut}_{\mathbf{Z}(\mathbb{B})}(E, e)$ is abelian.*

Proof. In every (small) tensor category, the endomorphism monoid of the tensor unit is abelian. See [Kas95, XI.2.4], for example. \square

PROPOSITION 2.6. *For every (small) bicategory \mathbb{B} , its Drinfeld center $\mathbf{Z}(\mathbb{B})$ has a canonical structure of a braided tensor category. The braiding*

$$(P, p) \otimes (Q, q) \longrightarrow (Q, q) \otimes (P, p)$$

is defined by the morphisms

$$p(Q(x)): P(x) \otimes Q(x) \longrightarrow Q(x) \otimes P(x)$$

for objects $x \in \mathbb{B}$.

Proof. See again [Kas95, XIII.4.2] or [JS91a, Proposition 4] for the case of tensor categories. \square

COROLLARY 2.7. *The monoid $\text{Iso}(\mathbf{Z}(\mathbb{B}))$ is abelian.*

Proof. In every (small) braided tensor category, the monoid of isomorphism classes of objects is abelian. \square

REMARK 2.8. The Drinfeld center is invariant under equivalences of bicategories. This is shown for tensor categories in [Müg03], even under the more general hypothesis that the tensor categories are weakly monoidal Morita equivalent. This will not be needed in the following.

3 SYMMETRIES, DEFORMATIONS, AND OBSTRUCTIONS

In this section, we will explain a systematic method how to compute the two primary invariants of the Drinfeld center $\mathbf{Z}(\mathbb{B})$ of every bicategory \mathbb{B} as a braided tensor category: The abelian monoid $\text{Iso}(\mathbf{Z}(\mathbb{B}))$ of isomorphism classes of objects under \otimes , and the abelian group $\text{Aut}_{\mathbf{Z}(\mathbb{B})}(E, e)$ of automorphisms of its unit of object. We will also explain the relation of the Drinfeld center to a more primitive construction: the center of the classifying category. These are connected by the homomorphism

$$\text{Iso}(\mathbf{Z}(\mathbb{B})) \longrightarrow \mathbf{Z}(\mathbf{Ho}(\mathbb{B})) \tag{3.1}$$


that sends the isomorphism class of an object (P, p) , where P is the family $P = (P(x) \in \mathbf{Mor}_{\mathbb{B}}(x, x) \mid x \in \mathbb{B})$, to the family $[P]$ of isomorphism classes, that is $[P] = ([P(x)] \in \mathbf{Iso} \mathbf{Mor}_{\mathbb{B}}(x, x) \mid x \in \mathbb{B})$.

DEFINITION 3.1. The canonical homomorphism (3.1) is called the *characteristic homomorphism*.

In general, the characteristic homomorphism need not be either injective or surjective. As an explanation of this phenomenon, we will see that the characteristic homomorphism can be interpreted as a fringe homomorphism of a spectral sequence $(E_r^{s,t} \mid r \geq 1)$ with an associated obstruction theory. This spectral sequence will compute the abelian monoid $\mathbf{Iso}(\mathbf{Z}(\mathbb{B}))$ from $E_{\infty}^{s,t}$ with $t - s = 0$ and the abelian group $\mathbf{Aut}_{\mathbf{Z}(\mathbb{B})}(E, e)$ from $E_{\infty}^{s,t}$ with $t - s = 1$.

REMARK 3.2. The spectral sequence that we are about to construct can be motivated by the spectral sequence one of us has constructed to compute the homotopy groups of homotopy coherent centers of simplicial categories, see [Szy] and compare Remark 2.2. For the purposes of this discussion, let us begin by ignoring all non-invertible objects and morphisms. Then the groups $\mathbf{Iso}(\mathbf{Z}(\mathbb{B}))$ and $\mathbf{Aut}_{\mathbf{Z}(\mathbb{B})}(E, e)$ that we are trying to compute are the (only) non-zero homotopy groups of the nerve (or classifying space) of the Drinfeld center. On the other hand, the nerve construction can also be used to produce a simplicial category from any bicategory, and it seems plausible that the homotopy coherent center of that simplicial category, which is a space, is equivalent to the nerve of the Drinfeld center of the bicategory that we started with. Then the theory in [Szy] can be applied. See also Remarks 3.3 and 5.7. It should be noted that our deductions in here will be entirely elementary, and do not make use of algebraic topology.

The non-zero part of the E_1 page of our spectral sequence is not very populated. There are only five terms $E_1^{s,t}$ that can be non-zero, and only three d_1 differentials between them. The situation can be illustrated as follows.



Then d_2 is the last differential that may be non-zero, and working through the spectral sequence is only a two-stage process. However, we note in advance

that our terms will not necessarily carry the structure of abelian groups, as one might be used to in spectral sequences.

REMARK 3.3. The range where our spectral sequence is non-trivial can be also motivated along the lines of Remark 3.2: Since for the nerve of a groupoid only π_0 (the set of isomorphism classes of objects) and π_1 (their automorphism groups) can be non-trivial, the spectral sequence in [Szy] for the homotopy coherent center of the corresponding simplicial category will have non-zero entries in at most five places: those with $0 \leq t \leq 1$ and $-1 \leq t - s \leq 1$.

We will now describe the terms and the differentials on the E_1 page, so as to obtain a description of the E_2 page.

3.1 THE TERMS WITH $t = 0$

Let us first look at the two terms with $t = 0$.

DEFINITION 3.4. We define

$$E_1^{0,0} = \prod_{x \in \mathbb{B}} \text{Iso } \mathbb{B}(x, x), \tag{3.2}$$

which is a monoid, and

$$E_1^{1,0} = \prod_{y, z \in \mathbb{B}} \text{Iso } \mathbf{Fun}(\mathbb{B}(y, z), \mathbb{B}(y, z)), \tag{3.3}$$

which is a monoid as well.

REMARK 3.5. Let us point out a common source of confusion: Each functor $F: \mathbb{B}(y, z) \rightarrow \mathbb{B}(y, z)$ induces a map $\text{Iso}(\mathbb{B}(y, z)) \rightarrow \text{Iso}(\mathbb{B}(y, z))$ between the sets of isomorphism classes, and this map only depends on the isomorphism class of the functor F . This gives us a (tautological) homomorphism

$$\tau(y, z): \text{Iso } \mathbf{Fun}(\mathbb{B}(y, z), \mathbb{B}(y, z)) \longrightarrow \text{Map}(\text{Iso } \mathbb{B}(y, z), \text{Iso } \mathbb{B}(y, z))$$

of monoids, but this homomorphism is neither injective nor surjective in general. It is the source that is relevant in (3.3), not the less useful target.

There are two natural homomorphisms $d'_1, d''_1: E_1^{0,0} \rightarrow E_1^{1,0}$ of monoids, given by sending a family $([P(x)] \mid x \in \mathbb{B})$ of isomorphism classes of objects to either the equivalence class $[M \mapsto P(z) \otimes M]$ of the endo-functor $M \mapsto P(z) \otimes M$ or to the equivalence class of the endo-functor $[M \mapsto M \otimes P(y)]$ respectively. The differential $d_1: E_1^{0,0} \rightarrow E_1^{1,0}$ should be thought of as the difference of them:

DEFINITION 3.6. The monoid $E_2^{0,0}$ is defined as the equalizer of d'_1 and d''_1 .

PROPOSITION 3.7. *There are injections*

$$E_2^{0,0} \leq \mathbf{Z}(\mathbf{Ho}(\mathbb{B})) \leq E_1^{0,0}$$

of monoids.

Proof. On the one hand, an element in $E_2^{0,0}$ is an element $E_1^{0,0}$ that lies in the equalizer of d'_1 and d''_1 . These are the families $([P(x)] \mid x \in \mathbb{B})$ of isomorphism classes of objects such that the two endo-functors $M \mapsto P(z) \otimes M$ and $M \mapsto P(z) \otimes M$ are *naturally* isomorphic. On the other hand, an element in the center $Z(\mathbf{Ho}(\mathbb{B}))$ of the classifying category is just a family $([P(x)] \mid x \in \mathbb{B})$ such that the objects $P(z) \otimes M$ and $P(z) \otimes M$ are isomorphic (perhaps not naturally) for all M .

Finally, notice that the center $Z(\mathbf{Ho}(\mathbb{B}))$ of the classifying category can be considered as the equalizer of the maps $\prod_{y,z} \tau(y,z)d'_1$ and $\prod_{y,z} \tau(y,z)d''_1$, where the maps $\tau(y,z)$ are the tautological maps defined in Remark 3.5. \square

3.2 THE TERMS WITH $t = 1$

Let us now look at the three terms with $t = 1$.

DEFINITION 3.8. We define

$$E_1^{0,1} = \prod_{x \in \mathbb{B}} \text{Aut}_{\mathbb{B}(x,x)}(\text{Id}(x)) \tag{3.4}$$

and

$$E_1^{1,1} = \prod_{y,z \in \mathbb{B}} \text{Aut}_{\mathbf{End}(\mathbb{B}(y,z))}(\text{id}), \tag{3.5}$$

which are both abelian groups.

Again, there are two distinguished homomorphisms $d'_1, d''_1: E_1^{0,1} \rightarrow E_1^{1,1}$, this time of abelian groups. One can be described as sending a family

$$u = (u(x): \text{Id}(x) \rightarrow \text{Id}(x) \mid x \in \mathbb{B})$$

of automorphisms to the natural transformation $(u(z) \otimes \text{id}(M) \mid M)$ and the other sends it to $(\text{id}(M) \otimes u(y) \mid M)$. Actually the targets are slightly different, but the tensor structure can be used to compare the two, using the diagram

$$\begin{array}{ccccc} \text{Id}(z) \otimes M & \xleftarrow{\cong} & M & \xleftarrow{\cong} & M \otimes \text{Id}(y) \\ \downarrow u(z) \otimes \text{id}(M) & & \downarrow & & \downarrow \text{id}(M) \otimes u(y) \\ \text{Id}(z) \otimes M & \xleftarrow[\cong]{} & M & \xleftarrow[\cong]{} & M \otimes \text{Id}(y). \end{array}$$

The differential $d_1: E_1^{0,1} \rightarrow E_1^{1,1}$ is the difference of d'_1 and d''_1 .

DEFINITION 3.9. We define $E_2^{0,1}$ to be the equalizer of the two homomorphisms d'_1 and d''_1 , this is the kernel of the difference $d_1 = d'_1 - d''_1$.

Direct inspection gives the following result.

PROPOSITION 3.10. *There is an isomorphism*

$$E_2^{0,1} \cong \text{Aut}_{\mathbf{Z}(\mathbb{B})}(E, e)$$

of abelian groups.

This already finishes the calculation of one of the basic invariants of $\mathbf{Z}(\mathbb{B})$ as a braided tensor category, the (abelian) automorphism group of its tensor unit.

3.3 MEASURING THE FAILURE OF INJECTIVITY

We will now proceed to calculate the (abelian) monoid $\text{Iso}(\mathbf{Z}(\mathbb{B}))$ of isomorphism classes of objects as well. In order to do so, we need to describe the remaining group on the E_1 page.

Given three objects x, y, z of \mathbb{B} , it will be useful to write

$$\mathbf{F}(x, y, z) = \mathbf{Fun}(\mathbb{B}(y, z) \times \mathbb{B}(x, y), \mathbb{B}(x, z))$$

as an abbreviation for the functor category.

DEFINITION 3.11. For any family $P = (P(x) \mid x \in \mathbb{B})$ we define

$$E_1^{2,1}(P) = \prod_{x,y,z \in \mathbb{B}} \text{Aut}_{\mathbf{F}(x,y,z)}(P \otimes ? \otimes ??), \tag{3.6}$$

where $P \otimes ? \otimes ?? : \mathbb{B}(y, z) \times \mathbb{B}(x, y) \rightarrow \mathbb{B}(x, z)$ denotes the functor

$$(M, N) \mapsto P(z) \otimes M \otimes N.$$

This is a group that may not be abelian.

It is clear that any isomorphism $P \cong Q$ of families also determines an isomorphism $E_1^{2,1}(P) \cong E_1^{2,1}(Q)$ of groups. For $P = E$, the group

$$E_1^{2,1}(E) \cong \prod_{x,y,z \in \mathbb{B}} \text{Aut}_{\mathbf{F}(x,y,z)}(\otimes)$$

receives three homomorphisms from the abelian group $E_1^{1,1}$: One sends a family

$$f = (f(M) : M \rightarrow M \mid y, z \in \mathbb{B}, M \in \mathbb{B}(y, z)) \in E_1^{1,1}$$

of natural automorphisms of the identity to the family

$$(f(M) \otimes \text{id}(N) \mid x, y, z \in \mathbb{B}, M \in \mathbb{B}(y, z), N \in \mathbb{B}(x, y)),$$

and the other ones are given similarly by

$$(\text{id}(M) \otimes f(N) \mid x, y, z \in \mathbb{B}, M \in \mathbb{B}(y, z), N \in \mathbb{B}(x, y))$$

and

$$(f(M \otimes N) \mid x, y, z \in \mathbb{B}, M \in \mathbb{B}(y, z), N \in \mathbb{B}(x, y)),$$

respectively.

Lead by Equation (2.1), we define a subgroup of the abelian group $E_1^{1,1}$ by what should be thought of as the alternating sum of these three homomorphisms:

DEFINITION 3.12.

$$Z_1^{1,1} = \{f \in E_1^{1,1} \mid f(M \otimes N) = (\text{id}(M) \otimes f(N))(f(M) \otimes \text{id}(N))\} \quad (3.7)$$

Note that

$$(\text{id}(M) \otimes f(N))(f(M) \otimes \text{id}(N)) = f(M) \otimes f(N),$$

so that we can rewrite this definition as

$$Z_1^{1,1} = \{f \in E_1^{1,1} \mid f(M \otimes N) = f(M) \otimes f(N)\}. \quad (3.8)$$

DEFINITION 3.13. We define

$$B_1^{1,1} = \{(u \otimes \text{id}(N))(\text{id}(M) \otimes u^{-1}) \mid u \in E_1^{0,1}\} \quad (3.9)$$

to be the image of the differential $d_1: E_1^{0,1} \rightarrow E_1^{1,1}$.

Recall that both of the groups $E_1^{0,1}$ and $E_1^{1,1}$ are abelian, so that taking the difference makes sense, and the image is a subgroup. It is then clear that we have $B_1^{1,1} \leq Z_1^{1,1}$.

DEFINITION 3.14. We define

$$E_2^{1,1} = Z_1^{1,1}/B_1^{1,1}, \quad (3.10)$$

which is also an abelian group.

PROPOSITION 3.15. *There is an isomorphism*

$$E_2^{1,1} \cong \text{Ker}(\text{Iso}(\mathbf{Z}(\mathbb{B})) \rightarrow \mathbf{Z}(\mathbf{Ho}(\mathbb{B})))$$

of abelian groups.

REMARK 3.16. Notice that $\text{Iso}(\mathbf{Z}(\mathbb{B}))$ and $\mathbf{Z}(\mathbf{Ho}(\mathbb{B}))$ are abelian monoids, and are not necessarily groups. However, each element from $\text{Iso}(\mathbf{Z}(\mathbb{B}))$ which maps into the identity element in $\mathbf{Z}(\mathbf{Ho}(\mathbb{B}))$ is invertible.

Proof. The kernel displayed above contains all isomorphism classes of objects $(P(x) \mid x \in \mathbb{B})$ in the Drinfeld center $\mathbf{Z}(\mathbb{B})$ such that there exists an isomorphism $P(x) \cong \text{Id}(x)$ for every object x of \mathbb{B} . We define

$$\begin{aligned} Z_1^{1,1} &\longrightarrow \text{Iso}(\mathbf{Z}(\mathbb{B})) \\ f &\longmapsto (P_f, p_f) \end{aligned}$$

as follows: Every element $f \in Z_1^{1,1}$ is a family of natural isomorphisms from the identity of $\mathbb{B}(y, z)$ to itself that is compatible with the tensor product as in (3.8). We define the image (P_f, p_f) of the element f to be the following central object: For every object $x \in \mathbb{B}$, we choose $P_f(x) = \text{Id}(x)$ to be the identity, and for every $M \in \mathbb{B}(x, y)$, the isomorphism $p_f: M \cong M \otimes P_f(x) \rightarrow P_f(y) \otimes M \cong M$


is given by $f(M): M \rightarrow M$. The fact that the family f is compatible with the tensor products ensures that this construction produces a central object.

The image (P_f, p_f) of the element f lies in the kernel of the homomorphism under consideration, and in fact, every element in the kernel is of this form. If the object (P_f, p_f) is isomorphic to the identity object, then there is a family of isomorphisms $u(x): P_f(x) = \text{Id}(x) \rightarrow \text{Id}(x)$ such that p_f is given by $(u(y) \otimes \text{id}(M))(\text{id}(M) \otimes u(x)^{-1})$, where we have used the identifications $\text{Id}(y) \otimes M \cong M \cong M \otimes \text{Id}(x)$ again. This just means that the element f lies in $B_1^{1,1}$. \square

The preceding proposition explains the potential failure of the injectivity of the characteristic homomorphism (3.1): If an element $Z(\mathbf{Ho}(\mathbb{B}))$ can be lifted to an element in $Z(\mathbb{B})$, then the abelian group $E_2^{1,1}$ acts on the different representatives in $\text{Iso}(Z(\mathbb{B}))$. However, in monoids, this action need neither be free nor transitive.

3.4 THE E_2 PAGE

The part of the E_2 page of the spectral sequence that can be non-trivial looks as follows.



The three terms that have already been calculated are

$$E_2^{0,1} \cong \text{Aut}_{Z(\mathbb{B})}(E, e)$$

in the column $t - s = 1$ and

$$E_2^{0,0} \leq Z(\mathbf{Ho}(\mathbb{B}))$$

$$E_2^{1,1} \cong \text{Ker}(\text{Iso}(Z(\mathbb{B})) \rightarrow Z(\mathbf{Ho}(\mathbb{B})))$$

in the column $t - s = 0$, so that there is an exact sequence

$$0 \longrightarrow E_2^{1,1} \longrightarrow \text{Iso}(Z(\mathbb{B})) \longrightarrow Z(\mathbf{Ho}(\mathbb{B}))$$

that describes $\text{Iso}(Z(\mathbb{B}))$, except for the image of the characteristic homomorphism.

One may guess that the image of the characteristic homomorphism would be the kernel of a differential

$$d_2: E_2^{0,0} \longrightarrow E_2^{2,1}, \quad (3.11)$$

but the situation is more complicated: The question mark in $E_2^{2,1}(?)$ in the figure indicates that we do not have a single group $E_2^{2,1}$ as the target of a differential (3.11), but rather an entire family $E_2^{2,1}(P)$ of groups, one as the target of a tailored differential that acts on $[P] \in E_2^{0,0}$. This will now be explained in detail.

3.5 OBSTRUCTIONS TO SURJECTIVITY

We now address the following question: When can an element in the center $Z(\mathbf{Ho}(\mathbb{B}))$ of the classifying category be lifted to an element in $\text{Iso}(\mathbf{Z}(\mathbb{B}))$ and hence in $\mathbf{Z}(\mathbb{B})$? Our proof of Proposition 3.7 already gives one condition: That element should lie in the submonoid $E_2^{0,0}$ that is cut out as the ‘kernel of d_1 .’ We may therefore right away start with an element in the monoid $E_2^{0,0}$. Recall from our Definition 3.6 that an element in the monoid $E_2^{0,0}$ is given by a family $(P(x) \mid x \in \mathbb{B})$ of objects for which the two functors $R_{P(x)}, L_{P(y)}: \mathbb{B}(x, y) \rightarrow \mathbb{B}(x, y)$ that are given on objects by $M \mapsto M \otimes P(x)$ and $M \mapsto P(y) \otimes M$, respectively, are naturally isomorphic. The family can be lifted to an object in the Drinfeld center $\mathbf{Z}(\mathbb{B})$ if and only if we can choose a family $p = (p_{x,y}: R_{P(x)} \rightarrow L_{P(y)} \mid x, y \in \mathbb{B})$ of (natural) isomorphisms of functors such that we have a natural isomorphism

$$p_{x,z}(M \otimes N) = (\text{id}(M) \otimes p_{x,y}(N))(p_{y,z}(M) \otimes \text{id}(N))$$

for every pair of objects $M \in \mathbb{B}(y, z)$ and $N \in \mathbb{B}(x, y)$. To measure the failure of a given p to comply to these needs, we may consider the composition

$$d_2(p)_{x,y,z}(M, N) = (\text{id}(M) \otimes p_{x,y}(N))(p_{y,z}(M) \otimes \text{id}(N))p_{x,z}^{-1}(M \otimes N), \quad (3.12)$$

which is an automorphism of the functor $(M, N) \mapsto P(z) \otimes M \otimes N$. If we let x, y , and z vary, then the family of the $d_2(p)_{x,y,z}$ defines an element $d_2(p)$ of the group $E_1^{2,1}(P)$ of our Definition 3.11. The automorphisms (3.12) is the identity automorphism if and only if (P, p) lies in the Drinfeld center $\mathbf{Z}(\mathbb{B})$. This leads us to regard the automorphisms (3.12) as the obstructions for (P, p) to be an object of the Drinfeld center.

Of course, it is possible that (P, p) will not be an object of the Drinfeld center, but (P, p') will be, for some other family $p' = (p'_{x,y})$ of isomorphisms. Since the group of automorphisms of the functor $(M, N) \mapsto P(z) \otimes M \otimes N$ is not abelian in general, we proceed as follows.

DEFINITION 3.17. We define the *obstruction differential* d_2 on P by

$$d_2(P) = \{ d_2(p) \mid p = (p_{x,y}: R_{P(x)} \rightarrow L_{P(y)} \mid x, y \in \mathbb{B}) \} \subseteq E_1^{2,1}(P),$$

where p runs over all possible isomorphisms of functors. We will say that $d_2(P)$ *vanishes* on P if it contains the identity automorphism.

The discussion preceding the definition proves the following result.

PROPOSITION 3.18. *The obstruction differential d_2 vanishes on P if for some choice of p the object (P, p) lies in the Drinfeld center $\mathbf{Z}(\mathbb{B})$.*

4 GROUPS AND THE CATEGORY OF BANDS

In this section, we present a detailed discussion of the various notions of centers, and in particular the Drinfeld center, in a situation that is genuinely different from tensor categories: categories and bicategories where the objects are (discrete) groups.

A size limitation needs to be chosen, and we can and will assume that all groups under consideration are finite. Therefore, let \mathbf{G} denote the category of all finite groups. Definition 2.1 requires a small category, so that it will be clear that the result is a set. However, our calculations will reveal that the result is a set anyway. We could also choose to work with a skeleton, and then note that the choice of skeleton does not affect the calculation, since any two skeleta are equivalent.

The category \mathbf{G} is the underlying category of a 2-category \mathbb{G} , where the category $\mathbf{Mor}_{\mathbb{G}}(G, H)$ is the groupoid of homomorphisms $G \rightarrow H$, which are the 1-morphisms of \mathbb{G} , and the 2-morphisms $h: \alpha \rightarrow \beta$ between two homomorphisms $\alpha, \beta: G \rightarrow H$ are the elements h in H that conjugate one into the other:

$$\{h \in H \mid h\alpha(g)h^{-1} = \beta(g) \text{ for all } g \in G\}.$$

REMARK 4.1. The classifying category $\mathbf{B} = \mathbf{Ho}(\mathbb{G})$ is sometimes called the category of *bands* in accordance with its use in non-abelian cohomology and the theory of gerbes, see [Gir71, IV.1]. It is customary to denote the conjugacy classes of homomorphisms $G \rightarrow H$ by

$$\text{Rep}(G, H) = \text{Iso } \mathbf{Mor}_{\mathbb{G}}(G, H) = \text{Mor}_{\mathbf{B}}(G, H).$$

These are the sets of morphisms in the classifying category $\mathbf{B} = \mathbf{Ho}(\mathbb{G})$. Note that Giraud used the notation $\text{Hex}(G, H)$ instead of our $\text{Rep}(G, H)$.

The automorphism group of any given homomorphism $\alpha: G \rightarrow H$ in the category $\mathbb{G}(G, H) = \mathbf{Mor}_{\mathbb{G}}(G, H)$ of homomorphisms is the centralizer

$$\text{Aut}_{\mathbb{G}(G, H)}(\alpha) = \{h \in H \mid h\alpha(g)h^{-1} = \alpha(g) \text{ for all } g \in G\} = \text{C}(\alpha) \quad (4.1)$$

of the image of α . This is a subgroup of the group H . We remark that the centralizers need not be abelian. For example, the centralizer of the constant homomorphism $G \rightarrow G$ is the entire group G , whereas the center of G reappears as the centralizer of the identity $G \rightarrow G$. These observations will be useful when we will determine the Drinfeld center of \mathbb{G} .

4.1 SOME ORDINARY CENTERS

Before we turn our attention towards the Drinfeld center, let us first describe the centers of the ordinary categories \mathbf{G} and $\mathbf{B} = \mathbf{Ho}(\mathbb{G})$.

PROPOSITION 4.2. *The centers of the categories \mathbf{G} and $\mathbf{B} = \mathbf{Ho}(\mathbb{G})$ are isomorphic to the abelian monoid $\{0, 1\}$ under multiplication.*

Proof. This is straightforward for the category \mathbf{G} of groups and homomorphisms. An element in the center thereof is a family $P = (P_G: G \rightarrow G)$ of homomorphisms that are natural in G . We can evaluate P on the full subcategory of cyclic groups, and since $\mathbf{Mor}(\mathbb{Z}/k, \mathbb{Z}/k) \cong \mathbb{Z}/k$, we see that P is determined by a profinite integer n in $\widehat{\mathbb{Z}}$: we must have $P_G(g) = g^n$ for all groups G and all of their elements g . But, if n is not 0 or 1, then there are clearly groups for which that map is not a homomorphism. In fact, we can take symmetric or alternating groups, as we will see in the course of the rest of the proof.

Let us move on to the center of the classifying category $\mathbf{B} = \mathbf{Ho}(\mathbb{G})$. Again, the homomorphisms $g \mapsto g^0$ and $g \mapsto g^1$ are in the center, and they still represent different elements, since they are not conjugate. In the classifying category, if $[P] = ([P_G]: G \rightarrow G)$ is an element in the center, testing against the cyclic groups only shows that there is a profinite integer n such that $P_G(g)$ is *conjugate* to g^n for each group G and each of its elements g . We will argue that no such family of homomorphisms P_G exists unless n is 0 or 1.

Let us call an endomorphisms $\alpha: G \rightarrow G$ on some group G of *conjugacy type* n if $\alpha(g)$ is conjugate to g^n for all g in G . We need to show that for all n different from 0 and 1 there exists at least one group G that does not admit an endomorphism of conjugacy type n .

If $|n| \geq 2$, then we choose $m \geq \max\{n, 5\}$ and consider the subgroup G of the symmetric group $S(m)$ generated by the elements of order n . Since the set of generators is invariant under conjugation, this subgroup is normal, and it follows that $G = A(m)$ (the subgroup of alternating permutations) or $G = S(m)$. An endomorphism $\alpha: G \rightarrow G$ of conjugacy type n would have to be trivial because it vanishes on the generators. But then g^n would be trivial for all elements g in $A(m) \leq G$, which is absurd.

If $n = -1$, then we first note that an endomorphism $\alpha: G \rightarrow G$ of conjugacy type -1 is automatically injective. Hence, if G is finite, then it is an automorphism. Therefore we choose a nontrivial finite group G of odd order such that its outer automorphism group is trivial. (Such groups exist, see [Hor74], [Dar75] or [Hei96] for examples that also have trivial centers.) If $\alpha: G \rightarrow G$ were an endomorphism of conjugacy type -1 , then this would be an inner automorphism by the assumption on G . Then $\text{id}: G \rightarrow G$, which represents the same class, would also be an endomorphism of conjugacy type -1 . In other words, every element g would be conjugate to its inverse g^{-1} , a contradiction since the order of G is odd. \square

4.2 THE DRINFELD CENTER

Now that we have evaluated the centers of the ordinary categories of groups and bands, we are ready to apply the obstruction theory and spectral sequence introduced in Section 3 in order to determine the Drinfeld center $\mathbf{Z}(\mathbb{G})$ of the 2-category \mathbb{G} of groups. The following result describes the two basic invariants of $\mathbf{Z}(\mathbb{G})$.

PROPOSITION 4.3. *The maps*

$$\mathbf{Z}(\mathbb{G}) \longrightarrow \text{Iso } \mathbf{Z}(\mathbb{G}) \longrightarrow \mathbf{Z}(\mathbf{B} = \mathbf{Ho}(\mathbb{G}))$$

are both isomorphisms, and the automorphism group of the identity object in $\mathbf{Z}(\mathbb{G})$ is trivial.

Proof. Let us start with the one entry in the spectral sequence that has $t = 0$. We already know that there is an upper bound $E_2^{0,0} \leq \mathbf{Z}(\mathbf{B} = \mathbf{Ho}(\mathbb{G}))$ by Proposition 3.7. The center of the classifying category has been determined in the preceding Proposition 4.2. That result also makes it clear that all elements in the center of the classifying category lift to the Drinfeld center of \mathbb{G} ; they even lift to the center of the underlying category \mathbf{G} . We deduce that the obstructions vanish.

Let us now deal with the two entries in the spectral sequence that have $t = 1$ and that determine the kernel of the map $\text{Iso } \mathbf{Z}(\mathbb{G}) \rightarrow \mathbf{Z}(\mathbf{B} = \mathbf{Ho}(\mathbb{G}))$ and the automorphism group of the identity object in $\mathbf{Z}(\mathbb{G})$. We have

$$E_1^{0,1} = \prod_F \text{Aut}_{\mathbb{G}(F,F)}(\text{id}) \cong \prod_F \mathbf{Z}(F)$$

by (4.1), and we record that this is an abelian group.

In order to determine the entry $E_1^{1,1}$, we start with the definition:

$$E_1^{1,1} = \prod_{G,H} \text{Aut}_{\mathbf{End}(\mathbb{G}(G,H))}(\text{id}).$$

We know that the category $\mathbb{G}(G, H) = \mathbf{Mor}_{\mathbb{G}}(G, H)$ is a groupoid, and as such it is equivalent to the sum of the groups $C(\alpha)$, where α runs through a system of representatives of $\text{Rep}(G, H)$ in $\mathbf{Mor}_{\mathbb{G}}(G, H)$. Since we are considering automorphisms of the identity object, we get

$$\text{Aut}_{\mathbf{End}(\mathbb{G}(G,H))}(\text{id}) \cong \prod_{[\alpha] \in \text{Rep}(G,H)} \mathbf{Z}C(\alpha),$$

the product of the centers of the centralizers. We note again that this is an abelian group. This leaves us with

$$E_1^{1,1} \cong \prod_{G,H} \prod_{[\alpha] \in \text{Rep}(G,H)} \mathbf{Z}C(\alpha).$$

The differential $d_1: E_1^{0,1} \rightarrow E_1^{1,1}$ is the difference of the two coface homomorphisms. Therefore, up to an irrelevant sign, it is given on a family $P = (P(F) \in Z(F) | F)$ by

$$(d_1 P)(\alpha) = P(H) - \alpha(P(G)) \in ZC(\alpha) \quad (4.2)$$

in the factor of $\alpha: G \rightarrow H$. It follows that the $E_2^{0,1}$ entry in the spectral sequence consists of those families P such that $P(H) = \alpha(P(G))$ for all G, H , and $\alpha: G \rightarrow H$. Taking α to be constant, we see that $P(F)$ has to be trivial for all F . This shows $E_2^{0,1} = 0$. Therefore, by Proposition 3.10, we deduce that $\text{Aut}_{\mathbf{Z}(\mathbb{G})}(E, e)$ is indeed trivial.

It remains to be shown that there are no more components than we already know. Proposition 3.15 says that these are indexed by the group $E_2^{1,1}$. This group can be calculated as follows. Its elements are represented by elements in the subgroup $Z_1^{1,1} \leq E_1^{1,1}$, the subgroup of elements $Q = (Q(\alpha) \in ZC(\alpha) | \alpha)$ such that

$$Q(\gamma\beta) = Q(\gamma) + \gamma(Q(\beta)), \quad (4.3)$$

again up to an irrelevant sign. We claim that each family with that property is already in the distinguished subgroup $B_1^{1,1}$, so that $Z_1^{1,1} = B_1^{1,1}$ and $E_2^{1,1} = Z_1^{1,1}/B_1^{1,1} = 0$.

That subgroup $B_1^{1,1}$ is the image of the differential d_1 . Therefore, to prove the claim, let us be given a family $Q = (Q(\alpha) \in ZC(\alpha) | \alpha)$ such that (4.3) holds. We can then evaluate this family at the unique homomorphisms $\alpha = \epsilon_F$ from the trivial group to F , for each finite group F , to obtain a family $P(F) = Q(\epsilon_F)$, and that family is our candidate for an element P to hit the element Q under the differential d_1 . And indeed, equation (4.3) for $\alpha = \gamma$ and $\beta = \epsilon_G$ gives

$$Q(\epsilon_H) = Q(\alpha\epsilon_G) = Q(\alpha) + \alpha(Q(\epsilon_G)).$$

Rearranging this yields the identity

$$Q(\alpha) = Q(\epsilon_H) - \alpha(Q(\epsilon_G)) = P(H) - \alpha(P(G)) = (d_1 P)(\alpha),$$

and this shows that Q is indeed in the image. We have proved the claim. \square

It seems reasonable to expect that similar arguments will determine the Drinfeld centers of related bicategories such as the ones coming from groupoids or topological spaces, etc. This will not be pursued further here. Instead, we will now turn our attention towards a class of examples that indicates the wealth of obstructions and nontrivial differentials that one can expect in general.

5 APPLICATIONS TO FUSION CATEGORIES

Fusion categories are tensor categories with particularly nice properties. They arise in many areas of mathematics and mathematical physics, such as operator algebras, conformal field theory, and Hopf algebras. A general theory of such

categories has been developed in [ENO05]. In this section, we use the theory developed so far in order to explain some constructions related to the centers of fusion categories that otherwise may seem to appear *ad hoc*. For completeness, let us start with the definition.

DEFINITION 5.1. A *fusion category* is a semisimple abelian tensor category (\mathbf{F}, \otimes, I) over a field \mathfrak{K} of characteristic zero, usually assumed to be algebraically closed, with finitely many simple objects, such that \otimes is bilinear, each object has a dual object, and the distinguished object I is simple.

We are here interested in the bicategory $\mathbb{B}(\mathbf{F})$ with one object that is associated with such a fusion category \mathbf{F} as explained in Example 1.5, and its Drinfeld center. Since fusion categories are special cases of tensor categories, this refers to the usual notion of a Drinfeld center of a tensor category, and as such it has been studied in other places. For example, the papers [Müg03], [Ost03], [GNN09], and [BV13] contain various results on the centers of fusion categories and related categories.

We will show that the spectral sequence introduced in Section 3 offers a systematic approach to the computation of the basic invariants of the Drinfeld center, by interpreting the different terms and differentials in the language of fusion categories.

5.1 THE FIRST PAGE

We start by identifying the terms on the first page of the spectral sequence.

PROPOSITION 5.2. *If a fusion category \mathbf{F} has n isomorphism classes of simple objects, then the monoid $E_1^{0,0}$ can be identified as a set with \mathbb{N}^n , the n -fold product of the monoid \mathbb{N} of non-negative integers. The multiplication is given by the fusion coefficients.*

Proof. The monoid $E_1^{0,0}$ is the set of isomorphism classes of objects in our category. If the different simple objects of \mathbf{F} are X_1, \dots, X_n , then there is a canonical identification of $E_1^{0,0}$ with \mathbb{N}^n , given by

$$(a_1, \dots, a_n) \longleftrightarrow X_1^{\oplus a_1} \oplus \dots \oplus X_n^{\oplus a_n}.$$

Notice that the direct sum on \mathbf{F} and the tensor product are two different operations. \square

An element of the monoid $E_1^{1,0}$ is just an isomorphism class of endofunctors $\mathbf{F} \rightarrow \mathbf{F}$.

PROPOSITION 5.3. *If a fusion category \mathbf{F} over a field \mathfrak{K} has n isomorphism classes of simple objects, with representatives X_1, \dots, X_n , then there are iso-*

morphisms

$$\begin{aligned} E_1^{0,1} &\cong \mathfrak{K}^\times \\ E_1^{1,1} &\cong (\mathfrak{K}^\times)^n \\ E_1^{2,1} &\cong \prod_{i,j} \text{Aut}(X_i \otimes X_j). \end{aligned}$$

REMARK 5.4. In every fusion category with simple objects X_1, \dots, X_n , the automorphism group of the object $X_1^{\oplus a_1} \oplus \dots \oplus X_n^{\oplus a_n}$ is the group $\prod_i \text{GL}_{a_i}(\mathfrak{K})$. All of the groups in the preceding proposition have this form.

Proof. The group $E_1^{0,1}$ is the automorphism group of the tensor identity I of \mathbf{F} . In a fusion category, the tensor identity is simple, and the endomorphism ring of each simple object is \mathfrak{K} . Therefore this group is isomorphic to \mathfrak{K}^\times .

The group $E_1^{1,1}$ is the automorphism group of the identity functor $\text{id}_{\mathbf{F}}: \mathbf{F} \rightarrow \mathbf{F}$. In the case of a fusion category, such an automorphism α is specified by giving $\alpha_i: X_i \rightarrow X_i$ for each $i = 1, \dots, n$. In other words, such an automorphism is given by a set of n invertible scalars, and the group is isomorphic with $(\mathfrak{K}^\times)^n$. Now, the two maps we have $E_1^{0,1} \rightarrow E_1^{1,1}$ are the same. They are given by the diagonal embedding $\mathfrak{K}^\times \rightarrow (\mathfrak{K}^\times)^n$.

The group $E_1^{2,1}$ is the automorphism group of the tensor product functor $\otimes: \mathbf{F} \times \mathbf{F} \rightarrow \mathbf{F}$. Any such automorphism is given by a set of n^2 invertible morphisms $\beta_{i,j}: X_i \otimes X_j \rightarrow X_i \otimes X_j$. We can thus identify this group with the product $\prod_{i,j} \text{Aut}(X_i \otimes X_j)$. \square

5.2 THE FIRST DIFFERENTIALS AND THE CENTER OF THE CLASSIFYING CATEGORY

The following result computes the first differential and the center of the classifying category.

PROPOSITION 5.5. *The abelian monoid $E_2^{0,0}$ is isomorphic to $Z(\mathbf{Ho}(\mathbb{B}(\mathbf{F})))$, the center of the classifying category.*

Proof. The first differential $E_1^{0,0} \rightarrow E_1^{1,0}$ is given by two maps, one which sends an object X to the functor $L_X: Y \mapsto X \otimes Y$ and the other one maps X to the functor $R_X: Y \mapsto Y \otimes X$. We are interested in the equalizer. This consists of all the (isomorphism classes of) objects X for which there exists an isomorphism (of functors) between L_X and R_X . For fusion categories, this is the same as the center of the classifying category. Indeed, since a fusion category \mathbf{F} is semi-simple, an additive functor $\mathbf{F} \rightarrow \mathbf{F}$ is determined by its restriction to simple objects, and the tensor product with a given object is an additive functor. \square

5.3 THE UNIVERSAL GRADING GROUP

For each fusion category \mathbf{F} , we define \mathbf{F}_{ad} to be the fusion subcategory of \mathbf{F} consisting of all the direct summands of all objects of the form $X_i \otimes X_i^*$. Notice

that we take only the products of simple objects with their duals. Then \mathbf{F} has a faithful grading by a group $U(\mathbf{F})$, the *universal grading group of \mathbf{F}* such that \mathbf{F}_{ad} is exactly the trivially graded component. Moreover, each other faithful grading of \mathbf{F} is a quotient of this grading. See [GN08] for more details.

PROPOSITION 5.6. *There are isomorphisms*

$$\begin{aligned} E_2^{0,1} &\cong \mathfrak{K}^\times \\ E_2^{1,1} &\cong \widehat{U(\mathbf{F})}, \end{aligned}$$

where $\widehat{U(\mathbf{F})}$ denotes the character group of the universal grading group of \mathbf{F} .

Proof. The first claim follows from the fact that the differential from $E_1^{0,1}$ to $E_1^{1,1}$ is zero. This fact also implies that the group $B_1^{1,1}$ is the trivial group. As for the second one, the three maps $E_1^{1,1} \rightarrow E_1^{2,1}$ are the following. The first is given by sending a family (α_i) to the family $(\alpha_i)_{i,j}$. The second sends a family (α_i) to the family $(\alpha_j)_{i,j}$. The third is more complicated: It sends a family (α_i) to $(\beta_{i,j})_{i,j}$, where $\beta_{i,j}$ acts by the scalar α_k on the X_k component inside $X_i \otimes X_j$. We can now offer two proofs, based on Proposition 3.15.

First, we can identify $Z_1^{1,1}$ and $\widehat{U(\mathbf{F})}$: For each $i = 1, \dots, n$, let $g_i \in U(\mathbf{F})$ be the degree of X_i by the universal grading. Then the element in $Z_1^{1,1}$ which corresponds to $\varphi \in \widehat{U(\mathbf{F})}$ is $(\varphi(g_i))$. We thus get an isomorphism $E_2^{1,1} \cong \widehat{U(\mathbf{F})}$. Second, we can also identify $\widehat{U(\mathbf{F})}$ directly with the kernel of the characteristic homomorphism $\text{Iso}(\mathbf{Z}(\mathbb{B}(\mathbf{F}))) \rightarrow \mathbf{Z}(\mathbf{Ho}(\mathbb{B}(\mathbf{F})))$: Characters of the universal grading group $U(\mathbf{F})$ are in one to one correspondence with objects of the Drinfeld center whose underlying object is the tensor unit. The central object corresponding to a character φ is (I, φ) , the tensor unit I , together with the isomorphism $I \otimes X_i \rightarrow X_i \otimes I$ given by the scalar $\varphi(g_i)$. Here we identify both objects with X_i in the canonical way, and the endomorphism ring of X_i is \mathfrak{K} . \square

In conclusion, we have the following exact sequence:

$$0 \longrightarrow \widehat{U(\mathbf{F})} \longrightarrow \text{Iso}(\mathbf{Z}(\mathbb{B}(\mathbf{F}))) \longrightarrow \mathbf{Z}(\mathbf{Ho}(\mathbb{B}(\mathbf{F}))) \tag{5.1}$$

for any fusion category \mathbf{F} . It identifies the character group of the universal grading group as the measure of the failure of the center of the classifying category to detect information that is contained in the richer Drinfeld center.

REMARK 5.7. The exact sequence (5.1) has been established (independently) in the work of Grossman, Jordan, and Snyder. Their preprint [GJS] gives two proofs: One is elementary and *ad hoc*; the other shows that the sequence can be embedded into the long exact sequence associated to a fibration of spaces. It would be interesting to relate that fibration to the tower of fibrations that is the origin of the spectral sequence in [Szy], compare Remark 3.2.

5.4 THE OBSTRUCTION DIFFERENTIAL

The last part which we need to understand in the spectral sequence is the obstruction differential on the second page, that is $d_2: E_2^{0,0} \rightarrow E_2^{2,1}(?)$. We have seen in Section 3.5 that the target $E_2^{2,1}(?)$ depends on the argument. Let us explain the obstruction that the differential encodes.

The monoid $E_2^{0,0}$ contains all objects X of \mathbf{F} for which the functors L_X and R_X are isomorphic. We will furnish a structure of a central object on X if we can find an isomorphism of functors $\alpha: L_X \rightarrow R_X$ such that the following diagram will be commutative for every $Y, Z \in \mathbf{F}$.

$$\begin{array}{ccccc}
 & & X \otimes (Y \otimes Z) & \xrightarrow{\alpha_Y \otimes \alpha_Z} & (Y \otimes Z) \otimes X \\
 & \nearrow & & & \searrow \\
 (X \otimes Y) \otimes Z & & & & Y \otimes (Z \otimes X) \\
 & \searrow & & & \nearrow \\
 & & (Y \otimes X) \otimes Z & \xrightarrow{\alpha_Y \otimes \text{id}_Z} & Y \otimes (X \otimes Z)
 \end{array}
 \tag{5.2}$$

In other words, if for all choices of α , the identity is in the set of loops of diagram (5.2).

Of course, different choices of α might furnish different central structures on X . If we take X to be the tensor unit, this obstruction indeed lies in $E_2^{2,1}(I)$.

5.5 VECTOR SPACES GRADED BY GROUPS

Let G be a finite group, and assume that our fusion category \mathbf{F} is \mathbf{Vec}_G^ω , that is, the category of G -graded vector spaces, in which the associativity constraint is deformed by the class $[\omega] \in H^3(G, \mathfrak{K}^\times)$ of a 3-cocycle ω , as in [ENO05, Section 2]. Briefly, the simple objects X_g in \mathbf{Vec}_G^ω are indexed by the elements g of the group G . The tensor product is given by $X_g \otimes X_h = X_{gh}$ and the associativity constraint

$$X_{ghk} = (X_g \otimes X_h) \otimes X_k \longrightarrow X_g \otimes (X_h \otimes X_k) = X_{ghk}$$

is given by the scalar $\omega(g, h, k)$. We can think of this category as the category of vector bundles over the set G , with deformed associativity constraints.

The automorphism group of the neutral object in the Drinfeld center is given by \mathfrak{K}^\times , as for every fusion category.

The abelian monoid of isomorphism classes of objects is more interesting in this case: We can view $E_1^{0,0}$ as the multiplicative monoid underlying the group semi-ring $\mathbb{N}G$, and then the elements of $E_2^{0,0}$ are exactly the central elements. The universal grading group $U(\mathbf{Vec}_G^\omega)$ is given by the group G . This shows that the characteristic homomorphism is not injective as soon as the group G admits a non-trivial character.

These observations determine the E_2 -page of the spectral sequence except for the obstruction differential.

Let us begin by understanding the obstruction differential d_2 in case we are taking a central element $z \in G$, and ask if the corresponding simple object X_z has a structure of a central object. By inspecting the diagram (5.2) above, we get that z and ω together furnish a 2-cocycle $\gamma_{\omega,z}$ on G , given by

$$\gamma_{\omega,z}(g, h) = \omega(g, h, z)\omega^{-1}(g, z, h)\omega(z, g, h).$$

Assume that an object X_z has a structure of a central object. We can then choose isomorphisms $\varphi_g: X_g \otimes X_z \rightarrow X_z \otimes X_g$ for every element g in G , such that diagram (5.2) is commutative, with $X = X_g$, $Y = X_h$ and $Z = X_z$. We have $X_z \otimes X_g = X_{zg} = X_{gz} = X_g \otimes X_z$. This means that the isomorphism φ_g can be given by a scalar. By inspecting the diagram (5.2) again, we see that its commutativity boils down to the equation $\partial(\varphi) = \gamma_{\omega,z}$. In other words, the object X_z will have a central structure if and only if the class of $\gamma_{\omega,z}$ vanishes in $H^2(G, \mathfrak{K}^\times)$. In that case, the different central structures on X_z are in one to one correspondence with characters of G , as in the case when z is the neutral element of our group G .

More generally, we can ask when the direct sum $X_z^{\oplus m}$ of m copies of X_z will have a central structure. For this, we need the twisted group algebra $\mathfrak{K}^{\gamma_{\omega,z}}G$. This \mathfrak{K} -algebra has a \mathfrak{K} -basis $\{u_g \mid g \in G\}$ and the multiplication is given by

$$u_g u_h = \gamma_{\omega,z}(g, h)u_{gh}.$$

The 2-cocycle condition ensures that this algebra is associative. The automorphism group of the object $X_z^{\oplus m}$ is isomorphic to the general linear group $GL_m(\mathfrak{K})$. The commutativity of the diagram (5.2) with $Z = X_z^{\oplus m}$ will mean that we can choose elements $v_g \in GL_m(\mathfrak{K})$ such that $v_g v_h = \gamma_{\omega,z}(g, h)v_{gh}$. In other words, the algebra $\mathfrak{K}^{\gamma_{\omega,z}}G$ has an m dimensional representation, and the different structures of central objects on $X_z^{\oplus m}$ correspond to the different isomorphism classes of those representations of the twisted group algebra $\mathfrak{K}^{\gamma_{\omega,z}}G$ that have dimension m . Since $\mathfrak{K}^{\gamma_{\omega,z}}G$ is finite-dimensional and semi-simple when $\text{char}(\mathfrak{K}) = 0$, there are only finitely many such isomorphism classes.

The general obstructions can be understood now in a similar way: If g_1, \dots, g_r are representatives of the conjugacy classes of G , then we denote by Y_i the direct sum of all X_g such that g is conjugate to g_i . We denote by y_i the corresponding sum in $\mathbb{N}G$. Then any central element in $\mathbb{N}G$ can be written as a sum $\sum_i a_i y_i$ for $a_i \in \mathbb{N}$. The possible central structures on the object $\bigoplus_i Y_i^{\oplus a_i}$ are given by tuples $([V_i])_i$ where $[V_i]$ is an isomorphism class of a representation of $\mathfrak{K}^{\gamma_{\omega,g_i}}C_G(g_i)$ of dimension a_i .

PROPOSITION 5.8. *For every group G , class $\omega \in H^3(G, \mathfrak{K}^\times)$, and central element $z \in Z(G)$ such that $\gamma_{\omega,z}$ is non trivial, the isomorphism class of the object X_z is not in the image of the characteristic homomorphism.*

EXAMPLE 5.9. Let $n \geq 2$ be any integer, and set $G = (\mathbb{Z}/n)^3$. We denote a \mathbb{Z}/n -basis of the abelian group G by $\{e_1, e_2, e_3\}$, and we write ζ for a primitive n -th root of unity in our field \mathfrak{K} . We consider the 3-cocycle ω defined by

$$\omega(e_1^{a_{11}} e_2^{a_{12}} e_3^{a_{13}}, e_1^{a_{21}} e_2^{a_{22}} e_3^{a_{23}}, e_1^{a_{31}} e_2^{a_{32}} e_3^{a_{33}}) = \zeta^{a_{11} a_{22} a_{33}}.$$

in $H^3(G, \mathfrak{K}^\times)$. It arises as the cup product of three \mathbb{Z}/n -linearly independent elements in the group $H^1(G, \mathfrak{K}^\times) \cong (\mathbb{Z}/n)^{\oplus 3}$. Since the group G is abelian, every element is central; we choose $z = e_1$. A direct calculation shows that

$$\gamma_{\omega, z}(e_1^{a_{11}} e_2^{a_{12}} e_3^{a_{13}}, e_1^{a_{21}} e_2^{a_{22}} e_3^{a_{23}}) = \zeta^{a_{12} a_{23} + a_{11} a_{22}},$$

which is non-trivial. Furthermore, the Wedderburn decomposition of the algebra $\mathfrak{K}^{\gamma_{\omega, z}} G$ is given by

$$\mathfrak{K}^{\gamma_{\omega, z}} G \cong \bigoplus_{i=1}^n M_n(\mathfrak{K})$$

Thus, the class of the m -fold direct sum $X_z^{\oplus m}$ will be in the image of the characteristic homomorphism if and only if m is a multiple of n .

6 THE BICATEGORY OF RINGS AND BIMODULES

In this section, we discuss another important example of a bicategory, this time one that has several objects, and that is not a (strict) 2-category, the bicategory \mathbb{M} of rings and bimodules, see [Ben67, 2.5] and [Mac71, XII.7]. The objects in \mathbb{M} are the (associative) rings A, B, C, \dots (with unit). The category $\mathbb{M}(A, B)$ is the category of (B, A) -bimodules M and their homomorphism. (As before in Section 4, some size limitation on the rings and bimodules is needed to keep the categories and their centers relatively small. We will not further comment on this, since it is again inessential to our calculations.) Composition in \mathbb{M} is given by the tensor product of bimodules, and the (A, A) -bimodule A is the identity object in the category $\mathbb{M}(A, A)$. Every morphism $f: A \rightarrow B$ in the (ordinary) category of rings gives rise to a (B, A) -bimodule B_f , where A acts via f . The identity object corresponds to the identity id_A in the category $\mathbb{M}(A, A)$. The (ordinary) category of rings has the ring \mathbb{Z} as an initial object, and we use this observation repeatedly in this section.

We can now begin to study our spectral sequence for the bicategory \mathbb{M} . As in Section 4, we start with the (ordinary) center of the classifying category.

PROPOSITION 6.1. *The center $Z(\mathbf{Ho}(\mathbb{M}))$ of the classifying category $\mathbf{Ho}(\mathbb{M})$ of the bicategory \mathbb{M} is the abelian monoid of isomorphism classes of abelian groups under the multiplication induced by the tensor product, with the isomorphism class of the infinite cyclic group as unit.*

Proof. Consider the monoid

$$E_1^{0,0} = \prod_{A \in \mathbb{M}} \text{Iso } \mathbb{M}(A, A).$$

An element in this monoid is given by a family of isomorphism classes of (A, A) -bimodules $(P(A) \mid A \in \mathbb{M})$. The monoid

$$E_1^{1,0} = \prod_{A,B \in \mathbb{M}} \text{Iso Fun}(\mathbb{M}(A, B), \mathbb{M}(A, B))$$

contains families of isomorphism classes of endofunctors of the category of (B, A) -bimodules, one such for every two rings A and B in \mathbb{M} . The first differential vanishes on the isomorphism class of the family $(P(A) \mid A \in \mathbb{M})$ if and only if the two functors $N \mapsto N \otimes_A P(A)$ and $N \mapsto P(B) \otimes_B N$ are isomorphic on the category of (B, A) -bimodules.

Consider in particular the case where $B = \mathbb{Z}$ is the initial ring, and the (\mathbb{Z}, A) -bimodule N is A itself. In this particular case, we see that the two (\mathbb{Z}, A) -bimodules $A \otimes_A P(A)$ and $P(\mathbb{Z}) \otimes A$ are isomorphic to each other and hence to $P(A)$. (Here and in the following we write $\otimes = \otimes_{\mathbb{Z}}$ for readability.) We see that, up to isomorphism, the entire family is determined by the abelian group $P(\mathbb{Z})$. Conversely, given any abelian group M , then the family of the $P(A) = M \otimes A$ admits the structure of a central object. In other words, the submonoid $E_2^{0,0}$ consists of all isomorphism class of families of the form

$$(P(A) \mid A \in \mathbb{M}) = (P(\mathbb{Z}) \otimes A \mid A \in \mathbb{M}),$$

where $P(\mathbb{Z})$ is an arbitrary abelian group. We thus have $E_2^{0,0} = \mathbf{Z}(\mathbf{Ho}(\mathbb{M}))$. It is clear from the definition of \mathbb{M} that the composition is given by the tensor product and the isomorphism class of the abelian group \mathbb{Z} , which corresponds to the family of isomorphism class of the (A, A) -bimodules A , is the unit. \square

We remark that we have seen in the course of the proof that the usual symmetry will furnish a central structure on any family of bimodules of the form $(P(A) = M \otimes A \mid A \in \mathbb{M})$. In particular, this holds for the unit given by the family $E = (E(A) = A)$ of (A, A) -bimodules A .

We can now direct our attention to the Drinfeld center of \mathbb{M} itself. We start by determining the automorphism group of its tensor unit.

PROPOSITION 6.2. *The automorphism group $\text{Aut}_{\mathbf{Z}(\mathbb{M})}(E, e)$ of the tensor unit (E, e) in the Drinfeld center $\mathbf{Z}(\mathbb{M})$ of the bicategory \mathbb{M} is cyclic of order 2.*

Proof. It is well known (and easy to prove) fact that the automorphism group of A as an (A, A) -bimodule is canonically isomorphic to the group $Z(A)^\times$ of invertible elements in the center of the ring A . So

$$E_1^{0,1} \cong \prod_{A \in \mathbb{M}} Z(A)^\times$$

follows. Similarly, any automorphism of the identity functor on the category of (B, A) -bimodules, which is isomorphic to the category of $B \otimes A^{\text{op}}$ -modules,

is determined by its action on the generator, so that it will be given as multiplication by an invertible element of the center of the ring $B \otimes A^{\text{op}}$. We thus have

$$E_1^{1,1} = \prod_{A,B \in \mathbb{M}} \mathbb{Z}(B \otimes A^{\text{op}})^\times.$$

Using these isomorphisms as identifications, the differential $d_1: E_1^{0,1} \rightarrow E_1^{1,1}$ then sends a family $(z_A \mid A \in \mathbb{M})$ to the family $(z_B \otimes z_A^{-1} \mid A, B \in \mathbb{M})$. It follows that, if the differential on the family $(z_A \mid A \in \mathbb{M})$ vanishes, then each element z_A must be the image in A of some invertible element in (the center of) \mathbb{Z} . Because the only such elements in \mathbb{Z} are ± 1 , we conclude that the only families in $E_2^{0,1}$ are $(1_A \mid A \in \mathbb{M})$ and $(-1_A \mid A \in \mathbb{M})$, and thus we have $\text{Aut}_{\mathbf{Z}(\mathbb{M})}(E, e) \cong E_2^{0,1} \cong \mathbb{Z}/2$. \square

It remains for us to describe the isomorphism classes of objects in the center $\mathbf{Z}(\mathbb{M})$ of \mathbb{M} . We will do this now by showing that the characteristic homomorphism is bijective, so that Proposition 6.1 gives the desired result.

PROPOSITION 6.3. *For the bicategory \mathbb{M} of rings and bimodules, the characteristic homomorphism $\text{Iso}(\mathbf{Z}(\mathbb{M})) \rightarrow \mathbf{Z}(\mathbf{Ho}(\mathbb{M}))$ is an isomorphism.*

Proof. We have already explained the fact that an element in the center $\mathbf{Z}(\mathbf{Ho}(\mathbb{M}))$ of the classifying category $\mathbf{Ho}(\mathbb{M})$ admits a structure of a central object that lives in $\mathbf{Z}(\mathbb{M})$. Therefore, all obstructions must vanish and the characteristic homomorphism is surjective.

The group $Z_1^{1,1}$ is the subgroup that consists of all families of invertible central elements $z_{A,B} \in \mathbb{Z}(B \otimes A^{\text{op}})^\times$ that satisfy the following condition: For every three rings A, B and C , the element

$$(\text{id}_C \otimes m_B \otimes \text{id}_A)(z_{B,C} \otimes z_{A,B}) \in C \otimes B \otimes A^{\text{op}}$$

is equal to $z_{A,B} \otimes 1$, where we identify $C \otimes B \otimes A^{\text{op}}$ with $C \otimes A^{\text{op}} \otimes B$ using the symmetry, and where we denote the multiplication $B \otimes B \rightarrow B$ by m_B .

Similarly to before, by studying this equation for the special case $B = \mathbb{Z}$ we learn that each such family must be of the form $z_{A,B} = z_B \otimes z_A^{-1}$ for some invertible central elements $z_A \in \mathbb{Z}(A)^\times$: Set $z_A = z_{A,\mathbb{Z}}$ and note that $z_{\mathbb{Z},A} = z_A^{-1}$.

In other words, a family in the group $Z_1^{1,1}$ will automatically be in the image of the differential d_1 , which is the subgroup $B_1^{1,1}$. We conclude that the group $E_2^{1,1}$ is trivial, and therefore the characteristic homomorphism is also injective in this case. \square

The following statement summarizes the results of this section.

THEOREM 6.4. *The Drinfeld center $\mathbf{Z}(\mathbb{M})$ of the bicategory \mathbb{M} of rings and bimodules is isomorphic to the category of abelian groups with the monoidal structure given by the usual tensor product of abelian groups, and the unit given by the group of integers.*

REMARK 6.5. The obvious generalization from the bicategory of rings (that is \mathbb{Z} -algebras) to the bicategory of R -algebras (over a given commutative ring R) can be proven with the same techniques; this only requires a more ornamented notation.

ACKNOWLEDGMENT

Both authors were supported by the Danish National Research Foundation through the Centre for Symmetry and Deformation (DNRF92).

REFERENCES

- [Bas68] H. Bass. Algebraic K-theory. W.A. Benjamin, Inc., New York-Amsterdam, 1968.
- [Ben67] J. Bénabou. Introduction to bicategories. Midwest Category Seminar 1–77. Lecture Notes in Mathematics 47. Springer, Berlin, 1967.
- [Bor94] F. Borceux. Handbook of categorical algebra. 1. Basic category theory. Encyclopedia of Mathematics and its Applications 50. Cambridge University Press, Cambridge, 1994.
- [Bou89] A. K. Bousfield. Homotopy spectral sequences and obstructions. Israel J. Math. 66 (1989) 54–104.
- [Ber84] J.-N. Bernstein. Le “centre” de Bernstein. Rédigé par P. Deligne. Travaux en Cours. Representations of reductive groups over a local field, 1–32. Hermann, Paris, 1984.
- [BV13] A. Bruguières, A. Virelizier. On the center of fusion categories. Pacific J. Math. 264 (2013) 1–30.
- [Dar75] R.S. Dark. A complete group of odd order. Math. Proc. Cambridge Philos. Soc. 77 (1975) 21–28.
- [ENO05] P.I. Etingof, D.A. Nikshych, V. Ostrik. On fusion categories. Ann. of Math. 162 (2005) 581–642.
- [ENO10] P.I. Etingof, D.A. Nikshych, V. Ostrik. Fusion categories and homotopy theory. With an appendix by E. Meir. Quantum Topol. 1 (2010) 209–273.
- [GNN09] S. Gelaki, D. Naidu, D. Nikshych. Centers of graded fusion categories. Algebra Number Theory 3 (2009) 959–990.
- [GN08] S. Gelaki, D. Nikshych. Nilpotent fusion categories. Adv. Math. 217 (2008) 1053–1071.

- [Gir71] J. Giraud. *Cohomologie non abélienne*. Die Grundlehren der mathematischen Wissenschaften 179. Springer-Verlag, Berlin-New York, 1971.
- [GJS] P. Grossman, D. Jordan, N. Snyder. Cyclic extensions of fusion categories via the Brauer-Picard groupoid.
<http://arxiv.org/abs/1211.6414>
- [Hei96] H. Heineken. Examples of complete groups of odd order. Proceedings of the 1st Panhellenic Algebra Conference (Athens, 1996). Bull. Greek Math. Soc. 38 (1996) 69–77.
- [Hor74] M.V. Horoševskiĭ. Complete groups of odd order. Algebra and Logic 13 (1974) 34–40.
- [JS91a] A. Joyal, R. Street. Tortile Yang-Baxter operators in tensor categories. J. Pure Appl. Algebra 71 (1991) 43–51.
- [JS91b] A. Joyal, R. Street. The geometry of tensor calculus. I. Adv. Math. 88 (1991) 55–112.
- [JS93] A. Joyal, R. Street. Braided tensor categories. Adv. Math. 102 (1993) 20–78.
- [KV94] M.M. Kapranov, V.A. Voevodsky. 2-categories and Zamolodchikov tetrahedra equations. Algebraic groups and their generalizations: quantum and infinite-dimensional methods (University Park 1991) 177–259. Proc. Sympos. Pure Math. 56, Part 2. Amer. Math. Soc., Providence, RI, 1994.
- [Kas95] C. Kassel. *Quantum Groups*. Graduate Texts in Mathematics 155. Springer-Verlag, New York, 1995.
- [KS74] G.M. Kelly, R. Street. Review of the elements of 2-categories. Category Seminar (Proc. Sem., Sydney, 1972/1973) 75–103. Lecture Notes in Math. 420. Springer, Berlin, 1974.
- [Lac10] S. Lack. A 2-categories companion. Towards higher categories, 105–191, IMA Vol. Math. Appl. 152. Springer, New York, 2010.
- [Mac71] S. Mac Lane. *Categories for the Working Mathematician*. Second Edition. Springer Verlag, 1971.
- [MP85] S. Mac Lane, R. Paré. Coherence for bicategories and indexed categories. J. Pure Appl. Algebra 37 (1985) 59–80.
- [Maj91] S. Majid. Representations, duals and quantum doubles of monoidal categories. Proceedings of the Winter School on Geometry and Physics (Srní 1990). Rend. Circ. Mat. Palermo 26 (1991) 197–206.

- [Müg03] M. Müger. From subfactors to categories and topology II: The quantum double of tensor categories and subfactors. *J. Pure Appl. Algebra* 180 (2003) 159-219.
- [Ost03] V. Ostrik. Module categories, weak Hopf algebras and modular invariants. *Transform. Groups* 8 (2003) 177–206.
- [Str96] R. Street. Categorical structures. *Handbook of algebra*, Vol. 1, 529–577. *Handb. Algebr.* 1. Elsevier/North-Holland, Amsterdam, 1996.
- [Szy] M. Szymik. Homotopy coherent centers versus centers of homotopy categories. <http://arxiv.org/abs/1305.3029>

Ehud Meir
Department
of Mathematical Sciences
University of Copenhagen
2100 Copenhagen Ø
DENMARK
meir@math.ku.dk

Markus Szymik
Department
of Mathematical Sciences
NTNU Norwegian University
of Science and Technology
7491 Trondheim
NORWAY
markus.szymik@math.ntnu.no

