

Markers of oxidative stress, skeletal muscle mass and function, and their responses to resistance exercise training in older adults

Ciriaco Carru^{a,b}, Mariasole Da Boit^c, Panagiotis Paliogiannis^a, Angelo Zinellu^a, Salvatore Sotgia^a, Rachael Sibson^d, Judith R. Meakin^e, Richard M. Aspden^d, Arduino A. Mangoni^f, Stuart R. Gray^g

^a Department of Biomedical Sciences, University of Sassari, Sassari, Italy; ^b Quality Control Unit, University Hospital (AOUSS), Sassari, Italy; ^c Faculty of Health and Life Sciences, De Montfort University, Leicester, United Kingdom; ^d Institute of Medical Sciences, University of Aberdeen, Aberdeen, United Kingdom; ^e Exeter MR Research Centre, University of Exeter, Exeter, United Kingdom; ^f Department of Clinical Pharmacology, College of Medicine and Public Health, Flinders University, Adelaide, Australia; ^g Institute of Cardiovascular and Medical Sciences, University of Glasgow, Glasgow, United Kingdom.

Funding: this work was supported by the Biotechnology and Biological Sciences Research Council, United Kingdom (grant number BB/J015911/1).

Corresponding author: Prof Ciriaco Carru, Department of Biomedical Sciences, University of 07100 Sassari, Sassari, Italy. Tel: +39079229775; Email: carru@uniss.it

Word count: 2,914

Abstract

Background: Oxidative stress (OS) negatively affects skeletal muscle homeostasis in experimental models of ageing. However, little is known about the associations between circulating OS markers and parameters of muscle mass and function, and their responses to exercise training, in humans.

Methods: Maximal voluntary contraction (MVC, primary outcome) and isokinetic torque of the knee extensors at $30^{\circ} \text{ s}^{-1}$ (MIT), muscle cross-sectional area (MCSA) and quality (MQ, secondary outcomes), and plasma concentrations of malondialdehyde (MDA, pro-OS), homocysteine (HCY, pro-OS), taurine (TAU, anti-OS), and protein sulfhydryl groups (PSH, anti-OS) were measured in 27 healthy older males and 23 females at baseline and after an 18-week resistance exercise program, with or without a nutritional intervention (fish oil vs. placebo).

Results: After adjusting for age, glomerular filtration rate, and nutritional intervention, there were no significant correlations between baseline OS markers and muscle parameters, barring a positive association between TAU and MIT in females ($r=0.53$, $P=0.035$) and between MDA and MCSA in males ($r=0.69$, $P=0.001$). Training did not significantly change OS markers, except for a reduction in MDA in females ($-0.27 \mu\text{mol/L}$, 95% CI -0.51 to -0.02 , $P=0.034$). In females, there were significant correlations between baseline MDA and exercise-induced changes in MVC ($P=0.018$), baseline TAU and changes in MCSA ($P=0.026$), and baseline HCY and changes in MCSA ($P=0.046$) and MQ ($P=0.022$). In males, baseline MDA was significantly associated with exercise-induced changes in MVC ($P=0.040$).

Conclusions: Plasma MDA, HCY, and TAU were significantly associated with baseline and/or exercise-induced changes in muscle mass and function in healthy older adults, primarily in females. Pending further confirmation in other populations, specific OS markers, particularly MDA, might predict muscle responses to resistance exercise programs in old age.

Key words: oxidative stress, muscle mass, muscle function, old age, exercise.

1. Introduction

Ageing is associated with intracellular and extracellular morphological and functional changes, resulting in the progressive impairment of organ function and the development of disease states. One typical example is the significant decline in neuromuscular function and performance associated with the reduction of skeletal muscle mass (sarcopenia), which results in impaired muscle strength and physical performance (Hunter et al., 2016, Cruz-Jentoft et al., 2010, Morley et al., 2011). The estimated prevalence of sarcopenia in old age ranges between 25-50% (Patel et al., 2013). The progressive increase in the number of frail older adults will inevitably lead to an increased prevalence of sarcopenia, and its associated health-care costs, worldwide (Muscaritoli et al., 2010, Visser and Schaap, 2011).

Many factors have been proposed to play a role in the pathogenesis of sarcopenia, including physical inactivity, alterations in protein metabolism and hormones, neurodegeneration, inflammation, and oxidative stress (OS) (Rolland et al., 2008). There is good evidence that the process of ageing is associated with increased concentrations of reactive oxygen and/or nitrogen species (RONS), and a reduction in endogenous antioxidants, in the skeletal muscle (Reid and Durham, 2002). This can lead to modifications of nucleic acids, proteins and lipids, resulting in molecular damage and/or dysfunction.

High concentrations of RONS have been shown to negatively affect muscle mass and function by altering the balance between protein synthesis and proteolysis (Fulle et al., 2004, Powers et al., 2011). However, the assessment of the effect of OS on skeletal muscle homeostasis has primarily been conducted in *in-vitro* studies and animal models, and the generalizability of these findings in humans is uncertain (Ji, 2015, Scicchitano et al., 2017). The investigation of circulating markers of OS in older adults might be particularly useful for the identification of those at risk of developing sarcopenia and/or responders to specific interventions, such as exercise training.

The aim of this study was to investigate the associations between predefined pro-OS and anti-OS markers and established parameters of skeletal muscle mass and function, and their responses to an exercise training and nutritional intervention program, in healthy older adults. Since the direct measurement of OS markers in tissue or body fluids is challenging, we measured plasma concentrations of oxidation target products such as malondialdehyde (MDA), low molecular thiol (HCY), and the antioxidants sulfhydryl group of proteins (PSH) and taurine (TAU).

2. Methods

2.1. Study population

We recruited 50 community-dwelling adults >65 years old (27 males and 23 females) that were not participating in any resistance exercise training, had no history of cardiovascular disease, cancer, arthritis, respiratory disease, metabolic disease, recent fractures and loss of mobility, and did not take regular analgesics or nutritional supplements. However, one female was prescribed angiotensin converting enzyme inhibitors for mild hypertension, and one male was prescribed allopurinol for gout. The study was approved by the University of Aberdeen College of Life Sciences and Medicine Ethics Review Board (CERB/2011/6/644) and registered at clinicaltrials.gov (ClinicalTrials.gov Identifier: NCT02843009). Written consent was obtained after explaining the aims, risks, and potential discomfort, in accordance with the declaration of Helsinki. Participants were part of a study investigating the effects of fish oil consumption on adaptations to resistance exercise, and were randomly assigned to either 3.0 g/day safflower oil or 3.0 g/day fish oil with data on other aspects of the project published elsewhere (Da Boit et al., 2017).

2.2. Resistance exercise training

Resistance exercise training was performed twice weekly for 18 weeks. Each session included four sets of nine repetitions for each exercise: leg press, leg extension, leg curl and calf press. The load for each exercise was set at 70% of the participant's one repetition maximum (1RM). This was assessed for each exercise at baseline and every six weeks, and the load adjusted accordingly.

2.3. Parameters of muscle mass and function

The following measurements were performed the morning after an overnight fast, as previously described (Da Boit et al., 2017). Measurements were made prior to and at least 48-h after the completion of the resistance exercise training intervention.

2.3.1. Knee extensor isometric and isokinetic torque

The maximal isometric torque of the knee extensor muscles of the right leg was determined during a maximal voluntary contraction (MVC) with the participant seated on a Biodex dynamometer with a knee angle of 73°. With the same seating position, maximal isokinetic torque (MIT) of the knee extensors was measured at 30° s⁻¹.

2.3.2. Magnetic resonance imaging (MRI)

Forty-five participants underwent MRI (3 others were claustrophobic and 2 had metal implants) on a Philips Achieva 3.0 Tesla whole body scanner using a 16-channel SENSE XL Torso coil. Muscle cross-section area (MCSA) was quantified mid-thigh. Muscle quality (MQ) was calculated as torque (knee extensor isometric strength) per unit MCSA.

2.4. Anthropometric and OS parameters

Body weight and height were measured, and body mass index (BMI) calculated, for each study participant at baseline. Blood samples were collected by venipuncture in K⁺ EDTA vacutainers, placed on ice and processed within 30 min. After samples were centrifuged for 10 min at 4 °C at 800g plasma was aliquoted and stored at -80°C until analysis. MDA was

determined by spectrophotometry according to an established method (Esterbauer and Cheeseman, 1990). Briefly, plasma was mixed with 0.075% thiobarbituric acid and incubated at 95 °C for 30 min. MDA concentration was assayed by measuring the absorbance at 535 nm using a MDA standard curve. Protein-SH (PSH) was assessed by spectrophotometry at 405 nm using 5,5', dithiobis-2-nitrobenzoic acid (Ellman, 1959). PSH concentration was determined using a glutathione standard curve and then normalized versus protein plasma quantity measured by the Lowry's method. The low molecular weight thiols HCY, and TAU were determined by capillary electrophoresis (CE) with Laser Induced Fluorescence detection (Carru et al., 2004, Zinellu et al., 2003, Zinellu et al., 2009). Following protein precipitation with trichloroacetic acid, samples were derivatized using 5-(Iodoacetamido)fluorescein as fluorophore for HCY, and with fluorescein isothiocyanate for TAU. Following dilution, samples were injected in CE. Plasma creatinine and creatine concentrations were measured using an Aquity UPLC coupled to a qToF Premier high-resolution mass spectrometer (Waters, Sydney, Australia). Estimated glomerular filtration rate (eGFR) was measured using the Modification of Diet in Renal Disease formula (Levey et al., 1999).

2.5. Statistical analysis

Results are expressed as means \pm SD, medians and interquartile ranges, or frequencies as appropriate. After testing for normal distribution, using the Kolmogorov-Smirnov test, between-group differences were assessed either by one-way ANOVA or Mann-Whitney U test. Differences between baseline and post-exercise muscle mass and function were assessed either by paired Student's *t*-test or Wilcoxon test. In each sex, the effects of exercise and nutritional intervention on OS markers were assessed by ANCOVA. Partial correlations, adjusted for age, eGFR, and nutritional intervention, assessed the relationship between baseline OS markers and parameters of skeletal muscle mass and function, and their changes after the exercise training (MVC: primary outcome; MIT, MCSA and MQ: secondary

outcomes). Analyses were performed using IBM SPSS Statistics Version 23, Release 23.0.0.2 (SPSS Inc., Armonk, NY, USA). A two-sided $P < 0.05$ indicated statistical significance.

3. Results

3.1. Baseline characteristics

MVC, MIT, MCSA, and plasma creatinine, HCY and PSH concentrations were significantly higher in males. By contrast, plasma creatine concentrations were significantly higher in females (Table 1). MQ and TAU and MDA concentrations were similar in both sexes.

3.2. Effect of exercise on muscle parameters and OS markers

The resistance exercise training program significantly increased MVC ($31 \pm 24\%$, $P < 0.001$), MIT ($12 \pm 18\%$, $P < 0.001$), MCSA ($3 \pm 5\%$, $P < 0.001$) and MQ ($27 \pm 26\%$, $P < 0.001$) in the whole group. There were no significant exercise-induced changes in OS markers, except for a significant reduction in MDA concentrations in females (Table 2).

3.3. Females

3.3.1. Baseline

There were no significant correlations between OS markers and parameters of muscle mass and function, barring a significant positive correlation between TAU and MIT (Table 3).

3.3.2. Exercise-induced changes

There was a significant positive correlation between baseline HCY and changes in MCSA, and a negative correlation between baseline HCY and changes in MQ (Table 4). Significant positive correlations were also observed between baseline TAU and changes in MCSA, and between baseline MDA and changes in MVC (Table 4).

3.4. Males

3.4.1. Baseline

There were no significant correlations between baseline OS markers and muscle parameters, barring a significant positive association between MDA and MCSA (Table 5).

3.4.2. Exercise-induced changes

There were no significant correlations between baseline OS markers and changes in muscle parameters, barring a significant positive association between MDA and changes in MVC (Table 6).

4. Discussion

Many efforts have been recently made to identify effective strategies to prevent and treat sarcopenia, including targeted pharmacological treatments, hormone replacement therapies, antioxidant-nutritional interventions, and physical exercise (Cruz-Jentoft et al., 2010, Morley et al., 2011). Among them, physical exercise has been shown to be particularly effective by increasing muscle strength, mass, and physical performance (Yoshimura et al., 2017). The identification of circulating biomarkers associated with parameters of muscle mass and function would be useful not only for sarcopenia risk stratification but also to predict the response to exercise training programs in older adults.

In this study, we investigated associations between predefined markers with either pro-OS or anti-OS activity and parameters of muscle structure and function, at baseline and after a resistance exercise training and nutritional intervention program, in a group of healthy older adults. At baseline, there were no significant correlations between OS markers and muscle parameters, except for a positive association between TAU and MIT in females and between MDA and MCSA in males. There were no significant exercise-induced changes in the concentrations of OS markers in either sex, except for a significant reduction in MDA in

females. In females, there were significant correlations between baseline MDA and exercise-induced changes in MVC, TAU and changes in MCSA, and between baseline HCY and changes in MCSA and MQ. In males, baseline MDA was significantly associated with exercise-induced changes in MVC. Therefore, the pro-OS MDA and HCY and the anti-OS TAU, but not PSH, showed significant associations with baseline and/or exercise-induced changes in skeletal muscle mass and function. In particular, baseline MDA concentrations predicted exercise-induced changes in MVC, the primary outcome, in both sexes.

MDA, the enzymatic and non-enzymatic product of prostaglandins breakdown, endoperoxide, and the end product of lipid peroxidation (Ayala et al., 2014), is useful for the monitoring of serum free radicals. The latter are difficult to measure directly because of their relatively low stability. Animal and human studies have investigated the effects of chronic (weeks to months) exercise on MDA concentrations. However, in these studies it was not possible to investigate the associations between plasma/serum MDA and skeletal muscle parameters because a) MDA concentrations were measured in specific tissues, and b) parameters of muscle mass and function were not assessed. Ravi Kiran et al. studied the effects of different intensities and durations of swim training sessions, six days/week for four weeks, on myocardial MDA concentrations in young and old rats. A significant exercise-induced reduction in myocardial MDA concentrations was observed in both groups (Ravi Kiran et al., 2006). Pangkahila et al. studied the effects of eight-week conventional training or balanced physical exercise on serum MDA, but not muscle parameters, in 24 males aged 18-30 years. MDA concentrations, measured before and after a training session at baseline and eight weeks later, were not significantly different over time and between the groups (Pangkahila et al., 2016). In our study, the exercise-associated reduction in plasma MDA concentrations was statistically significant in females but not in males. This suggests that resistance exercise training might have beneficial effects on lipid peroxidation in older individuals, particularly

women. However, the observed reduction in MDA concentrations might also be an indirect result of the improvement in the control of free radical production by the mitochondria (Daussin et al., 2012). Notably, the observed associations between baseline MDA concentrations and changes in MVC in both sexes support the potential role of MDA in the identification of those individuals that are more likely to respond, in terms of muscle strength, to resistance exercise training.

By contrast, the exercise program exerted opposite, albeit not significant, effects on plasma TAU concentrations in males (decreased) and females (increased). Baseline TAU concentrations were significantly correlated with baseline MIT and with exercise-induced changes in MSCA in females, but not in males. TAU plays an important role in several physiological functions, including cell signaling and development, membrane stability and receptor regulation, and various antioxidant and metabolic effects (Huxtable, 1992).

Furthermore, it regulates the release of Ca^{2+} from the sarcoplasmic reticulum and maintains the sensitivity of contractile elements to Ca^{2+} , affecting directly the excitation-contraction process and skeletal muscle strength (Spriet and Whitfield, 2015). Animal and human studies have previously investigated the effects of TAU supplementation on skeletal muscle mass and function and the effects of acute exercise on TAU concentrations, however no study has sought to determine the effects of chronic exercise on TAU concentrations in humans. In our study, 18-weeks of resistance exercise training significantly increased all parameters of skeletal mass and function in males, despite the trend towards a reduction in plasma TAU concentrations. This suggests that TAU concentrations are unlikely to play a significant role in the observed exercise-induced changes in muscle parameters in this group. By contrast, the significant correlations between baseline TAU and exercise-induced changes in MSCA in females suggest, similarly to MDA, a potential role of this sulfur-containing β -amino acid in the identification of female responders to exercise training programs.

Resistance exercise training did not significantly change plasma HCY concentrations in both sexes. However, baseline plasma HCY concentrations were positively associated with exercise-induced changes in MCSA, and negatively associated with changes in MQ, in females but not in males. Human studies have previously investigated the associations between HCY concentrations, physical exercise, and muscle strength. Steenge et al reported a non-significant reduction in plasma HCY concentrations ($-0.6 \mu\text{mol/L}$, 95% CI -1.9 to 0.5) in healthy women aged 19-38 years undergoing an eight-week period of resistance exercise, with three sessions each week involving leg squats, leg presses, leg curls, leg rowing, leg extensions, shoulder presses, sit-ups, and step-up exercises (Steenge et al., 2001). In a cross-sectional study, Kuo et al observed a significant negative association between plasma HCY concentrations and peak quadriceps strength in 1,677 older adults from the National Health and Nutrition Examination Survey 1999-2002 (Kuo et al., 2007). In another cross-sectional study of 2,919 older adults with hyperhomocysteinemia, Swart et al reported that plasma HCY concentrations were not independently associated with skeletal muscle mass, however they were independently associated with reduced handgrip strength in women ($P=0.02$) but not in men (Swart et al., 2013a). By contrast, in a study of 1,301 older adults from the Longitudinal Aging Study in Amsterdam, Swart et al observed an independent negative association between plasma HCY concentrations and reduced handgrip strength in men ($P<0.01$) but not in women. However, HCY concentrations did not predict the changes in handgrip strength after a three-year follow up (Swart et al., 2013b). We did not observe any significant correlation between plasma HCY concentrations and baseline and exercise-induced changes in parameters of muscle strength, MVC and MIT, in both sexes. However, similarly to TAU, the correlations with changes in MCSA and MQ in females warrant further research to clarify the potential predictive role of this highly reactive and pro-oxidant amino acid in subjects undergoing exercise training programs.

Although baseline PSH concentrations were significantly higher in males than in females, we did not observe any significant exercise-induced changes in PSH concentrations, or associations between PSH and parameters of muscle mass and function. In accordance to our findings, Margaritis et al did not observe any significant change in PSH concentrations in five healthy young males undergoing a six-month training program consisting of 29 training sessions, performed every three days, involving two types of exercise, the squat and the calf press. No specific parameters of muscle mass and function were assessed (Margaritis et al., 2009). Pending further confirmation in longitudinal studies, our results do not suggest a significant role of PSH in terms of sarcopenia risk stratification and identification of responders to exercise training programs, based on changes in muscle strength, volume, and quality.

The observation of sex-related differences in the markers studied, particularly TAU and HCY, suggests that the relationship between antioxidant and lipid peroxidation pathways and skeletal muscle homeostasis may be sex-specific. This, in turn, might help in the identification of tailored exercise programs in older males and females. However, the capacity of MDA to predict exercise-induced changes in MVC in both sexes suggests that this OS marker can predict treatment responders and that therapies targeting MDA and specific OS pathways might provide salutary effects on skeletal muscle homeostasis in old age.

Our study has some limitations, including its relatively small sample size and the recruitment of participants without significant medical history. Therefore, it is unknown whether our results can be generalized to other older populations, particularly frail patients with sarcopenia. Nevertheless, it also has significant strengths, including the rigorous assessment of established pro- and anti-OS markers and established parameters of muscle mass and function. Furthermore, we ruled out the potential confounding effect of co-morbidities and

medications that can affect skeletal muscle structure and function, and closely supervised the exercise training program in all participants.

In conclusion, the pro-oxidant markers MDA and HCY and the anti-oxidant marker TAU showed significant associations with baseline and/or exercise-induced changes in skeletal muscle mass and function in healthy older adults, particularly in females. Pending further confirmation in other study populations, these markers might be useful for sarcopenia risk stratification and prediction of responses to resistance exercise programs.

Table 1. Baseline characteristics

Parameter	Whole group (n=50)	Males (n=27)	Females (n=23)	P-value*
Age (years)	70 [67, 73]	69 [67, 72]	71 [68, 73]	0.544
Height (cm)	169 [159, 176]	175 [172, 179]	159 [155, 168]	<0.0001
Weight (Kg)	72.1±15.1	78.1±15.0	65.1±12.3	0.002
BMI (Kg/m ²)	25.4±4.4	25.6±4.2	25.3±4.7	0.821
eGFR (mL/min)	80±12	77±12	83±13	0.079
Creatine (µmol/L)	168±76	134±51	208±83	0.001
Creatinine (µmol/L)	79±12	85±10	71±9	<0.0001
MVC (N•m)	90 [75, 120]	115 [94, 138]	76 [68, 85]	<0.0001
MIT (N•m)	116±37	141±32	86±14	<0.0001
MCSA (cm ²)	51 [38, 64]	63 [58, 68]	37 [32, 43]	<0.0001
MQ (N•m/cm ²)	2.0±0.5	1.9±0.5	2.1±0.5	0.293
HCY (µmol/L)	13.6±3.7	14.8±4.2	12.4±2.6	0.031
TAU (µmol/L)	57.7±20.3	61.8±25.2	53.1±11.8	0.156
MDA (µmol/L)	2.78±0.59	2.93±0.53	2.61±0.63	0.067
PSH (µmol/L)	5.47±0.82	5.88±0.77	5.03±0.64	0.0001

Data shown as mean ± SD or median [IQR] as appropriate

BMI, body mass index; eGFR, estimated glomerular filtration rate; MVC, maximal voluntary contraction; MIT, maximal isokinetic torque; MCSA, muscle volume; MQ, muscle quality; HCY, homocysteine; TAU, taurine; MDA, malondialdehyde; PSH, sulphhydryl group of protein.

Significant differences between males and females (P<0.05) are highlighted in bold.

Table 2. Baseline vs. post-exercise mean (95% CI) differences in oxidative stress markers in females and males

	Mean (95% CI) difference	P-value
Females		
HCY	+0.15 (-0.57 to 0.86)	0.673#
TAU	+4.24 (-1.67 to 10.16)	0.149
MDA	-0.27 (-0.51 to -0.02)	0.034
PSH	-0.19 (-0.41 to 0.03)	0.094
Males		
HCY	+0.25 (-0.75 to 1.25)	0.613
TAU	-7.06 (-16.08 to 1.95)	0.119
MDA	-0.09 (-0.28 to 0.11)	0.356
PSH	+0.07 (-0.21 to 0.36)	0.600

HCY, homocysteine; TAU, taurine; MDA, malondialdehyde; PSH, sulphhydryl group of protein.

Significant differences ($P < 0.05$) are highlighted in bold.

#, time * nutritional intervention interaction ($P < 0.05$)

Table 3. Partial correlations* between baseline oxidative stress markers and parameters of muscle mass and function in females

	MVC (N•m)	MIT (N•m)	MCSA (cm²)	MQ (N•m.cm²)
HCY	r= +0.143 P=0.597	r= +0.004 P=0.988	r= -0.447 P=0.082	r= +0.476 P=0.062
TAU	r= +0.140 P=0.58	r= +0.53 P=0.035	r= +0.093 P=0.732	r= +0.073 P=0.789
MDA	r= -0.448 P=0.081	r= +0.003 P=0.99	r= +0.021 P=0.939	r= -0.348 P=0.186
PSH	r= +0.101 P=0.711	r= +0.384 P=0.142	r= +0.1 P=0.713	r= +0.055 P=0.841

*, adjusted for age and estimated glomerular filtration rate

MVC, maximal voluntary contraction; MIT, maximal isokinetic torque; MCSA, muscle volume; MQ, muscle quality; HCY, homocysteine; TAU, taurine; MDA, malondialdehyde; PSH, sulphydril group of protein.

Significant correlations (P<0.05) are highlighted in bold.

Table 4. Partial correlations* between oxidative stress markers and percent exercise-induced changes in parameters of muscle mass and function in females

	$\Delta\%MVC$ (N•m)	$\Delta\%MIT$ (N•m)	$\Delta\%MCSA$ (cm ²)	$\Delta\%MQ$ (N•m.cm ⁻²)
HCY	r= -0.155 P=0.582	r= +0.066 P=0.816	r= +0.522 P=0.046	r= -0.585 P=0.022
TAU	r= +0.278 P=0.316	r= -0.112 P=0.692	r= +0.573 P=0.026	r= -0.009 P=0.974
MDA	r= +0.6 P=0.018	r= -0.105 P=0.709	r= +0.036 P=0.898	r= +0.093 P=0.741
PSH	r= +0.198 P=0.480	r= -0.258 P=0.353	r= +0.355 P=0.194	r= -0.404 P=0.135

*, adjusted for age, estimated glomerular filtration rate, and nutritional intervention

MVC, maximal voluntary contraction; MIT, maximal isokinetic torque; MCSA, muscle volume; MQ, muscle quality; HCY, homocysteine; TAU, taurine; MDA, malondialdehyde; PSH, sulphhydryl group of protein.

Significant correlations (P<0.05) are highlighted in bold.

Table 5. Partial correlations* between baseline oxidative stress markers and parameters of muscle mass and function in males

	MVC (N•m)	MIT (N•m)	MCSA (cm²)	MQ (N•m.cm⁻²)
HCY	r= +0.208 P=0.38	r= +0.151 P=0.526	r= +0.303 P=0.194	r= +0.72 P=0.762
TAU	r= -0.111 P=0.641	r= +0.316 P=0.175	r= -0.407 P=0.075	r= +0.321 P=0.167
MDA	r= +0.221 P=0.349	r= +0.293 P=0.211	r= +0.699 P=0.001	r= -0,122 P=0.608
PSH	r= -0.179 P=0.451	r= -0.130 P=0.585	r= +0.341 P=0.141	r= -0.367 P=0.111

*, adjusted for age and estimated glomerular filtration rate

MVC, maximal voluntary contraction; MIT, maximal isokinetic torque; MCSA, muscle volume; MQ, muscle quality; HCY, homocysteine; TAU, taurine; MDA, malondialdehyde; PSH, sulphydril group of protein.

Significant correlations (P<0.05) are highlighted in bold.

Table 6. Partial correlations* between oxidative stress markers and percent exercise-induced changes in parameters of muscle mass and function in males

	$\Delta\%MVC$ (N•m)	$\Delta\%MIT$ (N•m)	$\Delta\%MCSA$ (cm ²)	$\Delta\%MQ$ (N•m.cm ⁻²)
HCY	r= +0.387 P=0.101	r= +0.303 P=0.208	r= +0.113 P=0.645	r= +0.341 P=0.153
TAU	r= -0.043 P=0.86	r= -0.026 P=0.916	r= +0.388 P=0.101	r= -0.13 P=0.595
MDA	r= +0.475 P=0.04	r= +0.226 P=0.353	r= +0.019 P=0.937	r= +0.44 P=0.06
PSH	r= -0.93 P=0.705	r= -0.187 P=0.444	r= -0.184 P=0.451	r= -0.46 P=0.852

*, adjusted for age, estimated glomerular filtration rate, and nutritional intervention

MVC, maximal voluntary contraction; MIT, maximal isokinetic torque; MCSA, muscle volume; MQ, muscle quality; HCY, homocysteine; TAU, taurine; MDA, malondialdehyde; PSH, sulphydril group of protein.

Significant correlations (P<0.05) are highlighted in bold.

References

- Ayala, A., Munoz, M. F. & Arguelles, S. 2014. Lipid peroxidation: production, metabolism, and signaling mechanisms of malondialdehyde and 4-hydroxy-2-nonenal. *Oxid Med Cell Longev*, 2014, 360438. 10.1155/2014/360438
- Carru, C., Deiana, L., Sotgia, S., Pes, G. M. & Zinellu, A. 2004. Plasma thiols redox status by laser-induced fluorescence capillary electrophoresis. *Electrophoresis*, 25, 882-9. 10.1002/elps.200305768
- Cruz-Jentoft, A. J., Baeyens, J. P., Bauer, J. M., Boirie, Y., Cederholm, T., Landi, F., Martin, F. C., Michel, J. P., Rolland, Y., Schneider, S. M., Topinkova, E., Vandewoude, M., Zamboni, M. & European Working Group on Sarcopenia in Older, P. 2010. Sarcopenia: European consensus on definition and diagnosis: Report of the European Working Group on Sarcopenia in Older People. *Age Ageing*, 39, 412-23. 10.1093/ageing/afq034
- Da Boit, M., Sibson, R., Sivasubramaniam, S., Meakin, J. R., Greig, C. A., Aspden, R. M., Thies, F., Jeromson, S., Hamilton, D. L., Speakman, J. R., Hambly, C., Mangoni, A. A., Preston, T. & Gray, S. R. 2017. Sex differences in the effect of fish-oil supplementation on the adaptive response to resistance exercise training in older people: a randomized controlled trial. *Am J Clin Nutr*, 105, 151-158. 10.3945/ajcn.116.140780
- Daussin, F. N., Rasseneur, L., Bouitbir, J., Charles, A. L., Dufour, S. P., Geny, B., Burelle, Y. & Richard, R. 2012. Different timing of changes in mitochondrial functions following endurance training. *Med Sci Sports Exerc*, 44, 217-24. 10.1249/MSS.0b013e31822b0bd4
- Ellman, G. L. 1959. Tissue sulfhydryl groups. *Arch Biochem Biophys*, 82, 70-7.

- Esterbauer, H. & Cheeseman, K. H. 1990. Determination of aldehydic lipid peroxidation products: malonaldehyde and 4-hydroxynonenal. *Methods Enzymol*, 186, 407-21.
- Fulle, S., Protasi, F., Di Tano, G., Pietrangelo, T., Beltramin, A., Boncompagni, S., Vecchiet, L. & Fano, G. 2004. The contribution of reactive oxygen species to sarcopenia and muscle ageing. *Exp Gerontol*, 39, 17-24.
- Hunter, S. K., Pereira, H. M. & Keenan, K. G. 2016. The aging neuromuscular system and motor performance. *J Appl Physiol* (1985), 121, 982-995. 10.1152/jappphysiol.00475.2016
- Huxtable, R. J. 1992. Physiological actions of taurine. *Physiol Rev*, 72, 101-63.
- Ji, L. L. 2015. Redox signaling in skeletal muscle: role of aging and exercise. *Adv Physiol Educ*, 39, 352-9. 10.1152/advan.00106.2014
- Kuo, H. K., Liao, K. C., Leveille, S. G., Bean, J. F., Yen, C. J., Chen, J. H., Yu, Y. H. & Tai, T. Y. 2007. Relationship of homocysteine levels to quadriceps strength, gait speed, and late-life disability in older adults. *J Gerontol A Biol Sci Med Sci*, 62, 434-9.
- Levey, A. S., Bosch, J. P., Lewis, J. B., Greene, T., Rogers, N. & Roth, D. 1999. A more accurate method to estimate glomerular filtration rate from serum creatinine: a new prediction equation. Modification of Diet in Renal Disease Study Group. *Ann Intern Med*, 130, 461-70.
- Margaritis, I., Rousseau, A. S., Marini, J. F. & Chopard, A. 2009. Does antioxidant system adaptive response alleviate related oxidative damage with long term bed rest? *Clin Biochem*, 42, 371-9. 10.1016/j.clinbiochem.2008.10.026
- Morley, J. E., Abbatecola, A. M., Argiles, J. M., Baracos, V., Bauer, J., Bhasin, S., Cederholm, T., Coats, A. J., Cummings, S. R., Evans, W. J., Fearon, K., Ferrucci, L., Fielding, R. A., Guralnik, J. M., Harris, T. B., Inui, A., Kalantar-Zadeh, K., Kirwan, B. A., Mantovani, G., Muscaritoli, M., Newman, A. B., Rossi-Fanelli, F., Rosano, G.

- M., Roubenoff, R., Schambelan, M., Sokol, G. H., Storer, T. W., Vellas, B., von Haehling, S., Yeh, S. S., Anker, S. D., Society on Sarcopenia, C. & Wasting Disorders Trialist, W. 2011. Sarcopenia with limited mobility: an international consensus. *J Am Med Dir Assoc*, 12, 403-9. 10.1016/j.jamda.2011.04.014
- Muscaritoli, M., Anker, S. D., Argiles, J., Aversa, Z., Bauer, J. M., Biolo, G., Boirie, Y., Bosaeus, I., Cederholm, T., Costelli, P., Fearon, K. C., Laviano, A., Maggio, M., Rossi Fanelli, F., Schneider, S. M., Schols, A. & Sieber, C. C. 2010. Consensus definition of sarcopenia, cachexia and pre-cachexia: joint document elaborated by Special Interest Groups (SIG) "cachexia-anorexia in chronic wasting diseases" and "nutrition in geriatrics". *Clin Nutr*, 29, 154-9. 10.1016/j.clnu.2009.12.004
- Pangkahila, E. A., Adiputra, N., Pangkahila, W. & Yasa, I. W. P. S. 2016. Balanced Physical Exercise Increase Physical Fitness, Optimize Endorphin Levels, and Decrease Malondialdehyde Levels. *Bali Medical Journal*, 5. 10.15562/bmj.v5i3.337
- Patel, H. P., Syddall, H. E., Jameson, K., Robinson, S., Denison, H., Roberts, H. C., Edwards, M., Dennison, E., Cooper, C. & Aihie Sayer, A. 2013. Prevalence of sarcopenia in community-dwelling older people in the UK using the European Working Group on Sarcopenia in Older People (EWGSOP) definition: findings from the Hertfordshire Cohort Study (HCS). *Age Ageing*, 42, 378-84. 10.1093/ageing/afs197
- Powers, S. K., Smuder, A. J. & Criswell, D. S. 2011. Mechanistic links between oxidative stress and disuse muscle atrophy. *Antioxid Redox Signal*, 15, 2519-28. 10.1089/ars.2011.3973
- Ravi Kiran, T., Subramanyam, M. V., Prathima, S. & Asha Devi, S. 2006. Blood lipid profile and myocardial superoxide dismutase in swim-trained young and middle-aged rats: comparison between left and right ventricular adaptations to oxidative stress. *J Comp Physiol B*, 176, 749-62. 10.1007/s00360-006-0096-5

- Reid, M. B. & Durham, W. J. 2002. Generation of reactive oxygen and nitrogen species in contracting skeletal muscle: potential impact on aging. *Ann N Y Acad Sci*, 959, 108-16.
- Rolland, Y., Czerwinski, S., Abellan Van Kan, G., Morley, J. E., Cesari, M., Onder, G., Woo, J., Baumgartner, R., Pillard, F., Boirie, Y., Chumlea, W. M. & Vellas, B. 2008. Sarcopenia: its assessment, etiology, pathogenesis, consequences and future perspectives. *J Nutr Health Aging*, 12, 433-50.
- Scicchitano, B. M., Pelosi, L., Sica, G. & Musaro, A. 2017. The physiopathologic role of oxidative stress in skeletal muscle. *Mech Ageing Dev*. 10.1016/j.mad.2017.08.009
- Spriet, L. L. & Whitfield, J. 2015. Taurine and skeletal muscle function. *Curr Opin Clin Nutr Metab Care*, 18, 96-101. 10.1097/MCO.0000000000000135
- Steenge, G. R., Verhoef, P. & Greenhaff, P. L. 2001. The effect of creatine and resistance training on plasma homocysteine concentration in healthy volunteers. *Arch Intern Med*, 161, 1455-6.
- Swart, K. M., Enneman, A. W., van Wijngaarden, J. P., van Dijk, S. C., Brouwer-Brolsma, E. M., Ham, A. C., Dhonukshe-Rutten, R. A., van der Velde, N., Brug, J., van Meurs, J. B., de Groot, L. C., Uitterlinden, A. G., Lips, P. & van Schoor, N. M. 2013a. Homocysteine and the methylenetetrahydrofolate reductase 677C-->T polymorphism in relation to muscle mass and strength, physical performance and postural sway. *Eur J Clin Nutr*, 67, 743-8. 10.1038/ejcn.2013.97
- Swart, K. M., van Schoor, N. M., Heymans, M. W., Schaap, L. A., den Heijer, M. & Lips, P. 2013b. Elevated homocysteine levels are associated with low muscle strength and functional limitations in older persons. *J Nutr Health Aging*, 17, 578-84. 10.1007/s12603-013-0047-2

- Visser, M. & Schaap, L. A. 2011. Consequences of sarcopenia. *Clin Geriatr Med*, 27, 387-99.
10.1016/j.cger.2011.03.006
- Yoshimura, Y., Wakabayashi, H., Yamada, M., Kim, H., Harada, A. & Arai, H. 2017. Interventions for Treating Sarcopenia: A Systematic Review and Meta-Analysis of Randomized Controlled Studies. *J Am Med Dir Assoc*, 18, 553 e1-553 e16.
10.1016/j.jamda.2017.03.019
- Zinellu, A., Carru, C., Galistu, F., Usai, M. F., Pes, G. M., Baggio, G., Federici, G. & Deiana, L. 2003. N-methyl-D-glucamine improves the laser-induced fluorescence capillary electrophoresis performance in the total plasma thiols measurement. *Electrophoresis*, 24, 2796-804. 10.1002/elps.200305570
- Zinellu, A., Sotgia, S., Scanu, B., Chessa, R., Gaspa, L., Franconi, F., Deiana, L. & Carru, C. 2009. Taurine determination by capillary electrophoresis with laser-induced fluorescence detection: from clinical field to quality food applications. *Amino Acids*, 36, 35-41. 10.1007/s00726-007-0022-5