

The effects of plant-based diets on pancreatic beta-cell function: A systematic review

Robert K Janko, Philip Wilson, Chinedu Nworie

Plant-based diets have been associated with improvements in blood glucose and HbA_{1c} levels, as well as improvements in insulin resistance. This review aimed to systematically analyse the scientific literature on the effects of plant-based dietary interventions on pancreatic beta-cell function, as measured by insulin secretion. The three clinical trials that were included in the analysis showed a high risk of bias that was predominantly due to a lack of blinding in their designs. Compared with non-plant-based interventions, vegan and vegetarian diets showed a number of significant improvements in insulin secretion, HbA_{1c} and weight loss across the studies. Larger randomised controlled trials with uniform measurements of beta-cell function would be beneficial to allow for quantitative analysis of pooled results.

Vegetarian and vegan diets have been associated with several health benefits, including lower risk of certain types of cancer and cardiovascular disease, lower blood lipid levels and lower blood glucose levels, as well as better management of diabetes overall (Dinu et al, 2017). A number of epidemiological studies have shown that high consumers of fruits, vegetables and fibre have a reduced risk of type 2 diabetes (InterAct Consortium, 2015; Dinu et al, 2017). A 2009 analysis of the Adventist Health Study-2 revealed that the prevalence of type 2 diabetes amongst vegans was 2.9% as opposed to 7.6% among non-vegetarians (Tonstad et al, 2009). This was partly due to the significantly lower BMI in vegans; however, the association remained significant even after adjusting for BMI.

Insulin resistance and impaired insulin production are the driving force in the development of type 2 diabetes. Higher-fat diets have been shown to increase the accumulation of intramyocellular lipids, making people susceptible to insulin resistance (Sakurai et al, 2011). In addition to

their lower fat content, plant-based diets offer protection against type 2 diabetes due to their high antioxidant, polyphenol, fibre and micronutrient composition. A systematic review identified 111 plants that have the ability to improve insulin sensitivity (Eddouks et al, 2014). Accumulation of fat in the cells of the pancreas can have detrimental effects as it can lead to impaired beta-cell function (Heiskanen et al, 2018). Exercise is a known contributor of improved beta-cell function; however, diet may also exert beneficial effects on pancreatic beta-cells by reducing pancreatic fat mass, lowering hyperglycaemia and reducing hyperlipidaemia (Larson-Meyer et al, 2006).

The production of insulin by beta-cells is also impaired by the presence and accumulation of advanced glycation end-products (AGEs) which are produced from precursor molecules such as methylglyoxal. Methylglyoxal has been shown to induce oxidative stress and cell death in pancreatic cells (Su et al, 2017). AGEs are formed endogenously in the body; however, the preparation of animal-derived products such as meat and cheese

Citation: Janko RK, Wilson P, Nworie C (2021) The effects of plant-based diets on pancreatic beta-cell function: A systematic review. *Journal of Diabetes Nursing* 25: [early view publication]

Article points

1. Plant-based diets have previously been associated with a lower incidence of type 2 diabetes even after adjustment for BMI.
2. There have been few randomised controlled trials into the effects of plant-based diets on pancreatic beta-cell function.
3. This systematic review demonstrates significant improvements in insulin secretion, HbA_{1c} and weight loss with plant-based diets; however, the quality of evidence is low, and larger randomised controlled trials with uniform measurements of beta-cell function would be beneficial.

Key words

- Dietary interventions
- Obesity, diet and lifestyle
- Plant-based diets
- Systematic review

Authors

Robert Janko and Chinedu Nworie are Master's students; Philip Wilson is Researcher and physician, all at University of Aberdeen.



Read more
online

**The ADA nutrition
therapy consensus report:
A quick guide**

A condensed summary of the American Diabetes Association's expert consensus report on the role of nutrition in the management of diabetes and the prevention of type 2 diabetes, including the main dietary messages for healthcare professionals.

Journal of Diabetes Nursing
23: JDN103

[Click here to access](#)

with the use of high heat results in the formation of AGEs in concentrations which are much higher than in plant-based foods (Uribarri et al, 2010). Furthermore, insulin resistance increases the secretory demand on beta-cells, which results in a decline in beta-cell function (Cerf, 2013). Therefore, dietary interventions that are low in animal fat and contain high amounts of fibre and antioxidants may lead to an improvement in the regulation of fat and glucose homeostasis, thereby preserving or improving beta-cell function.

Animal and *in vitro* studies have shown that factors such as caloric restriction (fasting) and plant chemicals such as flavonoids may lead to the preservation and regeneration of islet beta-cells, and thus improvement or restoration of insulin synthesis (Cheng et al, 2017; Ghorbani et al, 2019). However, the number of human studies involving plant-based dietary interventions are limited, and there have not been any systematic reviews analysing these studies.

Collectively, there is sufficient evidence to suggest that plant-based diets may directly impact the insulin-producing beta-cells of the pancreas. There are several systematic reviews and meta-analyses evaluating the efficacy of vegan or vegetarian diets in the treatment or prevention of type 2 diabetes; however, none of the reviews directly focus on the effect of plant-based diets on beta-cell function. Therefore, this study was conducted to evaluate these effects.

Methods

This systematic review was registered on PROSPERO (CRD42020168191) and has been conducted in line with the Preferred Reporting Items for Systematic Review and Meta-Analyses (PRISMA) statement.

A comprehensive and systematic literature search was conducted in an effort to identify relevant randomised controlled trials or interventional studies. A search strategy was devised using the PICO method for search term generation, which was used to search the Medline, EMBASE and Scopus databases. The key words used in combination as MeSH terms and text words were as follows: "Obesity" OR "obese" OR "Diabetes Mellitus" AND "plant based" OR "plant-based" OR "plant based" OR "plant food" OR "plant food" OR "vegan" OR "vegetarian" AND "beta

cell function" AND "beta-cell function" AND "b-cell function" AND "insulin secretion" AND "pancreatic function" AND "β cells". The systematic search was limited to human studies.

The references of included studies were also manually searched for potentially relevant articles.

Inclusion/exclusion criteria

Studies were included if they met the following inclusion criteria:

- Study designs had to be randomised controlled trials or non-randomised dietary intervention trials.
- Studies had to have a plant-based or vegan diet intervention group.
- Studies had to report C-peptide or insulin concentrations at baseline and post-intervention.

Studies were excluded if they were not published in a peer-reviewed journal. Studies that had enrolled participants with type 1 diabetes, were either ongoing or unpublished, did not include a plant-based intervention or were published in a non-English language were also excluded.

Data extraction

Two reviewers utilised a data extraction form adopted for this systematic review to extract all relevant data from the articles that met the inclusion criteria. The data extraction form was adopted from the SIGN (2015) data extraction form.

The systematic search of the literature was carried out by RKJ, who also screened relevant abstracts and titles, and completed data extraction. A second author, CN, was involved in data extraction. Any disagreement during data extraction was resolved by consensus.

The following data were extracted from relevant articles: author and publication date, country, participant characteristics relating to baseline health status (i.e. presence or absence of obesity or type 2 diabetes), sample size, intervention diet, control diet, measure of beta-cell function, study duration and length of follow-up, attrition rate during follow-up, method of randomisation and blinding, study design, funding source and conflict of interest.

Risk of bias assessment

Individual randomised controlled trials were assessed for risk of bias using the RevMan 5.3

software, which is based on the Cochrane Risk of Bias tool. Studies were judged for risk of bias with regard to the following criteria: random sequence generation (selection bias), allocation concealment (selection bias), blinding of participants and personnel (performance bias), incomplete outcome data (attrition bias), selective reporting (reporting bias), and other bias, which included sponsorship bias.

Results

The initial systematic search of the literature yielded 1131 titles and abstracts. Of these, 24 papers were read in full and three have been included in this systematic review (Figure 1).

A publication by Lim et al (2011) was excluded from this systematic review upon receipt of clarification via email from the original authors, who confirmed that the intervention diet contained meat protein. The design of this study was non-randomised and there was no control group.

The main results and characteristics of the included studies are summarised in Table 1.

All three of the included trials had the same lead author, and one of the studies only investigated the effect of a single meal on pancreatic beta-cell function, meaning that the intervention period differed significantly between the three trials. Therefore, it was decided that a quantitative synthesis in the form of meta-analysis would not be appropriate.

Among the three studies included, all had a randomised controlled trial design (Kahleova et al, 2011; 2018; 2019). The sample sizes of the individual studies ranged between 20 and 75 participants. The participants comprised adults with type 2 diabetes in two of the trials (Kahleova et al, 2011; 2019) and overweight adults in the other (Kahleova et al, 2018). All three trials controlled for caloric intake, and the two longer-term studies utilised calorie-restricted diets in both groups. Kahleova et al (2011) reduced caloric intakes in both groups by 500 kcal/day based on the measurement of resting energy expenditure of each subject.

The study by Kahleova et al (2019) involved two meal tests with a 1-week washout period between the meals, during which participants were asked to consume their regular diets. The other two trials were 16 and 24 weeks in duration.




Figure 1. PRISMA flow diagram of included studies.

Kahleova et al (2019) aimed to investigate if a plant-based meal stimulated postprandial incretin and insulin release. Kahleova et al (2018) aimed to test the effect of a dietary intervention on beta-cell function and insulin resistance in overweight adults with no history of diabetes. The third study hypothesised that a plant-based diet would be more effective in improving insulin resistance than a conventional diet for people with diabetes (Kahleova et al, 2011). Even though the main study objective concerned insulin resistance, it reported parameters relating to beta-cell function, such as fasting plasma C-peptide concentration.

Among people with type 2 diabetes in the trial by Kahleova et al (2011), 43% in the plant-based group were required to reduce their diabetes medication, compared with 5% in the control group ($P < 0.001$). This was done by a study physician in cases of repeat hypoglycaemia.

Kahleova et al (2019) examined the effects of a vegetarian burger versus a meat burger, matched for energy and macronutrient content, on postprandial incretin and insulin secretion. This was a randomised crossover trial that only recruited men with type 2 diabetes ($n = 20$).

Table 1. Results and characteristics of included studies.

Study reference and country of origin	Participant characteristics	Overall sample size	Intervention versus control diet	Measure of beta-cell function	Daily caloric intake	Study design and duration	Attrition during follow-up	Main results
Kahleova et al (2019) Czech Republic	Adults with type 2 diabetes	20	Plant-based tofu burger versus processed meat burger	C-peptide by deconvolution	Intervention 514.9 kcal/day Control 513.6 kcal/day	Randomised crossover study 1 day	All participants completed the study	Total insulin secretion (nmol/m²) Intervention: 68.2 (SD 66.1–70.4) Control: 62.2 (SD 60.2–64.3) Insulin secretion at fixed blood glucose of 5 mmol/L (pmol/min/m²) Intervention: 110.3 (SD 95.2–126.1) Control: 99.0 (SD 84.3–114.3)
Kahleova et al (2018) USA	Overweight adults	75	Low-fat plant-based diet (n=38) versus no dietary change (n=37)	Fasting plasma C-peptide	Intervention Baseline: 1851 kcal/day (SD 1695–2007) 16 weeks: 1450 kcal/day (SD 1249–1652) Control Baseline: 1923 kcal/day (SD 1627–2219) 16 weeks: 1582 kcal/day (SD 1368–1795)	Randomised controlled trial 16 weeks	Intervention 1 discontinued Control 2 discontinued	Change in fasting plasma C-peptide Intervention: Decrease from 2.6 ng/mL (SD 2.3–2.9) to 2.1 ng/mL (SD 1.9–2.4) Control: Increase from 2.5 ng/mL (SD 2.2–2.8) to 3.0 ng/mL (SD 2.2–3.8) Beta-cell glucose sensitivity (pmol/min/m² per mmol/L blood glucose) Intervention: Increase from 108.5 (SD 80.8–142.4) to 213.8 (SD 167.9–268.3) Control: Increase from 107.5 (SD 76.4–146.5) to 181.1 (SD 135.3–237.2)
Kahleova et al (2011) Czech Republic	Adults with type 2 diabetes	74	Vegetarian diet versus conventional diabetes diet	Plasma C-peptide levels by deconvolution	Both diets calorie-restricted by 500 kcal/day according to individual resting energy expenditure	Randomised controlled trial 24 weeks	84% of participants in each group completed all 24 weeks	Fasting plasma C-peptide change from baseline to 24 weeks (mIU/L) Intervention: -0.18 (±0.46) Control: 0 (±1.08)

SD=standard deviation.

The study's primary outcome was the area under the curve for postprandial insulin secretion. The vegetarian meal stimulated increased secretion of immunoreactive insulin (by 30.5%) and C-peptide (by 7.1%) compared with the meat meal ($P<0.001$ for both comparisons). Ingestion of the vegetarian meal resulted in significantly increased total insulin secretion 68.2 nmol/m^2 vs 62.2 nmol/m^2). Insulin secretion at 5 mmol/L glucose was 11.4% higher after the vegetarian meal compared with the meat meal ($P=0.04$). In addition, rate sensitivity increased by 55% ($P=0.04$), and the potentiation factor by 13.3% ($P=0.02$) after the vegetarian meal compared with the meat meal.

Kahleova et al (2018) observed a decrease in basal insulin secretion in the intervention group at 16 weeks (treatment effect $-54.2 \text{ pmol/min/m}^2$). The intervention group also had increased beta-cell glucose sensitivity; however, this did not reach statistical significance compared with the control group. The researchers found a dose-response increase in insulin secretion as a function of plasma glucose concentrations in the intervention group compared with controls. Changes in beta-cell function correlated negatively with changes in BMI ($r=-0.25$; $P=0.04$), but not with changes in visceral fat.

In Kahleova et al (2011), the intervention diet consisted of vegetables, fruits, grains, legumes and nuts, and a single portion of low-fat yogurt per day was permitted. The control group was administered a conventional diet for diabetes based on the guidelines of the Diabetes and Nutrition Study Group and the European Association for the Study of Diabetes. Fasting plasma C-peptide fell by $0.18 \pm 0.46 \text{ mIU/L}$ in the intervention group at 24 weeks. HbA_{1c} levels in the intervention group decreased significantly from baseline to 24 weeks (reductions of 7.1 mmol/mol vs 2.3 mmol/mol). The intervention group lost significantly more weight than the control group (6.2 kg vs 3.2 kg).

The risk of bias analysis revealed that all included trials were at a high risk of bias, mainly owing to the lack of blinding in the trials. Neither researchers nor participants were blinded to the intervention in any of the trials, as shown in Figure 2. All of the trials used randomisation; however, allocation concealment was unclear in two of the trials (Kahleova et al, 2011; 2018).

Discussion

To the best of our knowledge, this is the first systematic review investigating the effect of plant-based diets on pancreatic beta-cells. The review included three randomised controlled trials. Owing to significant heterogeneity between the trials, a meta-analysis has not been conducted.

All included studies reported beta-cell function as measured by C-peptide concentration. C-peptide is secreted by pancreatic beta-cells together with insulin. It has been widely used as a reliable measure of beta-cell function due to its lower degradation rate than insulin (Acosta-Montano et al, 2018).

This systematic review suggests that glycaemic control and beta-cell function, as measured by insulin secretion, can be improved by a plant-based diet. Postprandial insulin synthesis was significantly higher after a vegetarian burger compared with a meat burger (Kahleova et al, 2019). In the 24-week trial by Kahleova et al (2011), 43% of participants in the intervention group were required to reduce their diabetes medication. It is not clear from the data whether this was due to improved insulin sensitivity or production, or both; nonetheless, this was a clinically significant outcome.

Normal concentrations of C-peptide in people without diabetes are between 0.5 and 2.7 ng/mL .



Read more online

Dietary carbohydrate restriction as a management strategy for adults with type 2 diabetes: Exploring the opinions of dietitians

A survey to explore the opinions and practices of UK Registered Dietitians regarding low-carbohydrate diets and the suitability of the Eatwell Guide for people with type 2 diabetes.

Journal of Diabetes Nursing
223: JDN104

[Click here to access](#)

	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Blinding of participants and personnel (performance bias)	Blinding of outcome assessment (detection bias)	Incomplete outcome data (attrition bias)	Selective reporting (reporting bias)	Other bias
Kahleova et al (2011)	+	?	-	-	+	+	+
Kahleova et al (2018)	+	?	-	-	+	+	?
Kahleova et al (2019)	+	+	-	-	+	+	
Lim et al (2011)	-	-	-	-	+	+	

Figure 2. Risk of bias assessment of included studies.



Read more
online

**At a glance factsheet:
Dietary considerations in
type 1 diabetes**

This factsheet provides quick guidance on carbohydrate counting, weight management and exercise in type 1 diabetes.

Diabetes & Primary Care **22**:
3–4

[Click here to access](#)

Therefore, the reduction of 0.5 ng/mL in fasting levels in Kahleova et al (2018) was a clinically significant one, particularly when compared with the 0.5 ng/mL rise (an indirect indication of hyperinsulinaemia) seen in the control group.

The reduction in HbA_{1c} at 24 weeks observed by Kahleova et al (2011) was also clinically significant compared with the control group. HbA_{1c} levels and a reduction in diabetes medications may not be a direct measure of beta-cell function, but they are certainly a measure of glycaemia and diabetes management, and the plant-based diet groups experienced improvements in both measures in this trial.

Strengths and limitations

The main strength of this systematic review is that the trials included were all randomised controlled trials and we did not rely on observational data from cohort or case–control studies.

An important limitation is the lack of blinding in the design of the three studies. Although we intended to include non-randomised trial designs, the exclusion of the study by Lim et al (2011) meant that only randomised controlled trials have been included. A stricter selection was not possible due to the limited number of studies.

Another limitation of our systematic review is the lack of quantitative analysis. All three trials were led by the same researcher, which raises questions about the possible systematic bias that may have arisen. In addition, there was significant heterogeneity between the three trials in terms of study duration. The sample sizes across the included clinical trials were relatively small, thus decreasing the power of the individual studies. Therefore, it was decided that a meta-analysis would not be appropriate, and that a quantitative synthesis of the results to generate a pooled effect size was not possible.

Only one of the three included trials aimed to investigate the effects of plant-based diets on beta-cell function as the primary outcome. This is of importance as multiple outcomes within a study may increase the likelihood that the findings are a result of chance.

Consideration of potential confounders

A significant confounder in both of the longer trials (Kahleova et al, 2011; 2018) was caloric restriction;

however, both intervention and control groups were required to consume a calorically restricted diet in both of the trials. Calorie restriction on its own has been shown to improve beta-cell function as reported in the scientific literature. Lim et al (2011) showed that, within a week, a 600 kcal/day diet normalised beta-cell function, as measured by fasting plasma C-peptide and plasma insulin levels, as well as blood glucose levels, in people with type 2 diabetes. However, it seems there may be additional benefits gained from plant-based diets independent of caloric restriction, based on the evidence from this systematic review as well epidemiological research by Tonstad et al (2009), who reported that vegetarian diets offered protection against risk of type 2 diabetes, and that this association remained significant after adjusting for BMI.

Findings in the context of other research

Beta-cells have low glycogen stores as over 90% of the glucose that enter them is oxidised via glycolysis. This results in dependence on fatty acids for energy under fasting conditions, thus reducing intracellular lipid stores (Acosta-Montaña et al, 2018). This partly explains why beta-cell function seems to improve in response to calorie restriction or fasting. Saturated fats, found predominantly in animal fat, have been shown to reduce pancreatic beta-cell proliferation and induce lipoapoptosis, which gradually results in loss of beta-cell mass (Maedler et al, 2001). None of the included studies investigated beta-cell mass loss or examined the relationship between saturated fat intake and beta-cell mass.

Pancreatic beta-cells are particularly vulnerable to oxidative stress created by glucotoxicity, lipotoxicity or hypoxia, due to the lack of antioxidant enzymes present in the cells (Wang et al, 2017). The possible molecular mechanisms by which plant-based diets exert their effects on pancreatic beta-cells may be numerous; however, plant compounds such as flavonoids have been shown to preserve both the function and the survival of beta-cells through their antioxidant and anti-inflammatory properties (Ghorbani et al, 2019).

Vegan or vegetarian diets can easily be high in processed carbohydrates and fats; therefore, the quality of plant-based diets varies greatly. Although Kahleova et al (2011) did prepare and provide the meals for participants, the specific quantities and

ingredients of the meals are not reported. Kahleova et al (2019) did not investigate habitual intake as the intervention consisted of a single meal. It would be useful for future trials to focus on whole-food plant-based diets and report specific quantities of nutrients, minerals and vitamins consumed to allow for a comprehensive analysis.

Attrition rates in the intervention groups across the two longer studies were lower than in the control groups, suggesting that plant-based diets have high adherence rates and are sustainable. These factors are especially important as the success of dietary strategies for weight loss and diabetes management relies on how easy it is for individuals to remain on the diet over the long term. The longest-duration study, of 24 weeks, only included people who were willing to make dietary and lifestyle changes (Kahleova et al, 2011), which decreases the generalisability of the findings, as not all people with obesity or diabetes are willing to make such changes.

Restrictive diets may not be appropriate for all, and special considerations must be paid to ensure the nutritional adequacy of the diet. Of special concern are vitamin B12 and vitamin D, which have been shown to be deficient in vegan populations (Sakkas et al, 2020); therefore, plant-based dietary interventions should utilise supplements, or participants should be encouraged to consume fortified food items. Despite the restrictive nature of exclusively plant-based diets, the reviewed clinical trials showed high compliance rates.

Study quality and implications for future research

All trials were deemed to be at a high risk of bias as per the Cochrane Risk of Bias handbook (Higgins and Thomas, 2019), even though all had used adequate randomisation and the baseline characteristics of participants were comparable across studies. Based on this systematic review, further evidence is required in the form of randomised controlled trials with standardised measurements of beta-cell function and large sample sizes to determine whether plant-based diets can improve or restore beta-cell function. Unfortunately, blinding of participants and/or researchers in these trials will most likely be impractical or not possible.

Conclusion

This systematic review aimed to investigate whether a plant-based diet can positively affect pancreatic beta-cell function. Due to the limited number of clinical trials, we are unable to report a firm conclusion. A number of parameters relating to blood glucose management showed significant improvements in the plant-based groups from the included trials; however, larger randomised controlled trials with uniform measurements of beta-cell function (pre- and postprandial plasma C-peptide concentration over a period of time) would be beneficial to allow for quantitative analysis of pooled results. ■

Acosta-Montaño P, García-González V (2018) Effects of dietary fatty acids in pancreatic beta cell metabolism, implications in homeostasis. *Nutrients* **10**: 393

Cerf ME (2013) Beta cell dysfunction and insulin resistance. *Front Endocrinol (Lausanne)* **4**: 37

Cheng CW, Villani V, Buono R et al (2017) Fasting-mimicking diet promotes Ngn3-driven β -cell regeneration to reverse diabetes. *Cell* **168**: 775–88

Dinu M, Abbate R, Gensini GF et al (2017) Vegetarian, vegan diets and multiple health outcomes: a systematic review with meta-analysis of observational studies. *Crit Rev Food Sci Nutr* **57**: 3640–9

Eddouks M, Bidi A, El Bouhali B et al (2014) Antidiabetic plants improving insulin sensitivity. *J Pharm Pharmacol* **66**: 1197–214

Ghorbani A, Rashidi R, Shafiee-Nick R (2019) Flavonoids for preserving pancreatic beta cell survival and function: a mechanistic review. *Biomed Pharmacother* **111**: 947–57

Heiskanen MA, Motiani KK, Mari A et al (2018) Exercise training decreases pancreatic fat content and improves beta cell function regardless of baseline glucose tolerance: a randomised controlled trial. *Diabetologia* **61**: 1817–28

Higgins J, Thomas J (2019) *Cochrane Handbook for Systematic Reviews of Interventions* (version 6.0). Cochrane, London. Available at: <https://training.cochrane.org/handbook> (accessed 11.03.21)

InterAct Consortium (2015) Dietary fibre and incidence of type 2 diabetes in eight European countries: the EPIC-InterAct Study and a meta-analysis of prospective studies. *Diabetologia* **58**: 1394–408

Kahleova H, Matoulek M, Malinska H et al (2011) Vegetarian diet improves insulin resistance and oxidative stress markers more than conventional diet in subjects with type 2 diabetes. *Diabet Med* **28**: 549–59

Kahleova H, Tura A, Hill M et al (2018) A plant-based dietary intervention improves beta-cell function and insulin resistance in overweight adults: a 16-week randomized clinical trial. *Nutrients* **10**: 189

Kahleova H, Tura A, Klementova M et al (2019) A plant-based meal stimulates incretin and insulin secretion more than an energy- and macronutrient-matched standard meal in type 2 diabetes: a randomized crossover study. *Nutrients* **11**: 486

Larson-Meyer DE, Heilbronn LK, Redman LM et al (2006) Effect of calorie restriction with or without exercise on insulin sensitivity, beta-cell function, fat cell size, and ectopic lipid in overweight subjects. *Diabetes Care* **29**: 1337–44

Lim EL, Hollingsworth KG, Aribisala BS et al (2011) Reversal of type 2 diabetes: normalisation of beta cell function in association with decreased pancreas and liver triacylglycerol. *Diabetologia* **54**: 2506–14

Maedler K, Spinas GA, Dyntar D et al (2001) Distinct effects of saturated and monounsaturated fatty acids on beta-cell turnover and function. *Diabetes* **50**: 69–76

Sakkas H, Bozidis P, Touzios C et al (2020) Nutritional status and the influence of the vegan diet on the gut microbiota and human health. *Medicina (Kaunas)* **56**: 88

References (continued)

Sakurai Y, Tamura Y, Takeno K et al (2011) Determinants of intramyocellular lipid accumulation after dietary fat loading in non-obese men. *J Diabetes Investig* **2**: 310–7

SIGN (2015) *Checklists*. SIGN, Edinburgh. Available at: www.sign.ac.uk/what-we-do/methodology/checklists (accessed 17.03.21)

Suh KS, Choi EM, Jung WW et al (2017) Deoxyactein protects pancreatic β -cells against methylglyoxal-induced oxidative cell damage by the upregulation of mitochondrial biogenesis. *Int J Mol Med* **40**: 539–48

Tonstad S, Butler T, Yan R, Fraser GE (2009) Type of vegetarian diet, body weight, and prevalence of type 2 diabetes. *Diabetes Care* **32**: 791–6

Uribarri J, Woodruff S, Goodman S et al (2010) Advanced glycation end products in foods and a practical guide to their reduction in the diet. *J Am Diet Assoc* **110**: 911–16

Wang J, Wang H (2017) Oxidative stress in pancreatic beta cell regeneration. *Oxid Med Cell Longev* **2017**: 1930261