

THE HOMOLOGY OF THE TEMPERLEY-LIEB ALGEBRAS

RACHAEL BOYD AND RICHARD HEPWORTH

ABSTRACT. This paper studies the homology and cohomology of the Temperley-Lieb algebra $TL_n(a)$, interpreted as appropriate Tor and Ext groups. Our main result applies under the common assumption that $a = v + v^{-1}$ for some unit v in the ground ring, and states that the homology and cohomology vanish up to and including degree $(n - 2)$. To achieve this we simultaneously prove homological stability and compute the stable homology. We show that our vanishing range is sharp when n is even.

Our methods are inspired by the tools and techniques of homological stability for families of groups. We construct and exploit a chain complex of ‘planar injective words’ that is analogous to the complex of injective words used to prove stability for the symmetric groups. However, in this algebraic setting we encounter a novel difficulty: $TL_n(a)$ is not flat over $TL_m(a)$ for $m < n$, so that Shapiro’s lemma is unavailable. We resolve this difficulty by constructing what we call ‘inductive resolutions’ of the relevant modules.

Vanishing results for the homology and cohomology of Temperley-Lieb algebras can also be obtained from the existence of the Jones-Wenzl projector. Our own vanishing results are in general far stronger than these, but in a restricted case we are able to obtain additional vanishing results via the existence of the Jones-Wenzl projector.

We believe that these results, together with the second author’s work on Iwahori-Hecke algebras, are the first time the techniques of homological stability have been applied to algebras that are not group algebras.

CONTENTS

1. Introduction	2
2. Temperley-Lieb Algebras	10
3. Inductive resolutions	19
4. Planar injective words	26
5. Homological stability and stable homology	33
6. Sharpness	38
7. The case of $TL_2(a)$	40
8. High-acyclicity	42
9. Jones-Wenzl projectors and vanishing	59
References	64

2010 *Mathematics Subject Classification.* 20J06, 16E40 (primary), 20F36 (secondary) .
Key words and phrases. Homological stability, Temperley-Lieb algebras.

1. INTRODUCTION

In this work we study the homology and cohomology of the *Temperley-Lieb algebras*. In particular, we simultaneously prove that the algebras satisfy homological stability, and that their stable homology vanishes.

A sequence of groups and inclusions $G_0 \rightarrow G_1 \rightarrow G_2 \rightarrow \cdots$ is said to satisfy *homological stability* if for each degree d the induced sequence of homology groups

$$H_d(G_0) \rightarrow H_d(G_1) \rightarrow H_d(G_2) \rightarrow \cdots$$

eventually consists of isomorphisms. Homological stability can also be formulated for sequences of spaces. There are many important examples of groups and spaces for which homological stability is known to hold, such as symmetric groups [Nak60], general linear groups [Cha80, Maa79, vdK80], mapping class groups of surfaces [Har85, RW16] and 3-manifolds [HW10], automorphism groups of free groups [HV98, HV04], diffeomorphism groups of high-dimensional manifolds [GRW18], configuration spaces [Chu12, RW13], Coxeter groups [Hep16], Artin monoids [Boyd20], and many more. In almost all cases, homological stability is one of the strongest things we know about the homology of these families. It is often coupled with computations of the stable homology $\lim_{n \rightarrow \infty} H_*(G_n)$, which is equal to the homology of the G_n in the *stable range* of degrees, i.e. those degrees for which stability holds.

The homology and cohomology of a group G can be expressed in the language of homological algebra as

$$H_*(G) = \mathrm{Tor}_*^{RG}(\mathbb{1}, \mathbb{1}), \quad H^*(G) = \mathrm{Ext}_{RG}^*(\mathbb{1}, \mathbb{1}),$$

where R is the coefficient ring for homology and cohomology, RG is the group algebra of G and $\mathbb{1}$ is its trivial module. Thus the homology and cohomology of a group depend only on the group algebra RG and its trivial module $\mathbb{1}$. It is therefore natural to consider the homology and cohomology of an arbitrary algebra equipped with a ‘trivial’ module. Moreover, one may ask whether homological stability occurs in this wider context.


In [Hep20] the second author proved homological stability for *Iwahori-Hecke algebras* of type A . These are deformations of the group rings of the symmetric groups that are important in representation theory, knot theory, and combinatorics. There is a fairly standard suite of techniques used to prove homological stability, albeit with immense local variation, and the proof strategy of [Hep20] followed all the steps familiar from the setting of groups. As is typical, the hardest step was to prove that the homology of a certain (chain) complex vanishes in a large range of degrees.

In the present paper we will prove homological stability for the *Temperley-Lieb algebras*, and we will prove that the stable homology vanishes. However amongst the familiar steps in our proof lies a novel obstacle and — to counter it — a novel construction. At a certain point the usual techniques fail because Shapiro’s lemma


cannot be applied, as we will explain below. This is a new difficulty that never occurs in the setting of groups, but we are able to resolve it for the algebras at hand, and in fact our solution facilitates the unusually strong results that we are able to obtain. It is not surprising that the Iwahori-Hecke case is more straightforward than the Temperley-Lieb case: Iwahori-Hecke algebras are deformations of group rings, whereas the Temperley-Lieb algebras are significantly different.

To the best of our knowledge, the present paper and [Hep20] are the first time the techniques of homological stability have been applied to algebras that are not group algebras, and together they serve as proof-of-concept for the export of homological stability techniques to the setting of algebras. The moral of [Hep20] is that the ‘usual’ techniques of homological stability suffice, so long as the algebras involved satisfy a certain flatness condition. The moral of the present paper is that failure of the flatness condition can in some cases be overcome, using new ingredients and techniques, and can even lead to stronger results than in the flat scenario. Since the completion of this paper, we have extended our techniques to study the homology of the Brauer algebras in joint work with Patzt [BHP21].

1.1. Temperley-Lieb algebras. Let $n \geq 0$, let R be a commutative ring, and let $a \in R$. The *Temperley-Lieb algebra* $TL_n(a)$ is the R -algebra with basis (by which we will always mean R -module basis) given by the planar diagrams on n strands, taken up to isotopy, and with multiplication given by pasting diagrams and replacing closed loops with factors of a . The last sentence was intentionally brief, but we hope that its meaning becomes clearer with the following illustration of two elements $x, y \in TL_5(a)$



and their product $x \cdot y$.



The Temperley-Lieb algebras arose in theoretical physics in the 1970s [TL71]. They were later rediscovered by Jones in his work on von Neumann algebras [Jon83], and used in the first definition of the Jones polynomial [Jon85]. Kauffman gave the above diagrammatic interpretation of the algebras in [Kau87] and [Kau90].

The Temperley-Lieb algebra $TL_n(a)$ is perhaps best studied in the case where $a = v + v^{-1}$, for $v \in R$ a unit. In this case, it is a quotient of the Iwahori-Hecke algebra of type A_{n-1} with parameter $q = v^2$ (so it is closely related to the symmetric group) and it receives a homomorphism from the group algebra

of the braid group on n strands. It can also be described as the endomorphism algebra of $V_q^{\otimes n}$, where V_q is a certain 2-dimensional representation of the quantum group $U_q(\mathfrak{sl}_2)$. We recommend [RSA14] and [KT08] for further reading on $\mathrm{TL}_n(a)$, and [Wes95] and [GL96] for details on their representation theory.

1.2. Homology of Temperley-Lieb algebras. The Temperley-Lieb algebra $\mathrm{TL}_n(a)$ has a *trivial module* $\mathbb{1}$ consisting of a copy of R on which all diagrams other than the identity diagram act as multiplication by 0. It therefore has homology and cohomology groups $\mathrm{Tor}_*^{\mathrm{TL}_n(a)}(\mathbb{1}, \mathbb{1})$ and $\mathrm{Ext}_{\mathrm{TL}_n(a)}^*(\mathbb{1}, \mathbb{1})$.

Our first result is a vanishing theorem in the case that the parameter $a \in R$ is invertible.

Theorem A. *Let R be a commutative ring, and let a be a unit in R . Then $\mathrm{Tor}_d^{\mathrm{TL}_n(a)}(\mathbb{1}, \mathbb{1})$ and $\mathrm{Ext}_{\mathrm{TL}_n(a)}^d(\mathbb{1}, \mathbb{1})$ both vanish for $d > 0$.*

The next result holds regardless of whether or not a is invertible, and uses the common assumption that $a = v + v^{-1}$, $v \in R^\times$. However we see shortly that this assumption can be removed.

Theorem B. *Let R be a commutative ring, let $v \in R$ be a unit, let $a = v + v^{-1}$, and let $n \geq 0$. Then*

$$\mathrm{Tor}_d^{\mathrm{TL}_n(a)}(\mathbb{1}, \mathbb{1}) = 0 \quad \text{and} \quad \mathrm{Ext}_{\mathrm{TL}_n(a)}^d(\mathbb{1}, \mathbb{1}) = 0$$

for $1 \leq d \leq (n-2)$ if n is even, and for $1 \leq d \leq (n-1)$ if n is odd.

Thus the map $\mathrm{Tor}_d^{\mathrm{TL}_{n-1}(a)}(\mathbb{1}, \mathbb{1}) \rightarrow \mathrm{Tor}_d^{\mathrm{TL}_n(a)}(\mathbb{1}, \mathbb{1})$ is an isomorphism for $d \leq n-3$, so that we have homological stability, and $\lim_{n \rightarrow \infty} \mathrm{Tor}_*^{\mathrm{TL}_n(a)}(\mathbb{1}, \mathbb{1}) = 0$ in positive degrees, so the stable homology is trivial. The latter is reminiscent of Quillen's result on the vanishing stable homology of general linear groups of finite fields in defining characteristic [Qui72], and of Szymik-Wahl's result on the acyclicity of the Thompson groups [SW19]. Theorems A and B might lead us to expect that the homology and cohomology of the $\mathrm{TL}_n(a)$ are largely trivial, but in fact the results are as strong as possible, at least for n even:

Theorem C. *In the setting of Theorem B above, suppose further that n is even and that $a = v + v^{-1}$ is not a unit. Then $\mathrm{Tor}_{n-1}^{\mathrm{TL}_n(a)}(\mathbb{1}, \mathbb{1}) \neq 0$.*

Thus Theorem A does not extend to the case of a not invertible, and the stable range in Theorem B is sharp. In fact we can say more: When n is even, $\mathrm{Tor}_{n-1}^{\mathrm{TL}_n(a)}(\mathbb{1}, \mathbb{1}) \cong R/bR$ where b is a multiple of a (unfortunately our methods do not allow us to say anything more concrete about b).

Remark. One can compute $\mathrm{Tor}_1^{\mathrm{TL}_n(a)}(\mathbb{1}, \mathbb{1})$ directly using the method of [Wei94, Exercise 3.1.3]: it is R/aR for $n = 2$, and vanishes otherwise. We also compute the homology and cohomology of $\mathrm{TL}_2(a)$ by an explicit resolution: $\mathrm{Tor}_*^{\mathrm{TL}_2(a)}(\mathbb{1}, \mathbb{1})$ is R/aR in odd degrees, and the kernel R_a of $r \mapsto ar$ in positive even degrees,

so that if a is not invertible then $\mathrm{Tor}_*^{\mathrm{TL}_2(a)}(\mathbb{1}, \mathbb{1})$ is non-trivial in infinitely many degrees.

In [RW21], Randal-Williams shows that in fact you can remove our assumption that $a = v + v^{-1}$ for a unit $v \in R$, by applying Theorem C for an associated ring S . This yields the following strengthening of Theorem B.

Corollary. [RW21, Theorem B'] *Let R be a commutative ring, a be any element in R , and $n \geq 0$. Then*

$$\mathrm{Tor}_d^{\mathrm{TL}_n(a)}(\mathbb{1}, \mathbb{1}) = 0 \quad \text{and} \quad \mathrm{Ext}_{\mathrm{TL}_n(a)}^d(\mathbb{1}, \mathbb{1}) = 0$$

for $1 \leq d \leq (n - 2)$ if n is even, and for $1 \leq d \leq (n - 1)$ if n is odd.

Proof. The full proof can be found in [RW21], and uses the Base Change Spectral Sequence [Wei94, Section 5.6]. This is applied to the faithfully flat ring homomorphism $R \rightarrow S$ where $S = R[v]/(v^2 - a \cdot v + 1)$, which by construction has a unit v and element a such that $a = v + v^{-1}$. The results in Theorem B for the ring S can now be transferred to analogous results for the ring R . \square

1.3. Jones-Wenzl projectors. The *Jones-Wenzl projector* or *Jones-Wenzl idempotent* JW_n , if it exists, is the element of $\mathrm{TL}_n(a)$ uniquely characterised by the following two properties:

- $\mathrm{JW}_n \in 1 + I_n$
- $\mathrm{JW}_n \cdot I_n = 0 = I_n \cdot \mathrm{JW}_n$

where I_n is the two-sided ideal in $\mathrm{TL}_n(a)$ spanned by all diagrams other than the identity diagram. The Jones-Wenzl projector was first introduced by Jones [Jon83], was further studied by Wenzl [Wen87], and has since become important in representation theory, knot theory and the study of 3-manifolds.

The Jones-Wenzl projector exists if and only if the trivial module $\mathbb{1}$ is projective. Moreover, when the ground ring R is a field, there is a simple and explicit criterion for the existence of JW_n , given in terms of the parameter a . Thus, when this criterion holds, the vanishing of $\mathrm{Tor}_*^{\mathrm{TL}_n(a)}(\mathbb{1}, \mathbb{1})$ and $\mathrm{Ext}_{\mathrm{TL}_n(a)}^*(\mathbb{1}, \mathbb{1})$ in positive degrees follows immediately.

Our own Theorems A and B are in general far stronger than the vanishing results obtained from the existence of JW_n , as they do not require R to be a field, and the constraints are weaker. Indeed, in the case of n even, Theorems A and C are the final word on vanishing, since they imply that the homology and cohomology of $\mathrm{TL}_n(a)$ vanish in all positive degrees if and only if a is invertible. However, in the case of n odd and R a field, there are some situations where our theorems do not incorporate all vanishing results given by the existence of JW_n . These cases are encapsulated in the following.

Theorem D. *Let $n = 2k + 1$, and let R be a field whose characteristic does not divide $\binom{k}{t}$ for any $1 \leq t \leq k$. Let v be a unit in R and assume that $a = v + v^{-1} = 0$. Then $\mathrm{Tor}_*^{\mathrm{TL}_n(0)}(\mathbb{1}, \mathbb{1})$ and $\mathrm{Ext}_{\mathrm{TL}_n(0)}^*(\mathbb{1}, \mathbb{1})$ vanish in positive degrees.*

As with Theorem B, the assumption that $a = v + v^{-1}$ for v a unit can be removed in this result.

Combining Theorem D with Theorem A yields rather comprehensive vanishing results when R is a field with appropriate characteristic. For example, it now follows that when R is any field, the homology and cohomology of $\mathrm{TL}_3(v + v^{-1})$ vanish regardless of the choice of v . Similarly, the homology and cohomology of $\mathrm{TL}_5(v + v^{-1})$ will vanish over any field and for any value of v , except possibly in characteristic 2 when $v + v^{-1} = 0$. Since the first appearance of our paper, Sroka [Sro22] has used related techniques to show that when n is odd the Tor groups vanish in all positive degrees, for any choice of R .

The next few sections of this introduction will discuss the proofs of our main results in some detail.

1.4. Planar injective words. Several proofs of homological stability for the symmetric group [Maa79, Ker05, RW13] make use of the *complex of injective words*. This is a highly-connected complex with an action of the symmetric group \mathfrak{S}_n . Our main tool for proving Theorems B and C is the *complex of planar injective words* $W(n)$, a Temperley-Lieb analogue of the complex of injective words that we introduce and study here for the first time. It is a chain complex of $\mathrm{TL}_n(a)$ -modules, and in degree i it is given by the tensor product module $\mathrm{TL}_n(a) \otimes_{\mathrm{TL}_{n-i-1}(a)} \mathbb{1}$. This is analogous to the complex of injective words, whose i -simplices form a single \mathfrak{S}_n -orbit with typical stabiliser \mathfrak{S}_{n-i-1} , which is an alternative way of saying that the i -th chain group is isomorphic to $R\mathfrak{S}_n \otimes_{R\mathfrak{S}_{n-i-1}} \mathbb{1}$. We show the following high-acyclicity result. In order to construct appropriate differentials for $W(n)$ we exploit a homomorphism from the group algebra of the braid group on n strands, which is not necessarily apparent from the definition of $\mathrm{TL}_n(a)$. This is where the restriction of a to $a = v + v^{-1}$ is necessary.

Theorem E. $H_d(W(n))$ vanishes in degrees $d \leq (n - 2)$.

The complex $W(n)$ has rich combinatorial properties, analogous to those of the complex of injective words, that we explore in the companion paper [BH21]. In particular, Theorem E tells us that the homology of $W(n)$ is concentrated in the top degree $H_{n-1}(W(n))$, and in [BH21] we show that when R is Noetherian the rank of this top homology group is the n -th *Fine number* F_n [DS01], an analogue of the number of derangements on n letters. Furthermore we show that the differentials of $W(n)$ encode the *Jacobsthal numbers* [OEIS]. Finally in the semisimple case we show that $H_{n-1}(W(n))$ has descriptions firstly categorifying an alternating sum for the Fine numbers, and secondly in terms of standard Young tableaux. We call the $\mathrm{TL}_n(a)$ -module $H_{n-1}(W(n))$ the *Fineberg module*, and we denote it $\mathcal{F}_n(a)$. We know little about $\mathcal{F}_n(a)$ in general, though in the cases $n = 2, 3, 4$ we give examples describing it in terms of the cell modules of $\mathrm{TL}_n(a)$.

The proof of Theorem E is perhaps the most difficult technical result in this paper. It is obtained by filtering $W(n)$ and showing that the filtration quotients

are (suspensions of truncations of) copies of $W(n-1)$, and then proceeding by induction.

1.5. Spectral sequences and Shapiro's lemma. Let us now outline how we use the complex of planar injective words $W(n)$ to prove Theorems B and C. Following standard approaches to homological stability for groups, we consider a spectral sequence obtained from the complex $W(n)$. The E^1 -page of our spectral sequence consists of the groups $\mathrm{Tor}_j^{\mathrm{TL}_n(a)}(\mathbb{1}, \mathrm{TL}_n(a) \otimes_{\mathrm{TL}_{n-i-1}(a)} \mathbb{1})$. Furthermore, thanks to Theorem E, the spectral sequence converges to $\mathrm{Tor}_{*-n+1}^{\mathrm{TL}_n(a)}(\mathbb{1}, \mathcal{F}_n(a))$, where $\mathcal{F}_n(a) = H_{n-1}(W(n))$ is the Fineberg module. Our experience from homological stability tells us to apply *Shapiro's lemma*, or in this context a *change-of-rings isomorphism*, to identify

$$\mathrm{Tor}_*^{\mathrm{TL}_n(a)}(\mathbb{1}, \mathrm{TL}_n(a) \otimes_{\mathrm{TL}_{n-i-1}(a)} \mathbb{1}) \quad \text{with} \quad \mathrm{Tor}_*^{\mathrm{TL}_{n-i-1}(a)}(\mathbb{1}, \mathbb{1}).$$

This identification applied to the columns of our spectral sequence would allow us to implement an inductive hypothesis. However, such a change-of-rings isomorphism would only be valid if $\mathrm{TL}_n(a)$ were flat as a $\mathrm{TL}_{n-i-1}(a)$ -module, and this is not the case. This failure of Shapiro's lemma is a potentially serious obstacle to proceeding further. However, we are able to identify the columns of our spectral sequence by independent means, as follows:

Theorem F. *Let R be a commutative ring and let $a \in R$. Let $0 \leq m < n$. Then $\mathrm{Tor}_d^{\mathrm{TL}_n(a)}(\mathbb{1}, \mathrm{TL}_n(a) \otimes_{\mathrm{TL}_m(a)} \mathbb{1})$ and $\mathrm{Ext}_{\mathrm{TL}_n(a)}^d(\mathrm{TL}_n(a) \otimes_{\mathrm{TL}_m(a)} \mathbb{1}, \mathbb{1})$ both vanish for $d > 0$.*

In conjunction with a computation of the $d = 0$ case, this gives us the vanishing results of Theorem B. Moreover, in the case of n even we are able to analyse the rest of the spectral sequence (there is a single differential and a single extension problem) in sufficient detail to prove the sharpness result of Theorem C. This involves a careful study of the Fineberg module $\mathcal{F}_n(a)$. In general, our method identifies $\mathrm{Tor}_*^{\mathrm{TL}_n(a)}(\mathbb{1}, \mathbb{1})$ with $\mathrm{Tor}_{*-n}^{\mathrm{TL}_n(a)}(\mathbb{1}, \mathcal{F}_n(a))$, except in degrees $* = n - 1, n$ when n is even.

1.6. Inductive resolutions. It remains for us to discuss the proofs of Theorems A and F. These results are proved by a novel method that exploits the structure of the Temperley-Lieb algebras, and in particular they lie outwith the standard toolkit of homological stability. Moreover, it is Theorem F which allows us to overcome the failure of Shapiro's lemma.

The two theorems are very similar: Theorem A is an instance of the more general statement that $\mathrm{Tor}_*^{\mathrm{TL}_n(a)}(\mathbb{1}, \mathrm{TL}_n(a) \otimes_{\mathrm{TL}_m(a)} \mathbb{1})$ vanishes in positive degrees for $m \leq n$ and a invertible, while Theorem F states that the same groups vanish for $m < n$ and a arbitrary. These are both proved by strong induction on m . The initial cases $m = 0, 1$ are immediate because then $\mathrm{TL}_m(a) = R$ so $\mathrm{TL}_n(a) \otimes_{\mathrm{TL}_m(a)} \mathbb{1}$ is free. The induction step is proved by constructing and exploiting a resolution

of $\mathrm{TL}_n(a) \otimes_{\mathrm{TL}_m(a)} \mathbb{1}$ whose terms have the form $\mathrm{TL}_n(a) \otimes_{\mathrm{TL}_{m-1}(a)} \mathbb{1}$ and $\mathrm{TL}_n(a) \otimes_{\mathrm{TL}_{m-2}(a)} \mathbb{1}$, and then applying the inductive hypothesis. We call these resolutions *inductive resolutions* since they resolve the next module in terms of those already considered.

Our technique of inductive resolutions is generalised in a joint paper with Patzt, [BHP21], where we show that the homology of the Brauer algebras is isomorphic to the homology of the symmetric groups in a stable range when the parameter δ is not invertible, and in every degree when δ is invertible. This provides concrete evidence that the new techniques developed in this paper can be adapted to other algebras to obtain results of similar strength.

1.7. Discussion: Homological stability for algebras. As stated earlier, we regard the present paper, together with the results of [Hep20] on Iwahori-Hecke algebras, as proof-of-concept for the export of the techniques of homological stability to the setting of algebras. And, since the first appearance of this paper, these techniques have been extended to the setting of Brauer algebras in our joint work with Patzt [BHP21]. We hope that the present paper, together with [Hep20, BHP21], will be a springboard for further research in this direction.

One of the main motivations for studying the homology of groups, is that homology is a useful ‘measurement’ of the group. Put another way, homology is a powerful invariant, where the power comes from the fact that it is both informative, and (relatively) computable. The Tor and Ext groups of algebras are likewise strong invariants, and it is our hope that homology and cohomology of algebras can be utilised as a tool to answer questions in the fields where the algebras arise. For example, modern representation theory is rich in conjectures, and home to surprising isomorphisms between apparently very different algebras [BK09, BCH20]. Understanding the similarities and differences between naturally-arising algebras is precisely the kind of question that could be investigated via Tor and Ext-groups.

We will now discuss some questions arising from our work. Readers with experience in homological stability will be able to think of many new questions in this direction, so we will simply list some that are most prominent in our minds.

The Temperley-Lieb algebra can be regarded as an algebra of 1-dimensional cobordisms embedded in 2 dimensions, and the Brauer algebra can similarly be viewed as an algebra of 1-dimensional cobordisms embedded in infinite dimensions.

Question. Are there analogues of the Temperley-Lieb algebra consisting of d -dimensional cobordisms embedded in n dimensions? Does homological stability hold for these algebras? And can the stability be understood in an essentially geometric way?

And more generally:

Question. For which natural families of algebras does homological stability hold?

Candidate algebras, closely related to the existing cases, are: Iwahori-Hecke and Temperley-Lieb algebras of types B and D ; the periodic and dilute Temperley-Lieb algebras; and the blob, partition and Birman-Murakami-Wenzl algebras. We invite the reader to think of possibilities from further afield.

There have recently been advances in building general frameworks for homological stability proofs. In [RWW17] Randal-Williams and Wahl introduce a categorical framework that encapsulates, improves and extends several of the standard techniques used in homological stability proofs for groups. In [GKRW18] Galatius, Kupers and Randal-Williams introduce a framework that applies to E_k -algebras in simplicial modules. It exploits the notion of cellular E_k -algebras, and incorporates methods for proving *higher stability* results. This invites us to pose the following questions.

Question. Does the general homological stability machinery of Randal-Williams and Wahl [RWW17] generalise to an R -linear version, giving a general framework to prove that a family of R -algebras $A_0 \rightarrow A_1 \rightarrow A_2 \rightarrow \dots$ satisfies homological stability?

In this question, the most interesting issue is what form the resulting complexes will take. One might expect that for a family of algebras the relevant complexes will be constructed from tensor products, as with our complex $W(n)$. However it may happen, as in this paper, that flatness issues arise, in which case it seems unlikely that complexes built from the honest tensor products will be sufficient.

Question. Can the homological stability machinery of Galatius, Kupers and Randal-Williams [GKRW18] be applied in the setting of algebras?

It seems extremely likely that homology of Temperley-Lieb algebras will indeed fit into the framework of [GKRW18], by using appropriate simplicial models for the $\mathrm{Tor}_*^{\mathrm{TL}_n(a)}(\mathbb{1}, \mathbb{1})$, or more precisely for the chain complexes underlying these Tor groups. Again, the difficulty will lie in identifying and computing the associated *splitting complexes*, especially when flatness issues arise.

1.8. Outline. In Section 2 we recall the definition of the Temperley-Lieb algebra, the Jones basis, the relationship with Iwahori-Hecke algebras, and we establish results on the induced modules $\mathrm{TL}_n(a) \otimes_{\mathrm{TL}_m(a)} \mathbb{1}$ that will be important in the rest of the paper. Section 3 establishes our inductive resolutions and proves Theorems A and F. Section 4 introduces the complex of planar injective words $W(n)$ and the Fineberg module $\mathcal{F}_n(a)$. Sections 5 and 6 then use $W(n)$, in particular its high-acyclicity (Theorem E), to prove Theorems B and C. Section 7 investigates our results in the case of $\mathrm{TL}_2(a)$, computing the homology directly and also in terms of the Fineberg module $\mathcal{F}_2(a)$. Section 8 proves Theorem E. Section 9 investigates the vanishing results given by the Jones-Wenzl projectors and proves Theorem D.

1.9. Acknowledgements. The authors would like to thank the Max Planck Institute for Mathematics in Bonn for its support and hospitality. We are grateful to Robert Spencer and Paul Wedrich for interesting and helpful discussions about the representation theory of Temperley-Lieb algebras, and to Oscar Randal-Williams for his extension of Theorem B. We would also like to thank the anonymous referees for very insightful comments, which have much improved this paper.

2. TEMPERLEY-LIEB ALGEBRAS


In this section we will cover the basic facts about the Temperley-Lieb algebra that we will need for the rest of the paper. There is some overlap between the material recalled here and in [BH21]. In particular, we cover the definitions by generators and relations and by diagrams; we discuss the Jones basis for $\mathrm{TL}_n(a)$; we look at the induced modules $\mathrm{TL}_n(a) \otimes_{\mathrm{TL}_m(a)} \mathbb{1}$ that will be an essential ingredient in all that follows; and we discuss the homomorphism from the Iwahori-Hecke algebra of type A_{n-1} into $\mathrm{TL}_n(a)$. Historical references on Temperley-Lieb algebras were given in the introduction. General references for readers new to the $\mathrm{TL}_n(a)$ are Section 5.7 of Kassel and Turaev's book [KT08] on the braid groups, and especially Sections 1 and 2 of Ridout and Saint-Aubin's survey on the representation theory of the $\mathrm{TL}_n(a)$ [RSA14].

Definition 2.1 (The Temperley-Lieb algebra $\mathrm{TL}_n(a)$). Let R be a commutative ring and let $a \in R$. Let n be a non-negative integer. The *Temperley-Lieb algebra* $\mathrm{TL}_n(a)$ is defined to be the R -algebra with generators U_1, \dots, U_{n-1} and the following relations:


- (1) $U_i U_j = U_j U_i$ for $j \neq i \pm 1$
- (2) $U_i U_j U_i = U_i$ for $j = i \pm 1$
- (3) $U_i^2 = a U_i$ for all i .

Thus elements of the Temperley-Lieb algebra are formal sums of monomials in the U_i , with coefficients in the ground ring R , modulo the relations above. We often write $\mathrm{TL}_n(a)$ as TL_n . We note here that $\mathrm{TL}_0 = \mathrm{TL}_1 = R$.

There is an alternative definition of TL_n in terms of diagrams. In this description, an element of TL_n is an R -linear combination of *planar diagrams* (or 1-dimensional cobordisms). Each planar diagram consists of two vertical lines in the plane, decorated with n dots labelled $1, \dots, n$ from bottom to top, together with a collection of n arcs joining the dots in pairs. The arcs must lie between the vertical lines, they must be disjoint, and the diagrams are taken up to isotopy. For example, here are two planar diagrams in the case $n = 5$:




We will often omit the labels on the dots. Multiplication of diagrams is given by placing them side-by-side and joining the ends. Any closed loops created by this process are then erased and replaced with a factor of a . For example, the product xy of the elements x and y above is:



(We have subscribed to the heresy of [RSA14] by drawing planar diagrams that go from left to right rather than top to bottom.)

One can pass from the generators-and-relations definition of TL_n in Definition 2.1 to the diagrammatic description of the previous paragraph as follows. For $1 \leq i \leq n - 1$, to each U_i we associate the planar diagram shown below.



We refer to an arc joining adjacent dots as a *cup*. The relations for the Temperley-Lieb algebras are satisfied, and two of them are illustrated in Figure 1. The fact that this determines an isomorphism between the algebra defined by generators and relations, and the one defined by diagrams, is proved in [RSA14, Theorem 2.4], [KT08, Theorem 5.34], and [Kau05, Section 6].

In the rest of the paper we will refer to the diagrammatic point of view on the Temperley-Lieb algebra, but we will not rely on it for any proofs.

2.1. The Jones basis. From the diagrammatic point of view the Temperley-Lieb algebra TL_n has an evident R -basis given by the (isotopy classes of) planar diagrams. This is called the *diagram basis*. We now recall the analogue of the diagram basis given in terms of the U_i , which is called the *Jones basis* for TL_n , and we prove some additional facts about it that we will require later. See [KT08, Section 5.7], [RSA14, Section 2] or [Kau05, Section 6], but note that conventions vary, and see Remark 2.5 below in particular.

Definition 2.2 (Jones normal form). The *Jones normal form* for elements of $TL_n(a)$ is defined as follows. Let

$$n > a_k > a_{k-1} > \cdots > a_1 > 0 \quad n > b_k > b_{k-1} > \cdots > b_1 > 0$$




 (a) The relation $U_i^2 = aU_i$.

 (b) The relation $U_i U_{i+1} U_i = U_i$.

 FIGURE 1. Diagrammatic relations in TL_n .

be integers such that $b_i \geq a_i$ for all i . Let $\underline{a} = (a_k, \dots, a_1)$ and $\underline{b} = (b_k, \dots, b_1)$. Then set

$$x_{\underline{a}, \underline{b}} = (U_{a_k} \dots U_{b_k}) \cdot (U_{a_{k-1}} \dots U_{b_{k-1}}) \cdots (U_{a_1} \dots U_{b_1})$$

where the subscripts of the generators increase in each tuple $U_{a_i} \dots U_{b_i}$. A word written in the form $x_{\underline{a}, \underline{b}}$ is said to be written in *Jones normal form* for $\text{TL}_n(a)$.

Example 2.3. In TL_5 the words

$$U_1 U_2 U_3 U_4 = (U_1 U_2 U_3 U_4) = x_{(1), (4)}$$

$$U_4 U_3 U_2 U_1 = (U_4) \cdot (U_3) \cdot (U_2) \cdot (U_1) = x_{(4,3,2,1), (4,3,2,1)}$$

$$U_3 U_4 U_1 U_2 = (U_3 U_4) \cdot (U_1 U_2) = x_{(3,1), (4,2)}$$

$$U_2 U_3 U_1 U_2 = (U_2 U_3) \cdot (U_1 U_2) = x_{(2,1), (3,2)}$$

are in Jones normal form. The word $U_2 U_1 U_4 U_2 U_3$ is not, but it can be rewritten using the defining relations to give

$$U_2 U_1 U_4 U_2 U_3 = U_4 U_2 U_1 U_2 U_3 = U_4 U_2 U_3 = (U_4)(U_2 U_3) = x_{(4,2), (4,3)}.$$


Denote the subset of TL_n consisting of all $x_{\underline{a}, \underline{b}}$ with $\underline{a} = (a_1, \dots, a_k)$ and $\underline{b} = (b_1, \dots, b_k)$ by $\text{TL}_{n,k}$. Then the set

$$\text{TL}_{n,0} \sqcup \text{TL}_{n,1} \sqcup \cdots \sqcup \text{TL}_{n,n-1}$$

is a basis (recall that by basis we always mean R -module basis) of TL_n , called the *Jones basis*. For a proof of this fact see [KT08, Corollary 5.32], [RSA14, pp.967-969] or [Kau05, Section 6], though we again warn the reader that conventions vary.

There is an algorithm for taking a diagram and writing it as an element of the Jones basis, see [Kau05, Section 6]. We summarise the algorithm here. Let the i -th row of the diagram be the horizontal strip whose left and right ends lie between the dots i and $(i + 1)$ on each vertical line. Take a planar diagram, and ensure that it is drawn in minimal form: all arcs connecting the same side of the diagram to itself are drawn as semicircles, and all arcs from left to right are drawn without any cups, i.e. transverse to all vertical lines, and such that each arc of the diagram intersects each row transversely and at most once.

Proceed along each row of the diagram, connecting the consecutive arcs encountered with a dotted horizontal line labelled by the row in question. This is done in an alternating fashion: the first arc encountered is connected to the second by a dotted line, then the third is connected to the fourth, and so on. If we start with the elements x and y used earlier in this section, then this gives us the following:



A *sequence* in such a decorated diagram is taken by travelling right along the dotted arcs and up along the solid arcs from one dotted arc to the next, starting as far to the left as possible. The above diagrams each have two sequences, indicated in blue and red. The sequences in a diagram are linearly ordered by scanning from top to bottom and recording a sequence when one of its dotted lines is first encountered. So in the above diagrams the blue sequences precede the red ones. One now obtains a Jones normal form for the element by working through the sequences in turn, writing out the labels from left to right, and then taking the corresponding monomial in the U_i :

$$x = (U_4)(U_1U_2) = x_{(4,1),(4,2)}, \quad y = (U_2U_3U_4)(U_1U_2) = x_{(2,1),(4,2)}.$$

We now present a proof that the Jones basis spans, adding slightly more detail than we found in the references. The extra detail will be used in the next section.

Definition 2.4. Given a word $w = U_{i_1} \dots U_{i_n}$ in the U_i , define the *terminus* to be the subscript of the final letter of the word appearing, i_n , and denote it $t(w)$. Set $t(1) = \infty$ as a convention. Define the *index* of w to be the minimum subscript i_j appearing, and denote it $i(w)$.

Remark 2.5. Note here that the notions of Jones normal form and index in $TL_n(a)$ coincide with those of [KT08], under the bijection which sends the generator e_i of [KT08] to the generator U_{n-i} used in this paper, for $1 \leq i \leq n - 1$.

The following two lemmas are an enhancement of Lemmas 5.25 and 5.26 of [KT08].

Lemma 2.6. *Any word $w \in \text{TL}_n(a)$ is equal in $\text{TL}_n(a)$ to a scalar multiple of a word w' in which*

- (a) $i(w) = i(w')$ and $U_{i(w)}$ appears exactly once in w' ;
- (b) $t(w') = t(w)$.

Point (a) occurs as Lemma 5.25 of [KT08], and the following is a simple extension of the proof that appears there. We have opted to give our proof in full because, as well as the minor extension of the proof, our notation differs from that of [KT08] as in Remark 2.5.

Proof. We proceed by reverse induction on the index $i(w)$ of w , which lies in the range $1 \leq i(w) \leq n-1$. If $i(w) = n-1$, then $w = U_{n-1}^i$ for some $i \geq 1$, so that $w = a^{i-1}U_{n-1}$ is a scalar multiple of the word U_{n-1} . Since the words U_{n-1}^i and U_{n-1} have the same index and terminus, the result holds in this case.

Suppose that the claim holds for all words of index $> p$ and let w be a nonempty word of index p . Suppose that U_p appears in w at least twice. Then we may write $w = w_1U_pw'U_pw_2$ where $i(w') = \ell > p$.

If $\ell > p+1$, then all letters of w' commute with U_p , so that

$$w = w_1U_pw'U_pw_2 = w_1w'U_p^2w_2 = aw_1w'U_pw_2.$$

Thus we have reduced the number of occurrences of U_p in w while preserving the (nonempty) final portion U_pw_2 of the word, so that the terminus remains unchanged.

If $\ell = p+1$, then by the induction hypothesis we may assume that U_{p+1} appears only once in w' , so that $w' = w_3U_{p+1}w_4$ where w_3, w_4 are words of index $\geq p+2$. Therefore w_3, w_4 commute with U_p , and consequently

$$\begin{aligned} w &= w_1U_pw'U_pw_2 = w_1U_pw_3U_{p+1}w_4U_pw_2 \\ &= w_1w_3U_pU_{p+1}U_pw_4w_2 = w_1w_3U_pw_4w_2 = w_1w_3w_4U_pw_2. \end{aligned}$$

So again, we have reduced the number of occurrences of U_p in the word while preserving the final (nonempty) portion U_pw_2 , and in particular preserving the terminus.

Repeating the process of reducing the number of occurrences of U_p while preserving the terminus, we find that w is a scalar multiple of a word w' of the required form. \square

Lemma 2.7. *Any word $w \in \text{TL}_n(a)$ is equivalent in $\text{TL}_n(a)$ to a scalar multiple of a word w' such that*

- (a) w' is written in Jones normal form;
- (b) $t(w') \leq t(w)$;
- (c) if $t(w') < t(w)$ then $t(w') \leq t(w) - 2$.

Proof. As in the previous Lemma, point (a) occurs as [KT08, Lemma 5.26]. We refer the reader to that proof, with the following modifications:

- Invoke the bijection of generators of Remark 2.5. This amounts to replacing each occurrence of e_i with U_{n-i} , so for example the subscripts 1 and $n - 1$ are interchanged, and inequalities are ‘reversed’.
- Whenever the inductive hypothesis is used in [KT08, Lemma 5.25], instead use the statement of the present lemma as a stronger inductive hypothesis.
- At the point where [KT08, Lemma 5.25] is used in [KT08, Lemma 5.26], use instead Lemma 2.6.

With these modifications in place, one can simply observe how the terminus changes in the proof of [KT08, Lemma 5.26], to obtain the present strengthening of that result. \square

2.2. Induced modules of Temperley-Lieb Algebras.


Definition 2.8 (The trivial module $\mathbb{1}$). The *trivial* module $\mathbb{1}$ of the Temperley-Lieb algebra $\text{TL}_n(a)$ is the module consisting of R with the action of $\text{TL}_n(a)$ in which all of the generators U_1, \dots, U_{n-1} act as 0. We can regard $\mathbb{1}$ as either a left or right module over $\text{TL}_n(a)$, and we will usually do that without indicating so in the notation.

Definition 2.9 (Sub-algebra convention). For $m \leq n$, we will regard $\text{TL}_m(a)$ as the sub-algebra of $\text{TL}_n(a)$ generated by the elements U_1, \dots, U_{m-1} . We will often regard $\text{TL}_n(a)$ as a left $\text{TL}_n(a)$ -module and a right $\text{TL}_m(a)$ -module, so that we obtain the left $\text{TL}_n(a)$ -module $\text{TL}_n(a) \otimes_{\text{TL}_m(a)} \mathbb{1}$.

Remark 2.10. Elements of $\text{TL}_n(a) \otimes_{\text{TL}_m(a)} \mathbb{1}$ can always be written as elementary tensors of the form $y \otimes 1$, since in this module $x \otimes r = rx \otimes 1$ for all $r \in R$.

The modules $\text{TL}_n \otimes_{\text{TL}_m} \mathbb{1}$ are an essential ingredient in the rest of this paper: they will be the building blocks of all the complexes we construct in order to prove our main results, in particular the complex of planar injective words $W(n)$. The rest of this section will study them in some detail, in particular finding a basis for them analogous to the Jones basis.

Remark 2.11 ($\text{TL}_n(a) \otimes_{\text{TL}_m(a)} \mathbb{1}$ via diagrams). The elements of $\text{TL}_n(a) \otimes_{\text{TL}_m(a)} \mathbb{1}$ can be regarded as diagrams, just like the elements of $\text{TL}_n(a)$, except that now the first m dots on the right are encapsulated within a *black box*, and if any cups can be absorbed into the black box, then the diagram is identified with 0. For example, some elements of $\text{TL}_4(a) \otimes_{\text{TL}_3(a)} \mathbb{1}$ are depicted as follows:



The structure of $\mathrm{TL}_n(a) \otimes_{\mathrm{TL}_m(a)} \mathbb{1}$ as a left module for $\mathrm{TL}_n(a)$ is given by pasting diagrams on the left, and then simplifying, as in the following example for $n = 4$ and $m = 2$:

Definition 2.12 (The ideal I_m). Given $0 \leq m \leq n$, let I_m denote the left ideal of $\mathrm{TL}_n(a)$ generated by the elements U_1, \dots, U_{m-1} .

Lemma 2.13. $\mathrm{TL}_n(a) \otimes_{\mathrm{TL}_m(a)} \mathbb{1}$ and $\mathrm{TL}_n(a)/I_m$ are isomorphic as left $\mathrm{TL}_n(a)$ -modules via the maps

$$\mathrm{TL}_n(a) \otimes_{\mathrm{TL}_m(a)} \mathbb{1} \longrightarrow \mathrm{TL}_n(a)/I_m, \quad y \otimes r \longmapsto yr + I_m$$

and

$$\mathrm{TL}_n(a)/I_m \longrightarrow \mathrm{TL}_n(a) \otimes_{\mathrm{TL}_m(a)} \mathbb{1}, \quad y + I_m \longmapsto y \otimes 1.$$

Proof. Observe that the generators U_1, \dots, U_{m-1} of the left-ideal I_m in TL_n are precisely the generators of the subalgebra TL_m of TL_n . Thus the map $y \otimes r \mapsto yr + I_m$ is well defined because if $i = 1, \dots, m-1$ then elements of the form $yU_i \otimes r$ and $y \otimes U_i r$ both map to 0 in TL_n/I_m . And $y + I_m \mapsto y \otimes 1$ is well defined because elements of I_m are linear combinations of ones of the form $x \cdot U_i$ for $i = 1, \dots, m-1$, and $(x \cdot U_i) \otimes 1 = x \otimes (U_i \cdot 1) = x \otimes 0 = 0$ for $i = 1, \dots, m-1$. One can now check that the two maps are inverses of one another. \square

Remark 2.14. Lemma 2.13 justifies the description of $\mathrm{TL}_n(a) \otimes_{\mathrm{TL}_m(a)} \mathbb{1}$ in terms of diagrams with ‘black boxes’ that we gave in Remark 2.11. Indeed, I_m is precisely the span of those diagrams which have a cup on the right between the dots i and $i+1$ for some $i = 1, \dots, m-1$. But these are precisely the diagrams which are made to vanish by having a cup fall into the black box. Thus $\mathrm{TL}_n(a)/I_m$ has basis given by the remaining diagrams, i.e. the ones that are not rendered 0 by the black box.

Lemma 2.15. For $m \leq n$, the ideal I_m of $\mathrm{TL}_n(a)$ has basis consisting of those elements of $\mathrm{TL}_n(a)$ written in Jones normal form $x_{\underline{a}, \underline{b}}$, which have terminus $b_1 \leq m-1$ (and $k \neq 0$).

Proof. Recall that words of the form $x_{\underline{a}, \underline{b}}$ give a basis for TL_n . Then by definition any word $w \in I_m$ is of the form $w = x_{\underline{a}, \underline{b}} v$ for $v \in \langle U_1, \dots, U_{m-1} \rangle$ and $v \neq e$. Then we have that $t(w) \leq m-1$. Now apply Lemma 2.7 to w to complete the proof. \square

Lemma 2.16. For $m \leq n$, $\mathrm{TL}_n(a) \otimes_{\mathrm{TL}_m(a)} \mathbb{1}$ has basis given by $x_{\underline{a}, \underline{b}} \otimes \mathbb{1}$ such that the terminus $b_1 > m-1$.

Proof. From Lemma 2.13 $\mathrm{TL}_n \otimes_{\mathrm{TL}_m} \mathbb{1}$ is isomorphic to TL_n/I_m . Then elements of the form $x_{\underline{a},\underline{b}}$ give a basis for TL_n and elements of the form $x_{\underline{a},\underline{b}}$, which have terminus $b_1 \leq m-1$ give a basis for I_m by Lemma 2.15. Therefore a basis for the quotient is given by $x_{\underline{a},\underline{b}}$ such that the terminus $b_1 > m-1$, and under the isomorphism in Lemma 2.13 this gives the required basis. \square

Example 2.17. The Jones basis of $\mathrm{TL}_3(a)$ is:

$$1, \quad U_2, \quad U_1U_2, \quad U_1, \quad U_2U_1$$

So $\mathrm{TL}_3(a) \otimes_{\mathrm{TL}_2(a)} \mathbb{1}$ has basis consisting of those elements whose terminus is strictly greater than 1, namely:

$$1, \quad U_2, \quad U_1U_2.$$

(Recall that by convention the terminus of 1 is ∞ .)

Lemma 2.18. *For $m \leq n$, suppose that $y \in \mathrm{TL}_n(a)$ and that $y \cdot U_{m-1}$ lies in I_{m-1} . Then $y \cdot U_{m-1}$ lies in I_{m-2} .*

Proof. The product $y \cdot U_{m-1}$ is a linear combination of words ending with U_{m-1} , i.e. of words w with $t(w) = m-1$. By Lemma 2.7, this can be rewritten as a linear combination of Jones basis elements $x_{\underline{a},\underline{b}}$ whose terminus satisfies $t(x_{\underline{a},\underline{b}}) = m-1$ or $t(x_{\underline{a},\underline{b}}) \leq m-3$. Since $y \cdot U_{m-1} \in I_{m-1}$, this means that in fact no basis elements with terminus $m-1$ remain after cancellation, and therefore all remaining words have terminus $m-3$ or less, and so lie in I_{m-2} . \square

2.3. Iwahori-Hecke algebras.

Definition 2.19 (The Iwahori-Hecke algebra). Let $n \geq 0$ and let $q \in R^\times$. The *Iwahori-Hecke algebra* $\mathcal{H}_n(q)$ of type A_{n-1} is the algebra with generators

$$T_1, \dots, T_{n-1}$$

satisfying the following relations:

- $T_i T_j = T_j T_i$ for $i \neq j \pm 1$
- $T_i T_j T_i = T_j T_i T_j$ for $i = j \pm 1$
- $T_i^2 = (q-1)T_i + q$

Definition 2.20 (From Iwahori-Hecke to Temperley-Lieb). Now suppose that there is $v \in R^\times$ such that $q = v^2$. Then there are two natural homomorphisms

$$\theta_1, \theta_2: \mathcal{H}_n(q) \longrightarrow \mathrm{TL}_n(v + v^{-1}),$$

defined by $\theta_1(T_i) = vU_i - 1$ and $\theta_2(T_i) = v^2 - vU_i$ for $i = 1, \dots, n-1$. They induce isomorphisms

$$\bar{\theta}_1: \mathcal{H}_n(q)/I_1 \xrightarrow{\cong} \mathrm{TL}_n(v + v^{-1}), \quad \bar{\theta}_2: \mathcal{H}_n(q)/I_2 \xrightarrow{\cong} \mathrm{TL}_n(v + v^{-1}),$$

where I_1 is the two-sided ideal generated by elements of the form

$$T_i T_j T_i + T_i T_j + T_j T_i + T_i + T_j + 1$$

for $i = j \pm 1$, and I_2 is the two-sided ideal generated by elements of the form

$$T_i T_j T_i - q T_i T_j - q T_j T_i + q^2 T_i + q^2 T_j - q^3$$

for $i = j \pm 1$. See [FG97], Theorem 5.29 of [KT08], and Section 2.3 of [HMR09], though unfortunately conventions change from author to author. Another standard convention of setting $a = -(v+v^{-1})$ can easily be accounted for by swapping v with $-v^{\pm 1}$.

We will take an agnostic approach to the homomorphisms θ_1, θ_2 . We will choose one of them and denote it by simply

$$\theta: \mathcal{H}_n(q) \longrightarrow \mathrm{TL}_n(v + v^{-1}),$$

and denote by λ the constant term in $\theta(T_i)$, and by μ the coefficient of U_i in $\theta(T_i)$, so that

$$\theta(T_i) = \lambda + \mu U_i.$$

Then θ induces an isomorphism

$$\bar{\theta}: \mathcal{H}_n(q)/I \xrightarrow{\cong} \mathrm{TL}_n(v + v^{-1})$$

where I is the two-sided ideal generated by elements of the form

$$T_i T_j T_i - \lambda T_i T_j - \lambda T_j T_i + \lambda^2 T_i + \lambda^2 T_j - \lambda^3$$

for $i = j \pm 1$. And moreover, the elements $\theta(T_i)$ act on the trivial module $\mathbb{1}$ as multiplication by λ .

Definition 2.21. Let $v \in R^\times$. We define $s_1, \dots, s_{n-1} \in \mathrm{TL}_n(v + v^{-1})$ by setting

$$s_i = \theta(T_i) = \lambda + \mu U_i$$

and note that these elements satisfy the following properties:

- $s_i^2 = (v^2 - 1)s_i + v^2$ for all i ,
- $s_i s_j = s_j s_i$ for $i \neq j \pm 1$,
- $s_i s_j s_i = s_j s_i s_j$ for $i = j \pm 1$,
- $s_i s_j s_i - \lambda s_i s_j - \lambda s_j s_i + \lambda^2 s_i + \lambda^2 s_j - \lambda^3 = 0$ for $i = j \pm 1$,
- s_i acts on $\mathbb{1}$ as multiplication by λ .

Remark 2.22. There is a homomorphism from (the group algebra of) the braid group into $\mathrm{TL}_n(v + v^{-1})$ given on generators by $s_i \mapsto s_i$. This is the content of the second and third bullet points above, together with the fact that the s_i are invertible, which follows from the first bullet point (and the fact that v is a unit). Diagrammatically, this homomorphism can be viewed as a *smoothing expansion* from braided diagrams to planar diagrams: take a braid diagram, and then smooth each crossing in turn in the two possible ways, using appropriate weightings for each smoothing. For example, we can visualise the image of s_i in $\mathrm{TL}_n(v + v^{-1})$ as the standard braid group generator crossing strand i over strand $i+1$. There are two ways this crossing can be resolved to a planar diagram, and we equate s_i to the sum of these two states. They are the identity and U_i , as




FIGURE 2. Smoothings of s_i

shown in Figure 2. The coefficient of the identity is λ and the coefficient of U_i is μ , simply because we defined $s_i = \lambda + \mu U_i$. Similarly, we consider the image of s_i^{-1} as strand i crossing under strand $i + 1$, and when this is smoothed the coefficient of the identity is λ^{-1} and the coefficient of U_i is μ^{-1} , precisely because one can verify that $s_i^{-1} = \lambda^{-1} + \mu^{-1}U_i$ in $\text{TL}_n(v + v^{-1})$.

In principle we could describe how various Reidemeister moves affect the smoothing expansion, but it will not be necessary for the rest of the paper. Moreover, we will only encounter positive powers of s_i .

3. INDUCTIVE RESOLUTIONS

In this section we prove the following two theorems, which we recall from the introduction.

Theorem A. *Let R be a commutative ring and let a be a unit in R . Then $\text{Tor}_d^{\text{TL}_n(a)}(\mathbb{1}, \mathbb{1})$ and $\text{Ext}_{\text{TL}_n(a)}^d(\mathbb{1}, \mathbb{1})$ both vanish for $d > 0$.*

Theorem F. *Let R be a commutative ring and let $a \in R$. Let $0 \leq m < n$. Then $\text{Tor}_d^{\text{TL}_n(a)}(\mathbb{1}, \text{TL}_n(a) \otimes_{\text{TL}_m(a)} \mathbb{1})$ and $\text{Ext}_{\text{TL}_n(a)}^d(\text{TL}_n(a) \otimes_{\text{TL}_m(a)} \mathbb{1}, \mathbb{1})$ vanish for $d > 0$.*

In fact for Theorem A we will prove the following stronger claim:

Claim 3.1. *Suppose that the parameter $a \in R$ is invertible. Then for any $0 \leq m \leq n$, the groups $\text{Tor}_d^{\text{TL}_n(a)}(\mathbb{1}, \text{TL}_n(a) \otimes_{\text{TL}_m(a)} \mathbb{1})$ and $\text{Ext}_d^{\text{TL}_n(a)}(\text{TL}_n(a) \otimes_{\text{TL}_m(a)} \mathbb{1}, \mathbb{1})$ both vanish for $d > 0$.*

The similarity between Theorem F and Claim 3.1 is now clear. Both will be proved by induction on m , the initial cases $m = 0, 1$ being immediate because then TL_m is the ground ring R so that $\text{TL}_n \otimes_{\text{TL}_m} \mathbb{1} \cong \text{TL}_n$ is free. In order to produce an inductive proof, we construct resolutions of the modules $\text{TL}_n \otimes_{\text{TL}_m} \mathbb{1}$ whose terms are not free or projective or injective, but instead whose terms are the modules considered earlier in the induction, specifically $\text{TL}_n \otimes_{\text{TL}_{m-1}} \mathbb{1}$ and $\text{TL}_n \otimes_{\text{TL}_{m-2}} \mathbb{1}$. For this reason we refer to these resolutions as *inductive*

$$\begin{array}{ccc}
& \vdots & \\
& \downarrow (1-a^{-1}U_{m-1}) & \\
\mathrm{TL}_n \otimes_{\mathrm{TL}_{m-2}} \mathbb{1} & & 3 \\
& \downarrow a^{-1}U_{m-1} & \\
\mathrm{TL}_n \otimes_{\mathrm{TL}_{m-2}} \mathbb{1} & & 2 \\
& \downarrow (1-a^{-1}U_{m-1}) & \\
\mathrm{TL}_n \otimes_{\mathrm{TL}_{m-2}} \mathbb{1} & & 1 \\
& \downarrow a^{-1}U_{m-1} & \\
\mathrm{TL}_n \otimes_{\mathrm{TL}_{m-1}} \mathbb{1} & & 0 \\
& \downarrow 1 & \\
\mathrm{TL}_n \otimes_{\mathrm{TL}_m} \mathbb{1} & & -1
\end{array}$$

FIGURE 3. The complex $C(m)$.

resolutions. This approach is inspired by homological stability arguments, in which one considers complexes whose building blocks are induced up from the earlier objects in the sequence. The difference here is that our complexes are actual resolutions — they are acyclic rather than just acyclic up to a point — and because Shapiro’s lemma is unavailable we do not change the algebra we are working over, rather we change the algebra from which we are inducing our modules.

3.1. The inductive resolutions. In this subsection we establish the resolutions $C(m)$ and $D(m)$ of $\mathrm{TL}_n \otimes_{\mathrm{TL}_m} \mathbb{1}$ required to prove Claim 3.1 and Theorem F above.

Definition 3.2 (The complex $C(m)$). Let $2 \leq m \leq n$ and assume that a is invertible. We define a chain complex of left $\mathrm{TL}_n(a)$ -modules as in Figure 3. The degree is indicated in the right-hand column. The differentials of $C(m)$ are all given by extending the algebra over which the tensor product is taken, by right multiplying in the first factor by the indicated element of $\mathrm{TL}_n(a)$, or by a combination of the two. So, for example, the differential originating in degree 1 sends $x \otimes r \in \mathrm{TL}_n(a) \otimes_{\mathrm{TL}_{m-2}(a)} \mathbb{1}$ to $(x \cdot a^{-1}U_{m-1}) \otimes r \in \mathrm{TL}_n(a) \otimes_{\mathrm{TL}_{m-1}(a)} \mathbb{1}$. The complex is periodic of period 2 in degrees 1 and above, so that all entries are $\mathrm{TL}_n(a) \otimes_{\mathrm{TL}_{m-2}(a)} \mathbb{1}$ and the boundary maps between them alternate between $a^{-1}U_{m-1}$ and $(1 - a^{-1}U_{m-1})$. The boundary maps are well defined because U_{m-1} commutes inside $\mathrm{TL}_n(a)$ with all elements of $\mathrm{TL}_{m-2}(a)$.

$$\begin{array}{ccc}
& \vdots & \\
& \downarrow (1-U_{m-1}U_m) & \\
\mathrm{TL}_n \otimes_{\mathrm{TL}_{m-2}} \mathbb{1} & & 3 \\
& \downarrow U_{m-1}U_m & \\
\mathrm{TL}_n \otimes_{\mathrm{TL}_{m-2}} \mathbb{1} & & 2 \\
& \downarrow (1-U_{m-1}U_m) & \\
\mathrm{TL}_n \otimes_{\mathrm{TL}_{m-2}} \mathbb{1} & & 1 \\
& \downarrow U_{m-1} & \\
\mathrm{TL}_n \otimes_{\mathrm{TL}_{m-1}} \mathbb{1} & & 0 \\
& \downarrow 1 & \\
\mathrm{TL}_n \otimes_{\mathrm{TL}_m} \mathbb{1} & & -1
\end{array}$$

FIGURE 4. The complex $D(m)$.

Definition 3.3 (The complex $D(m)$). Let $2 \leq m < n$, and do not assume that a is invertible. We define a chain complex of left $\mathrm{TL}_n(a)$ -modules as in Figure 4. The degree is indicated in the right-hand column. The differentials of $D(m)$ are all given by extending the algebra over which the tensor product is taken, by right multiplying in the first factor by the indicated element of $\mathrm{TL}_n(a)$, or by a combination of the two. So, for example, the differential originating in degree 1 sends $x \otimes r \in \mathrm{TL}_n(a) \otimes_{\mathrm{TL}_{m-2}(a)} \mathbb{1}$ to $x \cdot U_{m-1} \otimes r \in \mathrm{TL}_n(a) \otimes_{\mathrm{TL}_{m-1}(a)} \mathbb{1}$. The complex is periodic of period 2 in degrees 1 and above, so that in that range all terms are $\mathrm{TL}_n(a) \otimes_{\mathrm{TL}_{m-2}(a)} \mathbb{1}$ and the boundary maps between them alternate between $U_{m-1}U_m$ and $(1 - U_{m-1}U_m)$. The boundary maps are well defined because U_{m-1} and $U_{m-1}U_m$ commute inside $\mathrm{TL}_n(a)$ with all elements of $\mathrm{TL}_{m-2}(a)$. Observe that the condition $m < n$ is necessary in order to ensure that U_m is actually an element of $\mathrm{TL}_n(a)$.

Lemma 3.4.

- (1) Let $2 \leq m \leq n$ and let a be invertible. Then $a^{-1}U_{m-1} \in \mathrm{TL}_n(a)$ is idempotent.
- (2) Let $2 \leq m < n$ and let a be arbitrary. Then $U_{m-1}U_m \in \mathrm{TL}_n(a)$ is idempotent.

Proof. We calculate

$$(a^{-1}U_i)^2 = a^{-2}U_i^2 = a^{-2}aU_i = a^{-1}U_i$$

and

$$U_{m-1}U_m \cdot U_{m-1}U_m = U_{m-1}U_mU_{m-1} \cdot U_m = U_{m-1}U_m. \quad \square$$

From now on in this section, we will attempt to talk about $C(m)$ and $D(m)$ at the same time. When we refer to $C(m)$, the relevant assumptions should be understood, namely that $2 \leq m \leq n$ and that $a \in R$ is a unit. And when we refer to $D(m)$, the assumptions that $2 \leq m < n$ but $a \in R$ is arbitrary should be understood. We trust that this will not be confusing.

Lemma 3.5. *$C(m)$ and $D(m)$ are indeed chain complexes.*

Proof. We give the proof for $C(m)$. The proof for $D(m)$ is similar. We must check that consecutive boundary maps of $C(m)$ compose to 0. In the case of the composite from degree 1 to -1 , the composition is given by

$$x \otimes r \mapsto (x \cdot a^{-1}U_{m-1}) \otimes r = x \otimes (a^{-1}U_{m-1} \cdot r) = x \otimes 0 = 0$$

this holds because the tensor product is over TL_m , which contains $a^{-1}U_{m-1}$. In the case of the remaining composites, this follows immediately from

$$(a^{-1}U_{m-1}) \cdot (1 - a^{-1}U_{m-1}) = 0 = (1 - a^{-1}U_{m-1}) \cdot (a^{-1}U_{m-1}),$$

which is a consequence of the fact that $a^{-1}U_{m-1}$ is idempotent (from Lemma 3.4). \square

Lemma 3.6. *The complexes $C(m)$ and $D(m)$ are acyclic.*

Proof. In degree -1 it is clear that the boundary map is surjective, for both $C(m)$ and $D(m)$.

In degree 0, we will give the proof for $C(m)$, the proof for $D(m)$ being similar. Suppose that $y \otimes 1 \in \mathrm{TL}_n \otimes_{\mathrm{TL}_{m-1}} \mathbb{1}$ lies in the kernel of the boundary map, or in other words that $y \otimes 1 \in \mathrm{TL}_n \otimes_{\mathrm{TL}_m} \mathbb{1}$ vanishes. This means that y lies in the left-ideal generated by the elements U_1, \dots, U_{m-1} . Since all but the last of these generators lie in TL_{m-1} , and we started with $y \otimes 1 \in \mathrm{TL}_n \otimes_{\mathrm{TL}_{m-1}} \mathbb{1}$, we may assume without loss that $y = y' \cdot U_{m-1}$ for some y' . But then

$$y \otimes 1 = y' \cdot U_{m-1} \otimes 1 = ay' \cdot (a^{-1}U_{m-1}) \otimes 1$$

does indeed lie in the image of the boundary map.

In degree 1, we give the proof for both complexes. First, for $C(m)$, suppose that $y \otimes 1 \in \mathrm{TL}_n \otimes_{\mathrm{TL}_{m-2}} \mathbb{1}$ lies in the kernel of the boundary map. It follows that $y \cdot (a^{-1}U_{m-1}) \otimes 1$ vanishes in $\mathrm{TL}_n \otimes_{\mathrm{TL}_{m-1}} \mathbb{1}$, which means that $y \cdot (a^{-1}U_{m-1})$ lies in the left ideal I_{m-1} generated by U_1, \dots, U_{m-2} . It follows from Lemma 2.18 that $y \cdot (a^{-1}U_{m-1})$ lies in the left ideal I_{m-2} generated by U_1, \dots, U_{m-3} , so that in $\mathrm{TL}_n \otimes_{\mathrm{TL}_{m-2}} \mathbb{1}$ the element $y \cdot (a^{-1}U_{m-1}) \otimes 1$ vanishes. Thus

$$y \otimes 1 = y \cdot (1 - a^{-1}U_{m-1}) \otimes 1$$

does indeed lie in the image of the boundary map. Second, for $D(m)$, suppose that $y \otimes 1 \in \mathrm{TL}_n \otimes_{\mathrm{TL}_{m-2}} \mathbb{1}$ lies in the kernel of the boundary map. Then, as




FIGURE 5. The complex $\mathbb{1} \otimes C(m)$

for $C(m)$, $y \cdot U_{m-1}$ lies in I_{m-2} . So $y \cdot U_{m-1}U_m$ also lies in the left ideal I_{m-2} since U_m commutes with the generators of I_{m-2} . Thus $y \cdot U_{m-1}U_m \otimes 1$ vanishes in $\text{TL}_n \otimes_{\text{TL}_{m-2}} \mathbb{1}$, so that $y \otimes 1 = y \cdot (1 - U_{m-1}U_m) \otimes 1$ does indeed lie in the image of the boundary map.

In degrees 2 and higher, acyclicity is an immediate consequence of the fact that $a^{-1}U_{m-1}$ and $U_{m-1}U_m$ are idempotents, by Lemma 3.4. \square

Lemma 3.7. *The complexes $\mathbb{1} \otimes_{\text{TL}_n(a)} C(m)$, $\mathbb{1} \otimes_{\text{TL}_n(a)} D(m)$, $\text{Hom}_{\text{TL}_n(a)}(C(m), \mathbb{1})$ and $\text{Hom}_{\text{TL}_n(a)}(D(m), \mathbb{1})$ are acyclic.*


Proof. We give the proof for $\mathbb{1} \otimes_{\text{TL}_n} C(m)$, the proof for the other parts being similar. The terms of $C(m)$ have the form $\text{TL}_n \otimes_{\text{TL}_{m-i}} \mathbb{1}$ where $i = 0, 1, 2$ depending on the degree. Thus $\mathbb{1} \otimes_{\text{TL}_n} C(m)$ has terms of the form $\mathbb{1} \otimes_{\text{TL}_n} (\text{TL}_n \otimes_{\text{TL}_{m-i}} \mathbb{1}) \cong \mathbb{1} \otimes_{\text{TL}_{m-i}} \mathbb{1} \cong \mathbb{1}$. Moreover, by tracing through this isomorphism, one sees that if a boundary map in $C(m)$ is labelled by an element $x \in \text{TL}_n$, then the corresponding boundary map in $\mathbb{1} \otimes_{\text{TL}_n} C(m)$ is simply the map $\mathbb{1} \rightarrow \mathbb{1}$ given by the action of x on $\mathbb{1}$. Thus $\mathbb{1} \otimes_{\text{TL}_n} C(m)$ is nothing other than the complex in Figure 5. (The right hand column indicates the degree.) This is visibly acyclic, and this completes the proof. \square

Remark 3.8 (Representation theory and the inductive resolutions). Schur-Weyl duality relates representations of TL_n with representations of the quantum group $U_q(\mathfrak{sl}_2)$, and it is possible to use this to construct our inductive resolutions via the representation theory of $U_q(\mathfrak{sl}_2)$. We will try to describe this briefly. We are indebted to an anonymous referee for explaining this connection to us.


One instance of Schur-Weyl duality is the following. Let V denote the standard representation of $U_q(\mathfrak{sl}_2)$. Then there is an isomorphism $\mathrm{TL}_n \cong \mathrm{End}_{U_q(\mathfrak{sl}_2)}(V^{\otimes n})$, and more generally there are isomorphisms $\mathrm{TL}(n, m) \cong \mathrm{Hom}_{U_q(\mathfrak{sl}_2)}(V^{\otimes n}, V^{\otimes m})$ that assemble into a monoidal functor on the *Temperley-Lieb category* TL . (The objects of TL are the non-negative integers, the morphism space $\mathrm{TL}(n, m)$ is the R -module spanned by planar diagrams with n marked points on the left and m marked points on the right, and composition is defined just like multiplication in TL_n .) See Webster’s appendix to [EL17].

One can write down exact sequences of $U_q(\mathfrak{sl}_2)$ -modules that, after applying Schur-Weyl duality, yield the inductive resolutions $C(m)$ and $D(m)$. We will not detail the construction of these sequences, except to say that each one relies on the construction of an appropriate splitting of some tensor power of V . The relevant splittings are constructed in each case as follows:

- In the case where a is invertible, the morphisms




in TL compose to give the identity morphism in $\mathrm{TL}(0, 0)$. (The two semicircles compose to the circle morphism from 0 to itself, and by the usual rule for composing diagrams, the circle morphism is a times the identity.) This then corresponds to a pair of maps $R = V^{\otimes 0} \rightarrow V^{\otimes 2}$ and $V^{\otimes 2} \rightarrow V^{\otimes 0} = R$ that compose to the identity, showing that $V^{\otimes 2}$ splits off a copy of R . Note that the map $V^{\otimes 2} \rightarrow V^{\otimes 2}$ that projects onto this copy of R is represented by the morphism




in TL . Compare this with the idempotent $a^{-1}U_{m-1}$ appearing in $C(n)$.

- When a is not invertible, we consider the morphisms



which compose to give the identity morphism in $\mathrm{TL}(1, 1)$. These diagrams correspond to a pair of maps $V \rightarrow V^{\otimes 3} \rightarrow V$ that compose to the identity, showing that $V^{\otimes 3}$ splits off a copy of V . Observe that the map $V^{\otimes 3} \rightarrow V^{\otimes 3}$ that projects to this copy of V is represented by the morphism



which can be compared to the idempotent $U_{m-1}U_m$ appearing in $D(n)$.

3.2. The spectral sequence of a double complex. Since the spectral sequence of a particular kind of double complex is used several times during this paper, we introduce and discuss it in this subsection.

We begin with the homological version. Suppose we have a chain complex Q_* of left TL_n -modules, such as $C(m)$ or $D(m)$, or the complex of planar injective words $W(n)$ to be introduced later. Then we choose a projective resolution P of $\mathbb{1}$ as a right module over TL_n , and we consider the double complex $P_* \otimes_{\mathrm{TL}_n} Q_*$. This is a homological double complex in the sense that both differentials reduce the grading. Associated to this double complex are two spectral sequences, $\{^I E^r\}$ and $\{^{II} E^r\}$, which both converge to the homology of the totalisation, $H_*(\mathrm{Tot}(P_* \otimes_{\mathrm{TL}_n} Q_*))$ as in Section 5.6 of [Wei94]. The first spectral sequence has E^1 -term given by $^I E_{i,j}^1 = H_j(P_i \otimes_{\mathrm{TL}_n} Q_*) \cong P_i \otimes_{\mathrm{TL}_n} H_j(Q_*)$ with $d^1: ^I E_{i,j}^1 \rightarrow ^I E_{i-1,j}^1$ induced by the differential $P_i \rightarrow P_{i-1}$. The isomorphism above holds because each P_i is projective and therefore flat. It follows that the E^2 -term is

$$^I E_{i,j}^2 = \mathrm{Tor}_i^{\mathrm{TL}_n}(\mathbb{1}, H_j(Q_*)).$$

The second spectral sequence has E^1 -term given by $^{II} E_{i,j}^1 = H_j(P_* \otimes_{\mathrm{TL}_n} Q_i)$, i.e.

$$^{II} E_{i,j}^1 = \mathrm{Tor}_j^{\mathrm{TL}_n}(\mathbb{1}, Q_i)$$

with $d^1: ^{II} E_{i,j}^1 \rightarrow ^{II} E_{i-1,j}^1$ induced by the boundary maps of Q_* .

We now consider the cohomological version. Suppose we have a chain complex Q_* of left TL_n -modules, again such as $C(m)$, $D(m)$ or $W(n)$ (the latter to be introduced later). Then we choose an injective resolution I^* of $\mathbb{1}$ as a left module over TL_n , and we consider the double complex $\mathrm{Hom}_{\mathrm{TL}_n}(Q_*, I^*)$. This is a cohomological double complex in the sense that both differentials increase the grading. Associated to this double complex are two spectral sequences, $\{^I E_r\}$ and $\{^{II} E_r\}$, both converging to the cohomology of the totalisation, $H^*(\mathrm{Tot}(\mathrm{Hom}_{\mathrm{TL}_n}(Q_*, I^*)))$ as in Section 5.6 of [Wei94]. The first spectral sequence has E_1 -term given by $^I E_1^{i,j} = H^j(\mathrm{Hom}_{\mathrm{TL}_n}(Q_*, I^i)) \cong \mathrm{Hom}_{\mathrm{TL}_n}(H_j(Q_*), I^i)$ with $d^1: ^I E_1^{i,j} \rightarrow ^I E_1^{i+1,j}$ induced by the differential of I^* . The isomorphism above holds because each I^i is injective, so that the functor $\mathrm{Hom}_{\mathrm{TL}_n}(-, I^i)$ is exact. It follows that the E_2 -term is

$$^I E_2^{i,j} = \mathrm{Ext}_{\mathrm{TL}_n}^i(H_j(Q_*), \mathbb{1}).$$

The second spectral sequence has E_1 -term $^{II} E_1^{i,j} = H^j(\mathrm{Hom}_{\mathrm{TL}_n}(Q_i, I^*))$, i.e.

$$^{II} E_1^{i,j} = \mathrm{Ext}_{\mathrm{TL}_n}^j(Q_i, \mathbb{1})$$

with differential $d^1: ^{II} E_1^{i,j} \rightarrow ^{II} E_1^{i+1,j}$ induced by the differential of Q_* .

3.3. Proof of Theorems A and F. We can now prove Claim 3.1 (which implies Theorem A) and Theorem F. The proofs of the two results will be almost identical except that the former uses the complex $C(m)$ and the latter uses the complex $D(m)$. Moreover, each result has a homological and cohomological part, referring to Tor and Ext respectively. In each case the two parts are proved similarly, by using either the homological or cohomological spectral sequence from Section 3.2 above. We will therefore only prove the homological part of Claim 3.1, i.e. we will prove that $\mathrm{Tor}_*^{\mathrm{TL}_n}(\mathbb{1}, \mathrm{TL}_n \otimes_{\mathrm{TL}_m} \mathbb{1})$ vanishes in positive degrees, leaving the details of the other parts to the reader.

Proof of Claim 3.1, Tor case. We prove the claim by fixing n and using strong induction on m in the range $n \geq m \geq 0$. As noted before, the initial cases $m = 0, 1$ of the induction are immediate since then TL_m is the ground ring and $\mathrm{TL}_n \otimes_{\mathrm{TL}_m} \mathbb{1} \cong \mathrm{TL}_n$ is free. We therefore fix m in the range $2 \leq m \leq n$.

We now employ the homological spectral sequences $\{^I E^r\}$ and $\{^{II} E^r\}$ of Section 3.2, in the case $Q = C(m)$. Then $^I E_{i,j}^2 = \mathrm{Tor}_i^{\mathrm{TL}_n}(\mathbb{1}, H_j(C(m))) = 0$ for all i and j , since $C(m)$ is acyclic by Lemma 3.6. Thus $\{^I E^r\}$ converges to zero, and the same must therefore be true of $\{^{II} E^r\}$, since both spectral sequences have the same target. In the second spectral sequence the E^1 -page

$$^{II} E_{i,j}^1 = \mathrm{Tor}_j^{\mathrm{TL}_n}(\mathbb{1}, C(m)_i)$$

is largely known to us. The bottom $j = 0$ row of $^{II} E^1$ is precisely the complex $\mathbb{1} \otimes_{\mathrm{TL}_n} C(m)$, which is acyclic by Lemma 3.7. And when $i \geq 0$, the term $C(m)_i$ is either $\mathrm{TL}_n \otimes_{\mathrm{TL}_{m-1}} \mathbb{1}$ or $\mathrm{TL}_n \otimes_{\mathrm{TL}_{m-2}} \mathbb{1}$, and our inductive hypothesis applies to these ($(m-1) < m$ and $(m-2) < m$) to show that $^{II} E_{i,j}^1 = \mathrm{Tor}_j^{\mathrm{TL}_n}(\mathbb{1}, C(m)_i) = 0$ when $j > 0$. See Figure 6 for a visualisation of the E^1 page. Altogether, this tells us that $^{II} E_{i,j}^2$ vanishes except for the groups

$$^{II} E_{-1,j}^2 = \mathrm{Tor}_j^{\mathrm{TL}_n}(\mathbb{1}, C(m)_{-1}) = \mathrm{Tor}_j^{\mathrm{TL}_n}(\mathbb{1}, \mathrm{TL}_n \otimes_{\mathrm{TL}_m} \mathbb{1})$$

for $j > 0$, which are concentrated in a single column and therefore not subject to any further differentials. Thus $^{II} E^2 = ^{II} E^\infty$. But we know that $^{II} E^\infty$ vanishes identically, so that the inductive hypothesis is proved, and so, therefore, is the proof of the homological part of Claim 3.1. \square

4. PLANAR INJECTIVE WORDS

Throughout this section we will consider the Temperley-Lieb algebra $\mathrm{TL}_n(a) = \mathrm{TL}_n(v + v^{-1})$, where $v \in R^\times$. We will make use of the elements s_1, \dots, s_{n-1} of Definition 2.21.

Definition 4.1. For $n \geq 0$ we define a chain complex $W(n)_*$ of left $\mathrm{TL}_n(a)$ -modules as follows. For i in the range $-1 \leq i \leq n-1$, the degree- i part of $W(n)_*$ is defined by

$$W(n)_i = \mathrm{TL}_n(a) \otimes_{\mathrm{TL}_{n-i-1}(a)} \mathbb{1}$$




FIGURE 6. The page ${}^{II}E^1$. The only differentials that affect the ${}^{II}E^2$ page are shown on the $j = 0$ row.

and in all other degrees we set $W(n)_i = 0$. Note that

$$W(n)_{-1} = \mathrm{TL}_n(a) \otimes_{\mathrm{TL}_n(a)} \mathbb{1} = \mathbb{1}.$$

For $i \geq 0$ the boundary map $d^i: W(n)_i \rightarrow W(n)_{i-1}$ is defined to be the alternating sum $\sum_{j=0}^i (-1)^j d_j^i$, where $d_j^i: W(n)_i \rightarrow W(n)_{i-1}$ is the map

$$d_j^i: \mathrm{TL}_n(a) \otimes_{\mathrm{TL}_{n-i-1}(a)} \mathbb{1} \longrightarrow \mathrm{TL}_n(a) \otimes_{\mathrm{TL}_{n-i}(a)} \mathbb{1}$$

defined by

$$d_j^i(x \otimes r) = (x \cdot s_{n-i+j-1} \cdots s_{n-i}) \otimes \lambda^{-j} r.$$

In the expression $s_{n-i+j-1} \cdots s_{n-i}$, the indices decrease from left to right. Thus, for example, the product is $s_{n-i+1} s_{n-i}$ when $j = 2$, it is s_{n-i} when $j = 1$, and it is trivial (the unit element) when $j = 0$ (the latter point can be regarded as a convention if one wishes). Recall that λ is indeed invertible since $\lambda = -1$ or v^2 and v is a unit. For notational purposes we will write $W(n)$ and only use a subscript when identifying a particular degree.

Observe that d_j is well-defined because the elements $s_{n-i}, \dots, s_{n-i+j-1}$ all commute with all generators of TL_{n-i-1} . We have depicted $W(n)$ in Figure 7.

Lemma 4.2. *The boundary maps of $W(n)$ satisfy $d^{i-1} \circ d^i = 0$.*

Proof. We will show that if $i \geq 1$ and $0 \leq j < k \leq i$, then the composite maps $d_j^{i-1} d_k^i, d_{k-1}^{i-1} d_j^i: W(n)_i \rightarrow W(n)_{i-2}$ coincide. (Thus the d_j^i satisfy the semi-simplicial identities, so $W(n)$ is a semi-simplicial R -module.) The fact that $d \circ d$ vanishes then follows. We have

$$d_j^{i-1} d_k^i(x \otimes r) = [x \cdot (s_{n-i+k-1} \cdots s_{n-i}) \cdot (s_{n-i+j} \cdots s_{n-i+1})] \otimes \lambda^{-(j+k)} r$$

$$\begin{array}{ccc}
& \mathrm{TL}_n \otimes_{\mathrm{TL}_0} \mathbb{1} & n-1 \\
& \downarrow d_0^{n-1} - d_1^{n-1} + \dots + (-1)^{n-1} d_{n-1}^{n-1} & \\
& \mathrm{TL}_n \otimes_{\mathrm{TL}_1} \mathbb{1} & n-2 \\
& \downarrow d_0^{n-2} - d_1^{n-2} + \dots + (-1)^{n-2} d_{n-2}^{n-2} & \\
& \vdots & \\
& \downarrow & \\
& \mathrm{TL}_n \otimes_{\mathrm{TL}_{n-3}} \mathbb{1} & 2 \\
& \downarrow d_0^2 - d_1^2 + d_2^2 & \\
& \mathrm{TL}_n \otimes_{\mathrm{TL}_{n-2}} \mathbb{1} & 1 \\
& \downarrow d_0^1 - d_1^1 & \\
& \mathrm{TL}_n \otimes_{\mathrm{TL}_{n-1}} \mathbb{1} & 0 \\
& \downarrow & \\
& \mathbb{1} & -1
\end{array}$$

FIGURE 7. The complex $W(n)$.

and

$$d_{k-1}^{i-1} d_j^i(x \otimes r) = [x \cdot (s_{n-i+j-1} \cdots s_{n-i}) \cdot (s_{n-i+k-1} \cdots s_{n-i+1})] \otimes \lambda^{-(j+k-1)} r.$$

Now we have:

$$\begin{aligned}
& (s_{n-i+k-1} \cdots s_{n-i}) \cdot (s_{n-i+j} \cdots s_{n-i+1}) \\
&= (s_{n-i+j-1} \cdots s_{n-i}) \cdot (s_{n-i+k-1} \cdots s_{n-i}) \\
&= (s_{n-i+j-1} \cdots s_{n-i}) \cdot (s_{n-i+k-1} \cdots s_{n-i+1}) \cdot s_{n-i}
\end{aligned}$$

Here, the first equality follows by taking the letters of the second parenthesis in turn, and ‘passing through’ the first parenthesis, using a single braid relation, with the result that the letter’s index is reduced by 1. Thus:

$$\begin{aligned}
d_j^{i-1} d_k^i(x \otimes r) &= [x \cdot (s_{n-i+k-1} \cdots s_{n-i}) \cdot (s_{n-i+j} \cdots s_{n-i+1})] \otimes \lambda^{-(j+k)} r \\
&= [x \cdot (s_{n-i+j-1} \cdots s_{n-i}) \cdot (s_{n-i+k-1} \cdots s_{n-i+1}) \cdot s_{n-i}] \otimes \lambda^{-(j+k)} r \\
&= [x \cdot (s_{n-i+j-1} \cdots s_{n-i}) \cdot (s_{n-i+k-1} \cdots s_{n-i+1})] \otimes s_{n-i} \cdot (\lambda^{-(j+k)} r) \\
&= [x \cdot (s_{n-i+j-1} \cdots s_{n-i}) \cdot (s_{n-i+k-1} \cdots s_{n-i+1})] \otimes \lambda^{-(j+k-1)} r \\
&= d_{k-1}^{i-1} d_j^i(x \otimes r)
\end{aligned}$$

where the third equality holds because this computation takes place in $W(n)_{i-2} = \mathrm{TL}_n \otimes_{\mathrm{TL}_{n-i+1}} \mathbb{1}$ and $s_{n-i} \in \mathrm{TL}_{n-i+1}$. \square

Remark 4.3. Let us explain the motivation for the definition of $W(n)$. Let \mathfrak{S}_n denote the symmetric group on n letters. The *complex of injective words* is the chain complex $\mathcal{C}(n)$ of \mathfrak{S}_n -modules, concentrated in degrees -1 to $(n-1)$, that in degree i is the free R -module with basis given by tuples (x_0, \dots, x_i) where $x_0, \dots, x_i \in \{1, \dots, n\}$ and no letter appears more than once. We allow the empty word $()$, which lies in degree -1 . The differential of $\mathcal{C}(n)$ sends a word (x_0, \dots, x_i) to the alternating sum $\sum_{j=0}^i (-1)^j (x_0, \dots, \widehat{x}_j, \dots, x_i)$. A theorem of Farmer [Far79] shows that the homology of $\mathcal{C}(n)$ vanishes in degrees $i \leq (n-2)$, and the same result has been proved since then by many authors [Maa79, BW83, Ker05, RW13]. The complex of injective words has been used by several authors to prove homological stability for the symmetric groups [Maa79, Ker05, RW13].

For this paragraph only, let us abuse our established notation and denote by $s_1, \dots, s_{n-1} \in \mathfrak{S}_n$ the elements defined by $s_i = (i \ i+1)$, the transposition of i with $i+1$. Then these elements satisfy the braid relations, i.e. the second and third identities of Definition 2.21. The complex of injective words $\mathcal{C}(n)$ can be rewritten in terms of the group ring $R\mathfrak{S}_n$ and the elements s_i . Indeed, it is shown in [Hep20] that $\mathcal{C}(n)_i \cong R\mathfrak{S}_n \otimes_{R\mathfrak{S}_{n-i-1}} \mathbb{1}$, where $\mathbb{1}$ is the trivial module of $R\mathfrak{S}_{n-i-1}$, and that under this isomorphism the differential $d^i: \mathcal{C}(n)_i \rightarrow \mathcal{C}(n)_{i-1}$ becomes the map


$$d^i: R\mathfrak{S}_n \otimes_{R\mathfrak{S}_{n-i-1}} \mathbb{1} \longrightarrow R\mathfrak{S}_n \otimes_{R\mathfrak{S}_{n-i}} \mathbb{1}$$

defined by $d^i(x \otimes 1) = \sum_{j=0}^i (-1)^j x \cdot (s_{n-i+j-1} \cdots s_{n-i}) \otimes 1$. (There are no constants λ in this expression). Comparing this description of $\mathcal{C}(n)$ with our definition of $W(n)$, we see that our complex of planar injective words is precisely analogous to the original complex of injective words, after systematically replacing the group algebras of symmetric groups with the Temperley-Lieb algebras. The lack of constants in the differential for $\mathcal{C}(n)$ is explained by the fact that the effect of s_i on $\mathbb{1}$ is multiplication by λ in the Temperley-Lieb setting, and multiplication by 1 in the symmetric group setting.

Since we regard the Temperley-Lieb algebra as the planar analogue of the symmetric group, we chose the name *planar injective words* for our complex $W(n)$. This seemed the least discordant way of giving our complex an appropriate name. See the next remark for a means of picturing the complex.

Remark 4.4. Let us describe a method for visualising $W(n)$. Recall from the diagrammatic description of $\mathrm{TL}_n(a) \otimes_{\mathrm{TL}_m(a)} \mathbb{1}$ when $m \leq n$ given in Remark 2.11 that elements of $W(n)_i$ can be regarded as diagrams where the first $n-i-1$ dots on the right are encapsulated within a black box, and if any cups can be absorbed into the black box, then the diagram is identified with 0. The differential $d^i: W(n)_i \rightarrow W(n)_{i-1}$ is then given by pasting special elements onto the right of a diagram, followed by taking their signed and weighted sum. These special elements each

enlarge the black box by an extra strand, and plumb one of the free strands into the new space in the black box: Here is an example for $n = 4$ and $i = 2$.



The resulting diagrams can be simplified using the smoothing rules for diagrams with crossings described in Remark 2.22. We leave it to the reader to make this description as precise as they wish, and note here that this is where the notion of *braiding*, so often seen in homological stability arguments, fits into our set up.

Remark 4.5. Readers who are familiar with the theory will recognise that $W(n)$ is the chain complex associated to an augmented semi-simplicial $\text{TL}_n(a)$ -module.

The main result about the complex of planar injective words is the following, which we recall from the introduction. It is analogous to the homological vanishing property of the complex of injective words first proved by Farmer [Far79].

Theorem E. *The homology of $W(n)$ vanishes in degrees $d \leq (n - 2)$.*

The proof of Theorem E is the most technical part of this work, and will be given in Section 8.

The complex of injective words on n letters has rich combinatorial features: its Euler characteristic is the number of derangements of $\{1, \dots, n\}$; when one works over \mathbb{C} , its top homology has a description as a virtual representation that categorifies a well-known alternating sum formula for the number of derangements; and again when one works over \mathbb{C} , its top homology has a compact description in terms of Young diagrams and counts of standard Young tableaux. In the associated paper [BH21] we establish analogues of these for the complex of planar injective words. In particular we show that when the ring R is Noetherian the rank of $H_{n-1}(W(n))$ is the n -th *Fine number* [DS01]. (The rank of the Temperley-Lieb algebra is the n -th *Catalan number*, which is the number of Dyck paths of length $2n$. The n -th Fine number is the number of Dyck paths of length $2n$ whose first peak occurs at an even height, and as we explain in [BH21], it is an analogue of the number of derangements.) We also discover a new feature of the complex: the differentials have an alternate expression in terms not of the s_i but of the U_i . This expression demonstrates a connection with the *Jacobsthal numbers*, and we will briefly explain the result for the top differential below. The top homology of the Tits building is known as the *Steinberg module*. This inspires the name in the following definition.

Definition 4.6. We define the n -th *Fineberg module* to be the $\text{TL}_n(a)$ -module $\mathcal{F}_n(a) = H_{n-1}(W(n))$. We often suppress the a and simply write \mathcal{F}_n .

The Fineberg module is an important ingredient in the full statement of our stability result, Theorem 5.1. In order to detect the non-zero homology group

appearing in Theorem C we need to study it in more detail using the connection with Jacobsthal numbers from [BH21].

The n -th *Jacobsthal number* J_n [OEIS] is (among other things) the number of sequences $n > a_1 > a_2 > \cdots > a_r > 0$ whose initial term has the opposite parity to n . Some examples, when $n = 4$, are $3, 1, 3 > 2, 3 > 1$ and $3 > 2 > 1$. (We allow the empty sequence, and say that by convention its initial term is $a_1 = 0$ and $r = 0$. Of course this only occurs when n is odd.) Another viewpoint of J_n in terms of compositions of n is given in [BH21].

Definition 4.7. Let $a = v + v^{-1}$ where $v \in R^\times$ is a unit. We define the *Jacobsthal element* in $\mathrm{TL}_n(a)$ as follows:

$$\mathcal{J}_n = (-1)^{n-1} \sum_{\substack{n > a_1 > \cdots > a_r > 0 \\ n - a_1 \text{ odd}}} \left(\frac{\mu}{\lambda}\right)^r U_{a_1} \cdots U_{a_r}$$

Recall we allow the empty sequence ($a_1 = 0$ and $r = 0$) when n is odd. This corresponds to a constant summand 1 in \mathcal{J}_n for odd n . Note that the number of irreducible terms in \mathcal{J}_n is J_n .

Example 4.8. In the cases $n = 1, 2, 3, 4$, and choosing $\theta = \theta_1$ so that $(\lambda, \mu) = (-1, v)$, we have:

$$\begin{aligned} \mathcal{J}_1 &= 1 \\ \mathcal{J}_2 &= vU_1 \\ \mathcal{J}_3 &= v^2U_2U_1 - vU_2 + 1 \\ \mathcal{J}_4 &= v^3U_3U_2U_1 - v^2U_3U_2 - v^2U_3U_1 + vU_3 + vU_1 \end{aligned}$$

Spencer has computed the Jacobsthal elements \mathcal{J}_n up to $n = 9$, these can be viewed at [Spe22].

Since \mathcal{F}_n is the homology of $W(n)$ in the top degree, it is simply the kernel of the top differential $d^{n-1}: W(n)_{n-1} \rightarrow W(n)_{n-2}$. There are identifications

$$W(n)_{n-1} = \mathrm{TL}_n(a) \otimes_{\mathrm{TL}_0(a)} \mathbb{1} \cong \mathrm{TL}_n(a) \text{ and } W(n)_{n-2} \cong \mathrm{TL}_n(a) \otimes_{\mathrm{TL}_1(a)} \mathbb{1} \cong \mathrm{TL}_n(a).$$

The following is shown in Theorem D of [BH21]:

Proposition 4.9. *Under the above identifications, the top differential of $W(n)$ is right-multiplication by \mathcal{J}_n . In particular, there is an exact sequence*


$$0 \longrightarrow \mathcal{F}_n(a) \longrightarrow \mathrm{TL}_n(a) \xrightarrow{-\mathcal{J}_n} \mathrm{TL}_n(a).$$

Remark 4.10. Note that Definition 4.7 gives a different value for the element \mathcal{J}_n than the one that appears in Definition 8.1 and Theorem D of [BH21]. This is because the proof of Theorem D of [BH21] contains a sign error: it assumes that $s_i = (\lambda - \mu U_i)$ rather than $s_i = (\lambda + \mu U_i)$ as it should have done. This error is fixed by replacing μ with $-\mu$ in the formula in Definition 8.1 of [BH21]. It is possible

to check Example 4.8 by hand to confirm that the signs in the present formula for \mathcal{J}_n are the correct ones.

The Fineberg module \mathcal{F}_n appears to be a new and interesting representation, and looks likely to be highly nontrivial for each choice of n . Let us illustrate this by computing \mathcal{F}_2 , \mathcal{F}_3 and \mathcal{F}_4 . We will continue with the choice $\theta = \theta_1$ so that $(\lambda, \mu) = (-1, v)$.

Our description will be phrased in terms of the cell modules of TL_n , which we describe briefly. A *half-diagram* (or *link state* in the language of [RSA14]) consists of a vertical line in the plane decorated with dots labelled $1, \dots, n$ from bottom to top, together with a collection of arcs in the plane, each of which either connects two dots, or is connected to a dot at one end, in such a way that each dot lies on precisely one arc. The arcs must lie to the right of the vertical line, they must be disjoint, and the half-diagrams are taken up to isotopy. Thus the half-diagrams on 4 dots are as follows.



The *cell module* $S(n, m)$ is the TL_n -module with R -basis consisting of the half-diagrams on n dots in which m arcs have free ends. The TL_n -module structure on $S(n, m)$ is obtained by pasting planar diagrams onto the left of half-diagrams and simplifying the result exactly as with composition in TL_n , with the extra condition that if pasting produces an arc with two free ends, then the resulting diagram is set to 0. In $S(4, 2)$, for example, we have:

$$U_1 \cdot \begin{array}{c} \text{---} \\ | \\ \bullet \\ | \\ \bullet \\ | \\ \bullet \\ | \\ \bullet \\ \text{---} \\ \text{---} \end{array} = a \cdot \begin{array}{c} \text{---} \\ | \\ \bullet \\ | \\ \bullet \\ | \\ \bullet \\ | \\ \bullet \\ \text{---} \\ \text{---} \end{array} \quad U_2 \cdot \begin{array}{c} \text{---} \\ | \\ \bullet \\ | \\ \bullet \\ | \\ \bullet \\ | \\ \bullet \\ \text{---} \\ \text{---} \end{array} = \begin{array}{c} \text{---} \\ | \\ \bullet \\ | \\ \bullet \\ | \\ \bullet \\ | \\ \bullet \\ \text{---} \\ \text{---} \end{array} \quad U_3 \cdot \begin{array}{c} \text{---} \\ | \\ \bullet \\ | \\ \bullet \\ | \\ \bullet \\ | \\ \bullet \\ \text{---} \\ \text{---} \end{array} = 0$$

(The reader is reminded that we label the dots from bottom to top.) Observe that $S(n, n) = \mathbb{1}$ is the trivial module for each n , and that $S(n, m)$ is nonzero only when $n - m$ is even.

Example 4.11 (The Fineberg module \mathcal{F}_2). The module \mathcal{F}_2 is the kernel of the map $\text{TL}_2 \rightarrow \text{TL}_2$, $x \mapsto x \cdot \mathcal{J}_2$. Now $\mathcal{J}_2 = vU_1$ as in Example 4.8, so that \mathcal{F}_2 is the R -module of rank 1 spanned by the element $a - U_1$. This is a copy of the trivial module $\mathbb{1} = S(2, 2)$.

Example 4.12 (The Fineberg module \mathcal{F}_3). The module \mathcal{F}_3 is the kernel of the map $\text{TL}_3 \rightarrow \text{TL}_3$, $x \mapsto x \cdot \mathcal{J}_3$, where $\mathcal{J}_3 = v^2U_2U_1 - vU_2 + 1$ as in Example 4.8. Thus \mathcal{F}_3 is the R -module of rank 2 with basis elements

$$\begin{aligned}
 \alpha &= U_1U_2 - vU_1, \\
 \beta &= U_2 - vU_2U_1.
 \end{aligned}$$

One can now check that there is an isomorphism of TL_3 -modules $\mathcal{F}_3 \cong S(3, 1)$ given by:

$$\mathcal{F}_3 \xrightarrow{\cong} S(3, 1) \quad \alpha \mapsto \begin{array}{c} \bullet \\ | \\ \bullet \\ | \\ \bullet \end{array} \begin{array}{l} \text{---} \\ \curvearrowright \\ \text{---} \end{array} \quad \beta \mapsto \begin{array}{c} \bullet \\ | \\ \bullet \\ | \\ \bullet \end{array} \begin{array}{l} \curvearrowleft \\ \text{---} \\ \text{---} \end{array}$$

Example 4.13 (The Fineberg module \mathcal{F}_4). The module \mathcal{F}_4 is the kernel of the map $\mathrm{TL}_4 \rightarrow \mathrm{TL}_4$, $x \mapsto x \cdot \mathcal{J}_4$, where $\mathcal{J}_4 = v^3 U_3 U_2 U_1 - v^2 U_3 U_2 - v^2 U_3 U_1 + v U_3 + v U_1$ as in Example 4.8. It is now possible to check (at length) that \mathcal{F}_4 is a free R -module of rank 6 with the following basis:

$$\begin{aligned} A &= U_3 U_1 - a U_3 U_1 U_2 \\ B &= U_2 U_3 U_1 - a U_2 U_3 U_1 U_2 \\ X &= U_1 U_2 U_3 - U_3 U_1 U_2 - a U_1 U_2 + U_1 \\ Y &= U_2 U_3 - U_2 U_3 U_1 U_2 - a U_2 + U_2 U_1 \\ Z &= U_3 U_2 U_1 - U_3 U_1 U_2 - a U_3 U_2 + U_3 \\ P &= U_3 U_1 U_2 - U_1 - U_3 + a. \end{aligned}$$

If we now define

$$M_0 = \mathrm{span}(A, B), \quad M_1 = \mathrm{span}(A, B, X, Y, Z), \quad M_2 = \mathrm{span}(A, B, X, Y, Z, P)$$

so that $M_0 \subseteq M_1 \subseteq M_2 = \mathcal{F}_4$, then one can check directly (by computing the effect of multiplying on the left by U_1, U_2, U_3) that M_0 and M_1 are submodules of \mathcal{F}_4 , and moreover, that we have isomorphisms:

$$\begin{aligned} M_0 &\xrightarrow{\cong} S(4, 0), & A &\mapsto \begin{array}{c} \bullet \\ | \\ \bullet \\ | \\ \bullet \\ | \\ \bullet \end{array} \begin{array}{l} \curvearrowright \\ \curvearrowright \\ \text{---} \end{array} & B &\mapsto \begin{array}{c} \bullet \\ | \\ \bullet \\ | \\ \bullet \\ | \\ \bullet \end{array} \begin{array}{l} \curvearrowleft \\ \curvearrowleft \\ \text{---} \end{array} \\ M_1/M_0 &\xrightarrow{\cong} S(4, 2), & X &\mapsto \begin{array}{c} \bullet \\ | \\ \bullet \\ | \\ \bullet \\ | \\ \bullet \end{array} \begin{array}{l} \text{---} \\ \text{---} \\ \curvearrowright \\ \text{---} \end{array} & Y &\mapsto \begin{array}{c} \bullet \\ | \\ \bullet \\ | \\ \bullet \\ | \\ \bullet \end{array} \begin{array}{l} \text{---} \\ \text{---} \\ \curvearrowleft \\ \text{---} \end{array} & Z &\mapsto \begin{array}{c} \bullet \\ | \\ \bullet \\ | \\ \bullet \\ | \\ \bullet \end{array} \begin{array}{l} \text{---} \\ \text{---} \\ \text{---} \\ \curvearrowright \end{array} \\ M_2/M_1 &\xrightarrow{\cong} \mathbb{1}, & P &\mapsto 1 \end{aligned}$$

Thus \mathcal{F}_4 has a filtration in which each of the three cell modules appears as precisely one of the filtration quotients. We emphasise that this result holds with *no* further assumptions on the ground ring R or on the parameter v .

5. HOMOLOGICAL STABILITY AND STABLE HOMOLOGY

The aim of this section is to prove the following result. Theorem B is an immediate consequence, and Theorem C will be proved in the next section as a corollary of it.

Theorem 5.1. *Let R be a commutative ring, let $v \in R$ be a unit, and let $a = v + v^{-1}$. Then for n odd we have:*

$$\mathrm{Tor}_i^{\mathrm{TL}_n(a)}(\mathbb{1}, \mathbb{1}) \cong \begin{cases} R & i = 0 \\ 0 & 1 \leq i \leq (n-1) \\ \mathrm{Tor}_{i-n}^{\mathrm{TL}_n(a)}(\mathbb{1}, \mathcal{F}_n(a)) & i \geq n \end{cases}$$

and for n even we have

$$\mathrm{Tor}_i^{\mathrm{TL}_n(a)}(\mathbb{1}, \mathbb{1}) \cong \begin{cases} R & i = 0 \\ 0 & 1 \leq i \leq (n-2) \\ \mathrm{Tor}_{i-n}^{\mathrm{TL}_n(a)}(\mathbb{1}, \mathcal{F}_n(a)) & i \geq (n+1) \end{cases}$$

for $i \neq n-1, n$, while in degrees $(n-1)$ and n there is an exact sequence

$$0 \rightarrow \mathrm{Tor}_n^{\mathrm{TL}_n(a)}(\mathbb{1}, \mathbb{1}) \rightarrow \mathbb{1} \otimes_{\mathrm{TL}_n(a)} \mathcal{F}_n(a) \xrightarrow{\mathcal{Q}_n} \mathbb{1} \rightarrow \mathrm{Tor}_{n-1}^{\mathrm{TL}_n(a)}(\mathbb{1}, \mathbb{1}) \rightarrow 0. \quad (1)$$

Analogous results hold for the Ext-groups. For n odd we have:

$$\mathrm{Ext}_{\mathrm{TL}_n(a)}^i(\mathbb{1}, \mathbb{1}) \cong \begin{cases} R & i = 0 \\ 0 & 1 \leq i \leq (n-1) \\ \mathrm{Ext}_{\mathrm{TL}_n(a)}^{i-n}(\mathcal{F}_n(a), \mathbb{1}) & i \geq n \end{cases}$$

and for n even we have

$$\mathrm{Ext}_{\mathrm{TL}_n(a)}^i(\mathbb{1}, \mathbb{1}) \cong \begin{cases} R & i = 0 \\ 0 & 1 \leq i \leq (n-2) \\ \mathrm{Ext}_{\mathrm{TL}_n(a)}^{i-n}(\mathcal{F}_n(a), \mathbb{1}) & i \geq (n+1) \end{cases}$$

for $i \neq n-1, n$, while in degrees $(n-1)$ and n there is an exact sequence

$$0 \rightarrow \mathrm{Ext}_{\mathrm{TL}_n(a)}^{n-1}(\mathbb{1}, \mathbb{1}) \rightarrow \mathbb{1} \xrightarrow{\mathcal{Q}^n} \mathrm{Hom}_{\mathrm{TL}_n(a)}(\mathcal{F}_n(a), \mathbb{1}) \rightarrow \mathrm{Ext}_{\mathrm{TL}_n(a)}^n(\mathbb{1}, \mathbb{1}) \rightarrow 0. \quad (2)$$

The central maps \mathcal{Q}_n and \mathcal{Q}^n of (1) and (2) respectively are described as follows. Regard $\mathcal{F}_n(a)$ as a left-submodule of $\mathrm{TL}_n(a)$ as in Proposition 4.9. Then the maps are

$$\mathcal{Q}_n: \mathbb{1} \otimes_{\mathrm{TL}_n(a)} \mathcal{F}_n(a) \longrightarrow \mathbb{1}, \quad x \otimes f \longmapsto x \cdot f$$

and

$$\mathcal{Q}^n: \mathbb{1} \longrightarrow \mathrm{Hom}_{\mathrm{TL}_n(a)}(\mathcal{F}_n(a), \mathbb{1}), \quad x \longmapsto (f \mapsto f \cdot x)$$

where $x \cdot f$ and $f \cdot x$ denote the action of $f \in \mathcal{F}_n(a) \subseteq \mathrm{TL}_n(a)$ on the right and left of $\mathbb{1}$, respectively.

In order to prove this theorem, we will use the complex of planar injective words $W(n)$ introduced in the previous section. Recall that the Fineberg module \mathcal{F}_n appearing in the statement is the top homology group $H_{n-1}(W(n))$.

Lemma 5.2. *The homology groups of both the complex $\mathbb{1} \otimes_{\mathrm{TL}_n(a)} W(n)$ and the complex $\mathrm{Hom}_{\mathrm{TL}_n(a)}(W(n), \mathbb{1})$ are concentrated in degree $(n-1)$, where in both cases they are given by $\mathbb{1}$ if n is even and 0 if n is odd.*

Proof. We have $W(n)_i = \mathrm{TL}_n \otimes_{\mathrm{TL}_{n-i-1}} \mathbb{1}$, and the boundary map $d^i: W(n)_i \rightarrow W(n)_{i-1}$ is given by $x \otimes r \mapsto x \cdot D_i \otimes r$, where $D_i = \sum_{j=0}^i (-1)^j s_{n-i+j-1} \cdots s_{n-i} \lambda^{-j}$.

By regarding $\mathbb{1}$ as both a left and right TL_n -module, we may regard $\mathbb{1} \otimes_{\mathrm{TL}_n} W(n)_i$ as a left TL_n -module. With this TL_n -module structure, we obtain $\mathbb{1} \otimes_{\mathrm{TL}_n} W(n)_i = \mathbb{1} \otimes_{\mathrm{TL}_n} (\mathrm{TL}_n \otimes_{\mathrm{TL}_{n-i-1}} \mathbb{1}) \cong \mathbb{1}$. Under these isomorphisms, the boundary map originating in degree i becomes the action on $\mathbb{1}$ of the element D_i . Similarly, $\mathrm{Hom}_{\mathrm{TL}_n}(W(n)_i, \mathbb{1}) = \mathrm{Hom}_{\mathrm{TL}_n}(\mathrm{TL}_n \otimes_{\mathrm{TL}_{n-i-1}} \mathbb{1}, \mathbb{1}) \cong \mathbb{1}$, and under these isomorphisms the boundary map originating in degree $(i-1)$ becomes the action of the element D_i on $\mathbb{1}$.

The action of $s_{n-i+j-1} \cdots s_{n-i}$ on $\mathbb{1}$ is simply multiplication by λ^j , with one factor of λ for each s term (recall $s_i = \mu U_i + \lambda$). Thus the action of D_i on $\mathbb{1}$ is nothing other than multiplication by $\sum_{j=0}^i (-1)^j$, which is 0 for i odd and 1 for i even.

So altogether $\mathbb{1} \otimes_{\mathrm{TL}_n} W(n)$ and $\mathrm{Hom}_{\mathrm{TL}_n}(W(n), \mathbb{1})$ are isomorphic to complexes with a copy of R in each degree $i = -1, \dots, (n-1)$ and with boundary maps alternating between the identity map and 0. In $\mathbb{1} \otimes_{\mathrm{TL}_n} W(n)$ the identity maps originate in even degrees, and in $\mathrm{Hom}_{\mathrm{TL}_n}(W(n), \mathbb{1})$ they originate in odd degrees. The claim now follows. \square

Proof of Theorem 5.1. We begin with the Tor-case.

In degree $d = 0$ the theorem holds trivially. Recall that P_* is a projective resolution of $\mathbb{1}$ as a right TL_n -module. We use the two homological spectral sequences $\{^I E^r\}$ and $\{^{II} E^r\}$ associated to $W(n)$ as described in Section 3.2.

Let us consider $\{^I E^r\}$. We have

$$^I E_{i,j}^2 = \begin{cases} \mathrm{Tor}_i^{\mathrm{TL}_n}(\mathbb{1}, \mathcal{F}_n) & j = (n-1) \\ 0 & j \neq (n-1) \end{cases}$$

and consequently the spectral sequence converges to $\mathrm{Tor}_{*-n+1}^{\mathrm{TL}_n}(\mathbb{1}, \mathcal{F}_n)$, for $* = i + j$. The same is therefore true of $\{^{II} E^r\}$.

Let us write $\varepsilon_n = H_{n-1}(\mathbb{1} \otimes_{\mathrm{TL}_n} W(n))$, so that by Lemma 5.2, ε_n is trivial for n odd and $\mathbb{1}$ for n even. Since \mathcal{F}_n consists of the cycles in $W(n)_{n-1}$, the map

$$\mathbb{1} \otimes_{\mathrm{TL}_n} \mathcal{F}_n \rightarrow \mathbb{1} \otimes_{\mathrm{TL}_n} W(n)_{n-1}$$

again lands in the cycles, giving us a map

$$\mathbb{1} \otimes_{\mathrm{TL}_n} \mathcal{F}_n \rightarrow H_{n-1}(\mathbb{1} \otimes_{\mathrm{TL}_n} W(n)) = \varepsilon_n.$$

When n is even and ε_n is identified with $\mathbb{1}$ as in the lemma, then this map simply becomes \mathcal{Q}_n as described in the statement of the theorem.




FIGURE 8. The page ${}^IIE^1$. The only differentials that affect the ${}^IIE^2$ page are shown on the $j = 0$ row.

We now know that $\{{}^IIE^r\}$ converges to $\text{Tor}_{*,-n+1}^{\text{TL}_n}(\mathbb{1}, \mathcal{F}_n)$. Its E^1 -page ${}^IIE_{i,j}^1 = \text{Tor}_j^{\text{TL}_n}(\mathbb{1}, W(n)_i)$ is largely known to us. Indeed, when $j = 0$ the terms are $\text{Tor}_0^{\text{TL}_n}(\mathbb{1}, W(n)_i) = \mathbb{1} \otimes_{\text{TL}_n} W(n)_i$, with d^1 -maps between them induced by the boundary maps of $W(n)$. In other words, the $j = 0$ part of ${}^IIE_{i,j}^1$ is precisely the complex $\mathbb{1} \otimes_{\text{TL}_n} W(n)$. When $0 \leq i \leq (n-1)$, the term $W(n)_i = \text{TL}_n \otimes_{\text{TL}_{n-i-1}} \mathbb{1}$ satisfies $0 \leq (n-i-1) < n$, so that by Theorem F we have

$${}^IIE_{i,j}^1 = \text{Tor}_j^{\text{TL}_n}(\mathbb{1}, \text{TL}_n \otimes_{\text{TL}_{n-i-1}} \mathbb{1}) = 0$$

for $j > 0$. When $i = -1$ we have $W(n)_{-1} = \mathbb{1}$ so that ${}^IIE_{-1,j}^1 = \text{Tor}_j^{\text{TL}_n}(\mathbb{1}, \mathbb{1})$ for $j > 0$. This is depicted in Figure 8.

By the description in the previous paragraph, we can now identify ${}^IIE_{*,*}^2$. The only possible differentials are in the $j = 0$ part, which is $\mathbb{1} \otimes_{\text{TL}_n} W(n)$, and whose homology is ε_n concentrated in degree $(n-1)$. Thus ${}^IIE_{*,*}^2$ is zero except for the following groups:

$${}^IIE_{i,j}^2 = \begin{cases} \text{Tor}_j^{\text{TL}_n}(\mathbb{1}, \mathbb{1}) & i = -1, j > 0 \\ \varepsilon_n & i = (n-1), j = 0 \end{cases}$$

as depicted in Figure 9.

From the E^2 -page onwards there is precisely one possible differential, namely $d^n: E_{n-1,0}^n \rightarrow E_{-1,n-1}^n$, which is a map $d^n: \varepsilon_n \rightarrow \text{Tor}_{n-1}^{\text{TL}_n}(\mathbb{1}, \mathbb{1})$. It forms part of an




FIGURE 9. The page ${}^{II}E^2$. This page stays constant until ${}^{II}E^n$ where the only possible further differential lies: this is shown in red. The $i + j = n - 1$ and $i + j = n - 2$ diagonals are indicated in blue.

exact sequence

$$0 \rightarrow {}^{II}E_{n-1,0}^\infty \rightarrow \varepsilon_n \xrightarrow{d^n} \text{Tor}_{n-1}^{\text{TL}_n}(\mathbb{1}, \mathbb{1}) \rightarrow {}^{II}E_{-1,n-1}^\infty \rightarrow 0$$

In ${}^{II}E_{*,*}^\infty$, each total degree has only one non-zero group, except (possibly) for total degree $(n - 1)$, where we have the two groups ${}^{II}E_{-1,n}^\infty$ and ${}^{II}E_{n-1,0}^\infty$. The relationship between the infinity-page of a spectral sequence and the sequence's target now give us a short exact sequence:

$$0 \rightarrow {}^{II}E_{-1,n}^\infty \rightarrow \text{Tor}_0^{\text{TL}_n}(\mathbb{1}, \mathcal{F}_n) \rightarrow {}^{II}E_{n-1,0}^\infty \rightarrow 0$$

The last two exact sequences combine to give us:

$$0 \rightarrow {}^{II}E_{-1,n}^\infty \rightarrow \text{Tor}_0^{\text{TL}_n}(\mathbb{1}, \mathcal{F}_n) \rightarrow \varepsilon_n \rightarrow \text{Tor}_{n-1}^{\text{TL}_n}(\mathbb{1}, \mathbb{1}) \rightarrow {}^{II}E_{-1,n-1}^\infty \rightarrow 0$$

The leftmost term is ${}^{II}E_{-1,n}^\infty = {}^{II}E_{-1,n}^2 = \text{Tor}_n^{\text{TL}_n}(\mathbb{1}, \mathbb{1})$. And ${}^{II}E_{-1,n-1}^\infty$ is the only group in total degree $(n - 2)$, and therefore coincides with $\text{Tor}_{(n-2)-n+1}^{\text{TL}_n}(\mathbb{1}, \mathcal{F}_n) = \text{Tor}_{-1}^{\text{TL}_n}(\mathbb{1}, \mathcal{F}_n) = 0$. And $\text{Tor}_0^{\text{TL}_n}(\mathbb{1}, \mathcal{F}_n) = \mathbb{1} \otimes_{\text{TL}_n} \mathcal{F}_n$. So the last exact sequence becomes:

$$0 \rightarrow \text{Tor}_n^{\text{TL}_n}(\mathbb{1}, \mathbb{1}) \rightarrow \mathbb{1} \otimes_{\text{TL}_n} \mathcal{F}_n \rightarrow \varepsilon_n \rightarrow \text{Tor}_{n-1}^{\text{TL}_n}(\mathbb{1}, \mathbb{1}) \rightarrow 0$$

When n is even, we claim that the map $\mathbb{1} \otimes_{\text{TL}_n} \mathcal{F}_n \rightarrow \varepsilon_n$ in this sequence is \mathcal{Q}_n . Let $\mathcal{F}_n[n-1]$ be the complex consisting of a copy of \mathcal{F}_n concentrated in degree $n-1$.

There is a natural inclusion of chain complexes $\mathcal{F}_n[n-1] \hookrightarrow W(n)$, and this leads to a map of double complexes and then of spectral sequences. The map $\mathbb{1} \otimes_{\text{TL}_n} \mathcal{F}_n \rightarrow \varepsilon_n$ can be identified using this map of spectral sequences.

It follows from the sequence that in the case n odd, when $\varepsilon_n = 0$, the final term satisfies $\text{Tor}_{n-1}^{\text{TL}_n}(\mathbb{1}, \mathbb{1}) = 0$, and the first two terms satisfy

$$\text{Tor}_n^{\text{TL}_n}(\mathbb{1}, \mathbb{1}) \cong \mathbb{1} \otimes_{\text{TL}_n} \mathcal{F}_n = \text{Tor}_0^{\text{TL}_n}(\mathbb{1}, \mathcal{F}_n)$$

as required.

The previous discussion determines what happens in total degrees $(n-1)$ and $(n-2)$. In total degrees d other than $(n-1)$ and $(n-2)$, and when $j > 0$, the only term on the E^∞ page is ${}^{II}E_{-1, d+1}^\infty = \text{Tor}_{d+1}^{\text{TL}_n}(\mathbb{1}, \mathbb{1})$, which must therefore equal $\text{Tor}_{d-n+1}^{\text{TL}_n}(\mathbb{1}, \mathcal{F}_n)$. Thus $\text{Tor}_d^{\text{TL}_n}(\mathbb{1}, \mathbb{1}) \cong \text{Tor}_{d-n}^{\text{TL}_n}(\mathbb{1}, \mathcal{F}_n)$ for $d \neq n, n-1$. This completes the proof.

For the Ext-case we use the two cohomological spectral sequences associated to $W(n)$ as in Section 3.2, and then proceed dually to the above. We leave the details to the reader. \square

6. SHARPNESS

We recall the statement of Theorem C from the introduction.

Theorem C. *Let n be even and suppose that a is not a unit. Then $\text{Tor}_{n-1}^{\text{TL}_n(a)}(\mathbb{1}, \mathbb{1})$ is non-zero.*

Let $\mathcal{I} \subseteq \text{TL}_n$ denote the left-ideal generated by all diagrams which have a cup on the right in positions other than 1, together with all multiples of a . Thus

$$\mathcal{I} = (\text{TL}_n \cdot a) + (\text{TL}_n \cdot U_2) + \cdots + (\text{TL}_n \cdot U_{n-1}).$$

Lemma 6.1. *Let n be even or odd, and let $1 \leq p \leq n-1$. Then $U_p \cdot \mathcal{J}_n \in \mathcal{I}$.*

Proof. Recall from Definition 4.7 that the monomials appearing in \mathcal{J}_n are those of the form $U_{i_1} \cdots U_{i_r}$ where $(n-1) \geq i_1 > i_2 > \cdots > i_r \geq 1$ and $i_1 \equiv (n-1) \pmod{2}$, and that such a monomial appears in \mathcal{J}_n with coefficient $(-1)^{n-1} \binom{\mu}{\lambda}^r$. We write $\mathcal{J}_n = K_n + L_n$ where K_n is the part of \mathcal{J}_n featuring monomials of the form $U_i U_{i-1} \cdots U_1$ for $i \equiv n-1 \pmod{2}$ in the range $1 \leq i \leq n-1$, and L_n is the part of \mathcal{J}_n featuring the remaining monomials.

If $U_{i_1} \cdots U_{i_r}$ is a monomial appearing in L_n , then it must either end in U_{i_r} for $n-1 \geq i_r > 1$ or end in a monomial of the form $U_{i_j} \cdot U_{i_{j-1}} \cdots U_1 = (U_{i_{j-1}} \cdots U_1) \cdot U_{i_j}$ for some $i_j \geq i_{j-1} + 2$, $i_{j-1} \geq 1$ and hence must lie in \mathcal{I} . Thus $L_n \in \mathcal{I}$, and to prove the lemma it will be sufficient to show that $U_p \cdot K_n \in \mathcal{I}$.

Now observe that

$$K_n = (-1)^{n-1} \sum_{\substack{0 \leq i \leq (n-1) \\ i \equiv n-1 \pmod{2}}} \binom{\mu}{\lambda}^i \cdot U_i U_{i-1} \cdots U_1.$$

(In the case $i = 0$ the product $U_i \cdots U_1$ is empty and therefore equal to 1. This term only appears in K_n when n is odd.) Suppose that $U_i \cdots U_1$ is a monomial appearing in the above sum. Then:

$$U_p \cdot (U_i \cdots U_1) = \begin{cases} (U_p \cdots U_1) \cdot (U_i \cdots U_{p+2}) & p \leq i - 2 \\ U_{i-1} \cdots U_1 & p = i - 1 \\ U_i \cdots U_1 \cdot a & p = i \\ U_{i+1} \cdots U_1 & p = i + 1 \\ (U_i \cdots U_1) \cdot U_p & p \geq i + 2 \end{cases}$$

Thus $U_p \cdot (U_i \cdots U_1) \in \mathcal{I}$ except for the cases $i = p - 1$, $i = p + 1$. When $p \equiv (n - 1) \pmod{2}$ these exceptional cases never occur, since we have assumed $i \equiv (n - 1) \pmod{2}$, and so $U_p \cdot K_n \in \mathcal{I}$ as required. And when $p \equiv n \pmod{2}$, we can compute the contribution from the two exceptional cases to find that, modulo \mathcal{I} , $U_p \cdot \mathcal{J}_n$ is equal to

$$\begin{aligned} & (-1)^{n-1} \left(\frac{\mu}{\lambda}\right)^{p-1} U_p \cdot (U_{p-1} \cdots U_1) + (-1)^{n-1} \left(\frac{\mu}{\lambda}\right)^{p+1} U_p \cdot (U_{p+1} \cdots U_1) \\ &= (-1)^{n-1} \left(\frac{\mu}{\lambda}\right)^{p-1} \cdot (U_p \cdots U_1) + (-1)^{n-1} \left(\frac{\mu}{\lambda}\right)^{p+1} \cdot (U_p \cdots U_1) \\ &= (-1)^{n-1} \left(\frac{\mu}{\lambda}\right)^p \left[\left(\frac{\mu}{\lambda}\right)^{-1} + \left(\frac{\mu}{\lambda}\right)^1 \right] \cdot (U_p \cdots U_1) \in \mathcal{I}. \end{aligned}$$

Now from Definition 2.20 we have either $(\mu, \lambda) = (v, -1)$ or $(\mu, \lambda) = (-v, v^2)$. In both cases the square bracket above evaluates to $-a$ (recall $a = v + v^{-1}$). Thus $U_p \cdot K_n$ is a multiple of a and therefore in \mathcal{I} as required. \square

Lemma 6.2. *Let n be even. Let $x \in \mathcal{F}_n(a)$, so that $x \cdot \mathcal{J}_n = 0$. Then the constant term of x is a multiple of a .*

Proof. Let b be the constant term of x , so that x is equal to b plus a linear combination of left-multiples of the elements U_1, \dots, U_{n-1} . Thus $x \cdot \mathcal{J}_n$ is equal to $b \cdot \mathcal{J}_n$ plus a linear combination of left-multiples of $U_1 \cdot \mathcal{J}_n, \dots, U_{n-1} \cdot \mathcal{J}_n$, all of which lie in \mathcal{I} by Lemma 6.1. Thus $x \cdot \mathcal{J}_n = b \cdot \mathcal{J}_n$ modulo \mathcal{I} .

As an R -module, the quotient TL_n/\mathcal{I} is isomorphic to the direct sum of copies of R/aR , with one summand for each monomial whose Jones normal form ends with U_1 . We have that

$$\mathcal{J}_n = (-1)^{n-1} \left[\left(\frac{\mu}{\lambda}\right) U_1 + \left(\frac{\mu}{\lambda}\right)^3 U_3 U_2 U_1 + \cdots \right] \text{ in } \text{TL}_n/\mathcal{I}$$

and it follows that

$$b \cdot \mathcal{J}_n = (-1)^{n-1} \left[b \left(\frac{\mu}{\lambda}\right) U_1 + b \left(\frac{\mu}{\lambda}\right)^3 U_3 U_2 U_1 + \cdots \right] \text{ in } \text{TL}_n/\mathcal{I},$$

so b must vanish in R/aR . \square

Lemma 6.3. *Let n be even. Then the image of the map*

$$\mathbb{1} \otimes_{\mathrm{TL}_n(a)} \mathcal{F}_n(a) \rightarrow \mathbb{1}, \quad 1 \otimes x \mapsto 1 \cdot x,$$

is contained in the ideal generated by a .

Proof. Since the elements U_p act on $\mathbb{1}$ as multiplication by 0, the map above simply sends $1 \otimes x$ to the constant term of x . But the previous lemma tells us that the constant term of x is a multiple of a . \square

Proof of Theorem C. Let n be even. From Theorem 5.1, we have the (fairly short) exact sequence

$$0 \rightarrow \mathrm{Tor}_n^{\mathrm{TL}_n}(\mathbb{1}, \mathbb{1}) \rightarrow \mathbb{1} \otimes_{\mathrm{TL}_n} \mathcal{F}_n \rightarrow \mathbb{1} \rightarrow \mathrm{Tor}_{n-1}^{\mathrm{TL}_n}(\mathbb{1}, \mathbb{1}) \rightarrow 0$$

and the image of $\mathbb{1} \otimes_{\mathrm{TL}_n} \mathcal{F}_n \rightarrow \mathbb{1}$ is contained in the ideal generated by a , and in particular does not contain the element 1, so that $\mathrm{Tor}_{n-1}^{\mathrm{TL}_n}(\mathbb{1}, \mathbb{1}) \neq 0$. \square

7. THE CASE OF $\mathrm{TL}_2(a)$

In this section we briefly consider the case $n = 2$, and fully compute the Tor and Ext groups. We do this first by a straightforward computation using an explicit free resolution. Then, in order to illustrate the theory developed in the paper, we re-prove the same result by explicitly computing the Fineberg module \mathcal{F}_2 and applying Theorem 5.1.

Proposition 7.1. *The homology and cohomology of $\mathrm{TL}_2(a)$ are as follows.*

$$\mathrm{Tor}_i^{\mathrm{TL}_2(a)}(\mathbb{1}, \mathbb{1}) = \begin{cases} R, & i = 0, \\ R/aR, & i > 0, i \text{ odd}, \\ R_a, & i > 0, i \text{ even}, \end{cases}$$

$$\mathrm{Ext}_{\mathrm{TL}_2(a)}^i(\mathbb{1}, \mathbb{1}) = \begin{cases} R, & i = 0, \\ R_a, & i > 0, i \text{ odd}, \\ R/aR, & i > 0, i \text{ even}, \end{cases}$$

where R_a denotes the kernel of the map $R \xrightarrow{a} R$. This holds for any choice of ground ring R and any choice of parameter $a \in R$.

Proof. We define a chain complex of left TL_2 -modules as follows. The degree is indicated in the right-hand column. The boundary maps are given by right-multiplication by the indicated element of TL_2 , except for the last, which is the

map $\mathrm{TL}_2 \rightarrow \mathbb{1}$, $x \mapsto x \cdot 1$.

$$\begin{array}{ccc}
 \vdots & & \\
 \downarrow (a-U_1) & & \\
 \mathrm{TL}_2 & 3 & \\
 \downarrow U_1 & & \\
 \mathrm{TL}_2 & 2 & \\
 \downarrow (a-U_1) & & \\
 \mathrm{TL}_2 & 1 & \\
 \downarrow U_1 & & \\
 \mathrm{TL}_2 & 0 & \\
 \downarrow & & \\
 \mathbb{1} & -1 &
 \end{array}$$

The composite of consecutive boundary maps is 0, due to the computation

$$U_1 \cdot (a - U_1) = 0 = (a - U_1) \cdot U_1,$$

and the fact that U_1 acts by 0 on $\mathbb{1}$. Moreover, this complex is acyclic, as one sees by considering the bases $1, U_1$ and $1, (a - U_1)$ of TL_2 . Thus the non-negative part of the complex above, which we denote by P_* , is a free resolution of the left TL_2 -module $\mathbb{1}$. Thus $\mathrm{Tor}_*^{\mathrm{TL}_2}(\mathbb{1}, \mathbb{1})$ and $\mathrm{Ext}_{\mathrm{TL}_2}^*(\mathbb{1}, \mathbb{1})$ are the homology of $\mathbb{1} \otimes_{\mathrm{TL}_2} P_*$ and the cohomology of $\mathrm{Hom}_{\mathrm{TL}_2}(P_*, \mathbb{1})$ respectively. Using the isomorphisms $\mathbb{1} \otimes_{\mathrm{TL}_2} \mathrm{TL}_2 \cong \mathbb{1}$, $a \otimes x \mapsto a \cdot x$ and $\mathrm{Hom}^{\mathrm{TL}_2}(\mathrm{TL}_2, \mathbb{1}) \cong \mathbb{1}$, $f \mapsto f(1)$ in every degree, and working out the induced boundary maps, we see that $\mathbb{1} \otimes_{\mathrm{TL}_2} P_*$ and $\mathrm{Hom}_{\mathrm{TL}_2}(P_*, \mathbb{1})$ are isomorphic to the complexes depicted below.

$$\begin{array}{ccc}
 \vdots & \vdots & \vdots \\
 \downarrow a & \uparrow a & \\
 R & R & 3 \\
 \downarrow 0 & \uparrow 0 & \\
 R & R & 2 \\
 \downarrow a & \uparrow a & \\
 R & R & 1 \\
 \downarrow 0 & \uparrow 0 & \\
 R & R & 0
 \end{array}$$

The homology and cohomology of these complexes are easily computed, and give the claim. \square

Proposition 7.2. *When $n = 2$ the Fineberg module satisfies $\mathcal{F}_2(a) \cong \mathbb{1}$, and the map $\mathbb{1} \otimes_{\text{TL}_2(a)} \mathcal{F}_2(a) \rightarrow \varepsilon_2 \cong \mathbb{1}$ is multiplication by a .*

Proof. We compute \mathcal{F}_2 explicitly in Example 4.11: $\mathcal{F}_2 \cong \langle a - U_1 \rangle \cong \mathbb{1}$. The map $\mathbb{1} \otimes_{\text{TL}_2} \mathcal{F}_2 \rightarrow \varepsilon_2 \cong \mathbb{1}$ is the composite map

$$\mathbb{1} \otimes_{\text{TL}_2} \mathcal{F}_2 \rightarrow \mathbb{1} \otimes_{\text{TL}_2} W(2)_1 = \mathbb{1} \otimes_{\text{TL}_2} (\text{TL}_2 \otimes_{\text{TL}_0} \mathbb{1}) \cong \mathbb{1}.$$

Under the central equality the basis element $a - U_1$ of $\mathcal{F}_2 \subset W(2)_1$ gets mapped to $a - U_1 = a$ in the tensor product. Therefore the composite map is given by multiplication by a , as required. \square

Corollary 7.3. *Suppose that $v \in R$ is a unit and that $a = v + v^{-1}$. Then the groups $\text{Tor}_i^{\text{TL}_2(a)}(\mathbb{1}, \mathbb{1})$ and $\text{Ext}_i^{\text{TL}_2(a)}(\mathbb{1}, \mathbb{1})$ are as described in Proposition 7.1.*

Proof. In the light of Proposition 7.2, the exact sequence from Theorem 5.1

$$0 \rightarrow \text{Tor}_2^{\text{TL}_2}(\mathbb{1}, \mathbb{1}) \rightarrow \mathbb{1} \otimes_{\text{TL}_2} \mathcal{F}_2 \rightarrow \mathbb{1} \rightarrow \text{Tor}_1^{\text{TL}_2}(\mathbb{1}, \mathbb{1}) \rightarrow 0$$

now becomes

$$0 \rightarrow \text{Tor}_2^{\text{TL}_2}(\mathbb{1}, \mathbb{1}) \rightarrow \mathbb{1} \otimes_{\text{TL}_2} \mathbb{1} \xrightarrow{a} \mathbb{1} \rightarrow \text{Tor}_1^{\text{TL}_2}(\mathbb{1}, \mathbb{1}) \rightarrow 0$$

from which one can compute $\text{Tor}_2^{\text{TL}_2}(\mathbb{1}, \mathbb{1}) = R_a$ and $\text{Tor}_1^{\text{TL}_2}(\mathbb{1}, \mathbb{1}) = R/aR$. For $i \geq 3$ we have the recursive formula

$$\text{Tor}_i^{\text{TL}_2}(\mathbb{1}, \mathbb{1}) = \text{Tor}_{i-2}^{\text{TL}_2}(\mathbb{1}, \mathcal{F}_2) \cong \text{Tor}_{i-2}^{\text{TL}_2}(\mathbb{1}, \mathbb{1})$$

which completes the proof. The Ext results similarly follow from Theorem 5.1. \square

8. HIGH-ACYCLICITY

In this final section we prove high connectivity of $W(n)$, Theorem E.

Theorem E. *$H_d(W(n))$ vanishes in degrees $d \leq (n - 2)$.*

8.1. A filtration. In this subsection we introduce a filtration of $W(n)$. We state a theorem relating the filtration quotients to $W(n - 1)$ (the proof of which is the topic of the next 3 subsections) and therefore by induction prove Theorem E.

Definition 8.1 (The filtration). We define a filtration F of $W(n)$,

$$F^0 \subseteq F^1 \subseteq \dots \subseteq F^n = W(n)$$

as follows.

- F^0 is defined to be the span of the elements of two kinds. We call elements of the first kind *basic elements* and these are of the form

$$x \otimes 1$$

in degrees i such that $-1 \leq i \leq n-2$, where x is represented by a monomial in the s_j not involving the letter s_1 . Elements of the second kind are those of the form

$$x \cdot (s_1 \cdots s_{n-i-1}) \otimes 1$$

in degrees i such that $0 \leq i \leq n-1$, where again x is represented by a monomial not involving the letter s_1 .

- F^k for $k \geq 1$ is defined to be the span of F^{k-1} together with terms of the form

$$x \cdot (s_1 \cdots s_{n-i-1+k}) \otimes 1$$

in degrees i such that $k \leq i \leq n-1$, where again x is represented by a monomial not involving s_1 .

Remark 8.2. Note that in the description of F^0 , it is possible for the product $s_1 \cdots s_{n-i-1}$ to be empty, i.e. the unit element, if the final index $(n-i-1)$ is zero ($i = n-1$). In contrast, in the description of F^k for $k \geq 1$, the product $s_1 \cdots s_{n-i-1+k}$ is never empty. This is one reason why it is important for us to treat F^0 quite separately from the other F^k , as is done in the remainder of this paper.

In Theorem 8.7 we show that each F^k is a subcomplex of $W(n)$. The fact that $F^n = W(n)$ will follow from Lemma 8.25.

Definition 8.3. Recall that the *cone* on a chain complex X (or, more precisely, the cone on the identity map of X) is the chain complex CX defined by $(CX)_i = X_i \oplus X_{i-1}$, and with differential defined by

$$d_{CX}^i(x, y) = (d_X^i(x) + y, -d_X^{i-1}(y)).$$

The *suspension* of a chain complex X is the complex ΣX defined by

$$(\Sigma X)_i = X_{i-1}$$

and with the same differential as X . The *truncation to degree p* of a chain complex X is the chain complex $\tau_p X$ defined by

$$(\tau_p X)_i = \begin{cases} X_i, & i \leq p \\ 0, & i > p \end{cases}$$

and with the same differential as X (in the relevant degrees).

Remark 8.4. Note that our definition of cone and suspension do not seem to match up very well. However, we have chosen our conventions in order to make the proof of the next theorem as direct as possible, and we believe that our choices are the best fit for this purpose.

Definition 8.5. Define the *shift map* σ to be the map

$$\sigma: \text{TL}_{n-1}(a) \rightarrow \text{TL}_n(a)$$

which sends each U_i to U_{i+1} for $1 \leq i \leq n-2$, and hence each s_i to s_{i+1} .

Lemma 8.6. *Each F^k consists of $\mathrm{TL}_{n-1}(a)$ -submodules of $W(n)$, where $\mathrm{TL}_{n-1}(a)$ acts via the shift map σ .*

Proof. Definition 8.1 defines each F^k as the span of certain ‘base elements’ of the form $y \otimes 1$ where $y \in \mathrm{TL}_n$ is represented by a monomial in the s_j subject to certain restrictions. Multiplying any such y on the left by any s_j for $1 < j \leq n-1$ does not affect whether it meets these restrictions. Since $s_j = \sigma(s_{j-1})$ for $1 < j \leq n-1$, this shows that the generators of TL_{n-1} send the base elements of each F^k to other base elements of F^k , and therefore F^k itself is stable under the action of TL_{n-1} . \square

Here is the main result of this section.

Theorem 8.7. *Each F^k is a subcomplex of $W(n)$. We identify*

$$F^0 \cong C(W(n-1)).$$

And for $k \geq 1$, we have

$$F^k/F^{k-1} \cong \tau_{n-1}\Sigma^{k+1}W(n-1).$$

Corollary 8.8 (Theorem E). *For each $n \geq 0$ the complex $W(n)$ is $(n-2)$ -acyclic, or in other words, its homology vanishes up to and including degree $(n-2)$.*

Proof. We prove this by induction on $n \geq 0$. One can verify the claim directly in the case $n = 0$. Fix $n \geq 1$ and suppose that the theorem has been proved for the previous case. Now $W(n)$ has the filtration $F^0 \subseteq F^1 \subseteq \dots \subseteq F^n$. We prove below that F^0 and all filtration quotients F^k/F^{k-1} are $(n-2)$ -acyclic, and then it follows (for example by using the short exact sequences $0 \rightarrow F^{k-1} \rightarrow F^k \rightarrow F^k/F^{k-1} \rightarrow 0$, or by using the spectral sequence of the filtration) that the same holds for $W(n)$ itself.

Observe that $F^0 \cong C(W(n-1))$, being isomorphic to a cone, is acyclic. Next, for $k \geq 1$ we have $F^k/F^{k-1} \cong \tau_{n-1}\Sigma^{k+1}W(n-1)$. The induction hypothesis states that $W(n-1)$ is $(n-3)$ -acyclic, so that $\Sigma^{k+1}W(n-1)$ is $(n-2+k)$ -acyclic and in particular $(n-2)$ -acyclic, so that $\tau_{n-1}\Sigma^{k+1}W(n-1)$ is also $(n-2)$ -acyclic. This completes the proof. \square

Remark 8.9 (Intuitions and motivations). The complex of planar injective words $W(n)$ is an analogue of the complex of injective words $\mathcal{C}(n)$, and Theorem E is the analogue for $W(n)$ of the well-known vanishing result for the homology of $\mathcal{C}(n)$; see Remark 4.3.

Our starting point in proving Theorem E was Kerz’s proof [Ker05] of the vanishing theorem for the homology of $\mathcal{C}(n)$. Kerz identifies within $\mathcal{C}(n)$ a subcomplex F^0 that is isomorphic to the cone $C(\mathcal{C}(n-1))$. This is then extended to a filtration $F^0 \subseteq F^1 \subseteq \dots \subseteq F^{n-1} \subseteq \mathcal{C}(n)$ in which each subsequent filtration quotient F^k/F^{k-1} is isomorphic to a direct sum of copies of the suspension $\Sigma^{k+1}\mathcal{C}(n-k-1)$. (In fact Kerz does not explicitly mention filtrations, but this is one way of framing his proof.) This permits an inductive proof of high-acyclicity as in Corollary 8.8.

Our proof of Theorem E began as an attempt to mimic Kerz's approach. There is an evident way to embed $W(n-1)$ into $W(n)$ — this is the span of the basic elements of F^0 — and this can be extended to an embedding of the cone $C(W(n-1))$ into $W(n)$ by considering the elements of the second kind in F^0 . The remainder of our proof is the result of attempting to extend this embedding to a complete filtration of $W(n)$. At this stage the parallels with [Ker05] begin to fail, but the Jones normal form gives us an extra tool. Using this we characterise the basis elements of $W(n)$ that are not in the image of the cone $C(W(n-1))$, and this characterisation gives a surprising separation into subcomplexes which 'look like' suspended and truncated copies of $W(n-1)$ — we build our filtration such that these are our filtration quotients F^k/F^{k-1} .

The final three subsections prove Theorem 8.7, by first setting up the required chain map for F^0 , then for F^k and then in the final section proving these chain maps are isomorphisms.

8.2. Proofs for F^0 . In this subsection we prove F^0 is a subcomplex of $W(n)$. We define a map from the cone $C(W(n-1))$ to F^0 and prove this is a well defined chain map.

Lemma 8.10. *F^0 is a subcomplex of $W(n)$.*

Proof. To prove the claim, we must take a generator of F^0 in degree i , and show that under the boundary map $d^i: W(n)_i \rightarrow W(n)_{i-1}$ this generator is mapped into F^0 . Since d^i is the alternating sum $d_0^i - d_1^i + \cdots + (-1)^i d_i^i$, it will suffice to fix j in the range $0 \leq j \leq i$, and show that d_j^i sends our generator into F^0 . Recall from Definition 4.1 the definition of d_j^i :

$$d_j^i(y \otimes r) = y \cdot (s_{n-i+j-1} \cdots s_{n-i}) \otimes \lambda^{-j} r.$$

Generators of F^0 come in two kinds. The first kind are the basic elements $x \otimes 1$ in degrees $-1 \leq i \leq n-2$ where x is represented by a monomial not featuring the letter s_1 . The map d_j^i only introduces a letter s_1 in the case $i = n-1$, which is excluded here, so that $d_j^i(x \otimes 1)$ is again a basic element and therefore also lies in F^0 .

The second kind of generators of F^0 are elements

$$x \cdot (s_1 \cdots s_{n-i-1}) \otimes 1$$

in degrees $0 \leq i \leq n-1$, where x is represented by a monomial not involving s_1 . In the case $j = 0$, we have

$$d_0^i(x \cdot (s_1 \cdots s_{n-i-1}) \otimes 1) = x \cdot (s_1 \cdots s_{n-i-1}) \otimes 1$$

but this lies in $W(n)_{i-1} = \mathrm{TL}_n \otimes_{\mathrm{TL}_{n-i}} \mathbb{1}$, hence is equal to $x \otimes \lambda^{n-i-1}$, and since x is represented by a monomial not involving s_1 , this does indeed lie in F^0 . (This

argument includes the special case $i = n - 1$, where the product $s_1 \cdots s_{n-i-1}$ is empty, but this clearly creates no issues.) In the case $j \geq 1$, we have

$$\begin{aligned} d_j^i(x \cdot (s_1 \cdots s_{n-i-1}) \otimes 1) &= x \cdot (s_1 \cdots s_{n-i-1}) \cdot (s_{n-i+j-1} \cdots s_{n-i}) \otimes \lambda^{-j} \\ &= x \cdot (s_1 \cdots s_{n-i-1}) \cdot (s_{n-i+j-1} \cdots s_{n-i+1}) \cdot s_{n-i} \otimes \lambda^{-j} \\ &= x \cdot (s_{n-i+j-1} \cdots s_{n-i+1}) \cdot (s_1 \cdots s_{n-i-1}) \cdot s_{n-i} \otimes \lambda^{-j} \\ &= x \cdot (s_{n-i+j-1} \cdots s_{n-i+1}) \cdot (s_1 \cdots s_{n-i}) \otimes \lambda^{-j} \\ &= (x \cdot (s_{n-i+j-1} \cdots s_{n-i+1})) \cdot (s_1 \cdots s_{n-(i-1)-1}) \otimes \lambda^{-j} \end{aligned}$$

which lies in F^0 since $x \cdot (s_{n-i+j-1} \cdots s_{n-i+1})$, does not involve the letter s_1 and so $d_j^i(x \cdot (s_1 \cdots s_{n-i-1}) \otimes 1)$ is a scalar multiple of a generator of F^0 , and thus in F^0 , as required. \square

Definition 8.11. Define a map

$$\Phi^0: C(W(n-1)) \longrightarrow F^0$$

as follows. Recall that

$$\begin{aligned} C(W(n-1))_i &= W(n-1)_i \oplus W(n-1)_{i-1} \\ &= (\mathrm{TL}_{n-1}(a) \otimes_{\mathrm{TL}_{n-i-2}(a)} \mathbb{1}) \oplus (\mathrm{TL}_{n-1}(a) \otimes_{\mathrm{TL}_{n-i-1}(a)} \mathbb{1}) \end{aligned}$$

and that

$$F_i^0 \subseteq W(n)_i = \mathrm{TL}_n(a) \otimes_{\mathrm{TL}_{n-i-1}(a)} \mathbb{1}.$$

We define Φ^0 in degree i by the rule

$$\Phi_i^0(x \otimes \alpha, y \otimes \beta) = \xi_i(x \otimes \alpha) + \eta_i(y \otimes \beta)$$

where

$$\begin{aligned} \xi_i: W(n-1)_i &\rightarrow W(n)_i \\ x \otimes \alpha &\mapsto \sigma(x) \otimes \lambda^{n-1} \alpha \end{aligned}$$

and

$$\begin{aligned} \eta_i: W(n-1)_{i-1} &\rightarrow W(n)_i \\ y \otimes \beta &\mapsto \sigma(y) \cdot (s_1 \cdots s_{n-i-1}) \otimes \lambda^i \beta. \end{aligned}$$

It is simple to check that the image of both maps lies in F_i^0 .

Lemma 8.12. *The maps ξ_i and η_i are well defined.*

Proof. In the case of ξ_i this is simple to verify, as the map $\sigma: \mathrm{TL}_{n-1} \rightarrow \mathrm{TL}_n$ is in fact a map of right-modules with respect to the map of algebras $\sigma: \mathrm{TL}_{n-i-2} \rightarrow \mathrm{TL}_{n-i-1}$.

In the case of η_i , the definition of $\eta_i(y \otimes \beta)$ as presented depends on y and β themselves, and we must check that it depends only on $y \otimes \beta$. Thus we must show that

$$\eta_i(ys_j \otimes \beta) = \eta_i(y \otimes \lambda\beta)$$

whenever $1 \leq j \leq n - i - 2$. And indeed

$$\begin{aligned}
\eta_i(ys_j \otimes \beta) &= \sigma(ys_j) \cdot (s_1 \cdots s_{n-i-1}) \otimes \lambda^i \beta \\
&= \sigma(y) \cdot s_{j+1} \cdot (s_1 \cdots s_{n-i-1}) \otimes \lambda^i \beta \\
&= \sigma(y) \cdot (s_1 \cdots s_{n-i-1}) \cdot s_j \otimes \lambda^i \beta \\
&= \sigma(y) \cdot (s_1 \cdots s_{n-i-1}) \otimes \lambda^{i+1} \beta \\
&= \eta_i(y \otimes \lambda \beta)
\end{aligned}$$

where the third equality holds since $2 \leq j + 1 \leq n - i - 1$ (a simple way to see this is to draw the s_i as braids), and the fourth holds since $j \leq n - i - 2$ and the tensor product is over TL_{n-i-1} . \square

Lemma 8.13. *The ξ_i and η_i interact with the boundary maps of $W(n)$ in the following way:*

- (1) $d_j^i \circ \xi_i = \xi_{i-1} \circ d_j^i$ for i in the range $-1 \leq i \leq n - 2$ and j in the range $0 \leq j \leq i$.
- (2) $d_0^i \circ \eta_i = \xi_{i-1}$ for i in the range $0 \leq i \leq n - 1$.
- (3) $d_{j+1}^i \circ \eta_i = \eta_{i-1} \circ d_j^{i-1}$ for i in the range $0 \leq i \leq n - 1$ and j in the range $0 \leq j \leq i - 1$.

Proof. For the first point, we have:

$$\begin{aligned}
d_j^i(\xi_i(x \otimes \alpha)) &= d_j^i(\sigma(x) \otimes \lambda^{n-1} \alpha) \\
&= \sigma(x) \cdot (s_{n-i+j-1} \cdots s_{n-i}) \otimes \lambda^{-j} \lambda^{n-1} \alpha \\
&= \sigma(x \cdot (s_{n-i+j-2} \cdots s_{n-i-1})) \otimes \lambda^{-j} \lambda^{n-1} \alpha \\
&= \xi_{i-1}(x \cdot (s_{n-i+j-2} \cdots s_{n-i-1}) \otimes \lambda^{-j} \alpha) \\
&= \xi_{i-1}(x \cdot (s_{(n-1)-i+j-1} \cdots s_{(n-1)-i}) \otimes \lambda^{-j} \alpha) \\
&= \xi_{i-1}(d_j^i(x \otimes \alpha)).
\end{aligned}$$

For the second point, we have:

$$\begin{aligned}
d_j^i(\eta_i(y \otimes \beta)) &= d_0^i(\sigma(y) \cdot (s_1 \cdots s_{n-i-1}) \otimes \lambda^i \beta) \\
&= \sigma(y) \cdot (s_1 \cdots s_{n-i-1}) \otimes \lambda^i \beta \\
&= \sigma(y) \otimes \lambda^{n-i-1} \lambda^i \beta \\
&= \sigma(y) \otimes \lambda^{n-1} \beta \\
&= \xi_{i-1}(y \otimes \beta),
\end{aligned}$$

where the third equality holds because the terms lie in $W(n)_{i-1} = \mathbb{T}\mathbb{L}_n \otimes_{\mathbb{T}\mathbb{L}_{n-i}} \mathbb{1}$. And for the third point we have:

$$\begin{aligned}
d_{j+1}^i \eta_i(y \otimes \beta) &= d_{j+1}^i(\sigma(y) \cdot (s_1 \cdots s_{n-i-1}) \otimes \lambda^i \beta) \\
&= \sigma(y) \cdot (s_1 \cdots s_{n-i-1}) \cdot (s_{n-i+(j+1)-1} \cdots s_{n-i}) \otimes \lambda^{-j-1} \lambda^i \beta \\
&= \sigma(y) \cdot (s_1 \cdots s_{n-i-1}) \cdot (s_{n-i+j} \cdots s_{n-i+1}) \cdot s_{n-i} \otimes \lambda^{i-j-1} \beta \\
&= \sigma(y) \cdot (s_{n-i+j} \cdots s_{n-i+1}) \cdot (s_1 \cdots s_{n-i}) \otimes \lambda^{i-j-1} \beta \\
&= \sigma(y \cdot (s_{n-i+j-1} \cdots s_{n-i})) \cdot (s_1 \cdots s_{n-(i-1)-1}) \otimes \lambda^{i-1} \lambda^{-j} \beta \\
&= \eta_{i-1}(y \cdot (s_{n-i+j-1} \cdots s_{n-i}) \otimes \lambda^{-j} \beta) \\
&= \eta_{i-1}(y \cdot (s_{(n-1)-(i-1)+j-1} \cdots s_{(n-1)-(i-1)}) \otimes \lambda^{-j} \beta) \\
&= \eta_{i-1}(d_j^{i-1}(y \otimes \beta))
\end{aligned}$$

where for the final equality we recall that the source of η_{i-1} is $W(n-1)_{i-2}$. \square

Lemma 8.14. Φ^0 is a chain map.

Proof. Referring to the definition of the differential on $C(W(n-1))$ (Definition 8.3), we see that in order to check that $d^i \circ \Phi_i^0 = \Phi_{i-1}^0 \circ d^i$, it is enough to show that $d^i \circ \xi_i(x \otimes \alpha) = \xi_{i-1}(d^i(x \otimes \alpha))$ and $d^i \circ \eta_i(y \otimes \beta) = \xi_{i-1}(y \otimes \beta) - \eta_{i-1}(d^{i-1}(y \otimes \beta))$. Using the previous lemma, for the first we have

$$\begin{aligned}
d^i \circ \xi_i(x \otimes \alpha) &= \sum_{j=0}^i (-1)^j d_j^i(\xi_i(x \otimes \alpha)) \\
&= \sum_{j=0}^i (-1)^j \xi_{i-1}(d_j^i(x \otimes \alpha)) \\
&= \xi_{i-1} \left(\sum_{j=0}^i (-1)^j d_j^i(x \otimes \alpha) \right) \\
&= \xi_{i-1}(d^i(x \otimes \alpha)).
\end{aligned}$$

And for the second we have

$$\begin{aligned}
d^i \circ \eta_i(y \otimes \beta) &= \sum_{j=0}^i (-1)^j d_j^i(\eta_i(y \otimes \beta)) \\
&= d_0^i(\eta_i(y \otimes \beta)) - \sum_{j=0}^{i-1} (-1)^j d_{j+1}^i \eta_i(y \otimes \beta) \\
&= \xi_{i-1}(y \otimes \beta) - \sum_{j=0}^{i-1} (-1)^j \eta_{i-1} d_j^{i-1}(y \otimes \beta) \\
&= \xi_{i-1}(y \otimes \beta) - \eta_{i-1} \left(\sum_{j=0}^{i-1} (-1)^j d_j^{i-1}(y \otimes \beta) \right) \\
&= \xi_{i-1}(y \otimes \beta) - \eta_{i-1}(d^{i-1}(y \otimes \beta)). \quad \square
\end{aligned}$$

8.3. Proofs for F^k , $k \geq 1$. In this subsection we prove, for $k \geq 1$, that F^k is a subcomplex of $W(n)$. We define a map from $\tau_{n-1} \Sigma^{k+1} W(n-1)$ to F^k/F^{k-1} and prove this is a well defined chain map. We start off with some elementary lemmas involving the s_j , which we require for later proofs.

Lemma 8.15. *Let $m \geq 1$ and $p \leq m$. Then*

$$s_1 \cdots s_m \cdots s_p = (s_m \cdots s_{p+1}) \cdot (s_1 \cdots s_m).$$

In the case $m = p$ the product $s_m \cdots s_{p+1}$ is empty and therefore equal to 1.

Lemma 8.16. *Let $p \geq 1$, $q \geq r \geq 1$. Then the product $(s_1 \cdots s_p) \cdot (s_q \cdots s_r)$ can be described as follows.*

(1) *When $r - 1 \leq p \leq q - 1$,*

$$(s_1 \cdots s_p) \cdot (s_q \cdots s_r) = (s_q \cdots s_{r+1}) \cdot (s_1 \cdots s_{p+1}).$$

(2) *When $p = q$, $(s_1 \cdots s_p) \cdot (s_q \cdots s_r)$ is a linear combination of terms of the form $(s_t \cdots s_{r+1}) \cdot (s_1 \cdots s_t)$ for $p \geq t \geq r + 1$, as well as $s_1 \cdots s_r$ and $s_1 \cdots s_{r-1}$.*

(3) *When $p \geq q + 1$,*

$$(s_1 \cdots s_p) \cdot (s_q \cdots s_r) = (s_{q+1} \cdots s_{r+1}) \cdot (s_1 \cdots s_p).$$

Proof. When $r - 1 \leq p \leq q - 1$,

$$\begin{aligned}
(s_1 \cdots s_p) \cdot (s_q \cdots s_r) &= (s_1 \cdots s_p) \cdot (s_q \cdots s_{p+2}) \cdot (s_{p+1} \cdots s_r) \\
&= (s_q \cdots s_{p+2}) \cdot (s_1 \cdots s_p) \cdot (s_{p+1} \cdots s_r) \\
&= (s_q \cdots s_{p+2}) \cdot (s_1 \cdots s_{p+1} \cdots s_r) \\
&= (s_q \cdots s_{p+2}) \cdot (s_{p+1} \cdots s_{r+1}) \cdot (s_1 \cdots s_{p+1}) \\
&= (s_q \cdots s_{r+1}) \cdot (s_1 \cdots s_{p+1}),
\end{aligned}$$

where we used Lemma 8.15 to obtain the fourth equality.

When $p = q$, we claim that

$$(s_1 \cdots s_p) \cdot (s_q \cdots s_r) = (s_1 \cdots s_p) \cdot (s_p \cdots s_r)$$

is a linear combination of terms of the form $(s_t \cdots s_{r+1}) \cdot (s_1 \cdots s_t)$ for $p \geq t \geq r+1$, as well as $s_1 \cdots s_r$ and $s_1 \cdots s_{r-1}$. We will prove this claim by induction on the difference $p - r$. When $p - r = 0$, we have

$$(s_1 \cdots s_p) \cdot (s_p \cdots s_r) = s_1 \cdots s_p \cdot s_p.$$

Now since s_p^2 is a linear combination of s_p and 1, this is a linear combination of $s_1 \cdots s_p = s_1 \cdots s_r$ and $s_1 \cdots s_{p-1} = s_1 \cdots s_{r-1}$ as required. Now let $p - r \geq 1$, and assume that the claim holds for all smaller values. Then

$$(s_1 \cdots s_p) \cdot (s_p \cdots s_r) = (s_1 \cdots s_{p-1}) \cdot s_p^2 \cdot (s_{p-1} \cdots s_r)$$

is a linear combination of

$$\begin{aligned} (s_1 \cdots s_{p-1}) \cdot s_p \cdot (s_{p-1} \cdots s_r) &= s_1 \cdots s_p \cdots s_r \\ &= (s_p \cdots s_{r+1}) \cdot (s_1 \cdots s_p) \end{aligned}$$

(where we used Lemma 8.15) and

$$(s_1 \cdots s_{p-1}) \cdot (s_{p-1} \cdots s_r).$$

The former is $(s_t \cdots s_{r+1}) \cdot (s_1 \cdots s_t)$ in the case $t = p$, while the induction hypothesis tells us that the latter is a linear combination of $(s_t \cdots s_{r+1}) \cdot (s_1 \cdots s_t)$ for $p - 1 \geq t \geq r + 1$, as well as $s_1 \cdots s_r$ and $s_1 \cdots s_{r-1}$. This completes the proof of the claim.

When $p \geq q + 1$,

$$\begin{aligned} (s_1 \cdots s_p) \cdot (s_q \cdots s_r) &= (s_1 \cdots s_{q+1}) \cdot (s_{q+2} \cdots s_p) \cdot (s_q \cdots s_r) \\ &= (s_1 \cdots s_{q+1}) \cdot (s_q \cdots s_r) \cdot (s_{q+2} \cdots s_p) \\ &= (s_1 \cdots s_{q+1} \cdots s_r) \cdot (s_{q+2} \cdots s_p) \\ &= (s_{q+1} \cdots s_{r+1}) \cdot (s_1 \cdots s_{q+1}) \cdot (s_{q+2} \cdots s_p) \\ &= (s_{q+1} \cdots s_{r+1}) \cdot (s_1 \cdots s_p) \end{aligned}$$

(where we again used Lemma 8.15 to obtain the fourth equality) as required. \square

Lemma 8.17. *For $k \geq 1$, F^k is a subcomplex of $W(n)$.*

Proof. We fix $k \geq 1$ and take a generator of F^k/F^{k-1} in degree i , where $k \leq i \leq n - 1$, and show that the boundary map $d^i: W(n)_i \rightarrow W(n)_{i-1}$ sends our generator into F^k . Since d is the alternating sum $d_0^i - d_1^i + \cdots + (-1)^i d_i^i$, it will suffice to fix j in the range $0 \leq j \leq i$, and show that d_j^i sends our generator into F^k . Recall from Definition 8.1 that our generator of F^k/F^{k-1} in degree i is $x \cdot (s_1 \cdots s_{n-i-1+k}) \otimes 1$, where x does not involve the letter s_1 . Note that

$$(n - i - 1 + k) = (n - 1) - i + k \geq (n - 1) - (n - 1) + 1 = 1,$$

so that the product $(s_1 \cdots s_{n-i-1+k})$ is not empty. We have

$$d_j^i(x \cdot (s_1 \cdots s_{n-i-1+k}) \otimes 1) = x \cdot (s_1 \cdots s_{n-i-1+k}) \cdot (s_{n-i-1+j} \cdots s_{n-i}) \otimes \lambda^{-j},$$

where the factor $(s_{n-i-1+j} \cdots s_{n-i})$ can be empty, in the case $j = 0$.

- First we consider the case $j = 0$. We find that

$$\begin{aligned} d_0^i(x \cdot (s_1 \cdots s_{n-i-1+k}) \otimes 1) &= x \cdot (s_1 \cdots s_{n-i-1+k}) \otimes 1 \\ &= x \cdot (s_1 \cdots s_{n-(i-1)-1+(k-1)}) \otimes 1 \end{aligned}$$

lies in F^{k-1} , and therefore in F^k as required.

- Now we consider the case $1 \leq j \leq (k-1)$. Then $(n-i-1+k) \geq (n-i-1+j)+1$, so that the third item of Lemma 8.16 applies and shows that

$$\begin{aligned} d_j^i(x \cdot (s_1 \cdots s_{n-i-1+k}) \otimes 1) &= x \cdot (s_1 \cdots s_{n-i-1+k}) \cdot (s_{n-i-1+j} \cdots s_{n-i}) \otimes \lambda^{-j} \\ &= x \cdot (s_{n-i+j} \cdots s_{n-i+1}) \cdot (s_1 \cdots s_{n-i-1+k}) \otimes \lambda^{-j} \\ &= x \cdot (s_{n-i+j} \cdots s_{n-i+1}) \cdot (s_1 \cdots s_{n-(i-1)-1+(k-1)}) \otimes \lambda^{-j} \end{aligned}$$

Since $n-i+1 \geq n-(n-1)+1=2$, the word $(s_{n-i+j} \cdots s_{n-i+1})$ does not involve s_1 , and consequently the element above lies in F^{k-1} , and therefore in F^k .

- Now we consider the case $j = k$. Then $(n-i-1+k) = (n-i-1+j)$ and so the second item of Lemma 8.16 applies and shows that

$$\begin{aligned} d_k^i(x \cdot (s_1 \cdots s_{n-i-1+k}) \otimes 1) &= \\ &= x \cdot (s_1 \cdots s_{n-i-1+k}) \cdot (s_{n-i-1+k} \cdots s_{n-i}) \otimes \lambda^{-k} \end{aligned}$$

is a linear combination of terms

$$x \cdot (s_t \cdots s_{n-i+1}) \cdot (s_1 \cdots s_t) \otimes \lambda^{-k}$$

for t in the range

$$(n-i+1) \leq t \leq (n-i-1+k) = (n-(i-1)-1+(k-1))$$

together with

$$x \cdot (s_1 \cdots s_{n-(i-1)-1}) \otimes \lambda^{-k}$$

and

$$x \cdot (s_1 \cdots s_{n-(i-1)-2}) \otimes \lambda^{-k} = x \otimes \lambda^{-k}.$$

Now $(s_t \cdots s_{n-i+1})$ does not involve s_1 , so the first of these terms lies in F^{k-1} , while the second and third lie in F^0 . So altogether we have the required result.

- Now we consider the case $k + 1 \leq j$. Here we have

$$(n - i - 1) \leq (n - i - 1 + k) + 1 \leq (n - i - 1 + j),$$

so that the first item of Lemma 8.16 applies and shows that

$$\begin{aligned} d_j^i(x \cdot (s_1 \cdots s_{n-i-1+k}) \otimes 1) &= x \cdot (s_1 \cdots s_{n-i-1+k}) \cdot (s_{n-i-1+j} \cdots s_{n-i}) \otimes \lambda^{-j} \\ &= x \cdot (s_{n-i-1+j} \cdots s_{n-i+1}) \cdot (s_1 \cdots s_{n-i-1+k+1}) \otimes \lambda^{-j} \\ &= x \cdot (s_{n-i-1+j} \cdots s_{n-i+1}) \cdot (s_1 \cdots s_{n-(i-1)-1+k}) \otimes \lambda^{-j}. \end{aligned}$$

Since $(s_{n-i-1+j} \cdots s_{n-i+1})$ does not involve s_1 , the element above lies in F^k as required. \square

Definition 8.18. Define a map

$$\Psi^k : \tau_{n-1}\Sigma^{k+1}W(n-1) \longrightarrow F^k/F^{k-1}$$

as follows. Note that for i in the range $k \leq i \leq (n-1)$,

$$\begin{aligned} [\tau_{n-1}\Sigma^{k+1}W(n-1)]_i &= W(n-1)_{i-k-1} \\ &= \text{TL}_{n-1}(a) \otimes_{\text{TL}_{(n-1)-(i-k-1)-1}(a)} \mathbb{1} \\ &= \text{TL}_{n-1}(a) \otimes_{\text{TL}_{n-i-1+k}(a)} \mathbb{1}, \end{aligned}$$

while $(F^k/F^{k-1})_i$ is a quotient of $\text{TL}_n(a) \otimes_{\text{TL}_{n-i-1}(a)} \mathbb{1}$. Define the degree i part of Ψ to be the map

$$\Psi_i^k : \text{TL}_{n-1}(a) \otimes_{\text{TL}_{n-i-1+k}(a)} \mathbb{1} \longrightarrow (F^k/F^{k-1})_i$$

given by

$$\Psi_i^k : x \otimes \alpha \longmapsto (-1)^{-i(k+1)} \sigma(x) \cdot (s_1 \cdots s_{n-i-1+k}) \otimes \lambda^i \alpha.$$

For later convenience, we will denote by ψ_i^k the map

$$\psi_i^k : x \otimes \alpha \longmapsto \sigma(x) \cdot (s_1 \cdots s_{n-i-1+k}) \otimes \lambda^i \alpha,$$

so that $\Psi_i^k = (-1)^{-i(k+1)} \psi_i^k$.

Lemma 8.19. *The map ψ_i^k is well defined (and the same therefore holds for Ψ_i^k).*

Proof. As presented above, the value of $\psi_i^k(x \otimes \alpha)$ depends on the choices of x and α , rather than on $x \otimes \alpha$. So to check that ψ_i^k is well-defined, we must check that $\psi_i^k(xs_p \otimes \alpha) = \psi_i^k(x \otimes \lambda\alpha)$ whenever $p \leq (n - i - 1 + k) - 1$. Let us write $q = (n - i - 1 + k)$, so that $p \leq q - 1$. (In particular we are assuming that $q \geq 2$.) Now

$$\begin{aligned} \psi_i^k(xs_p \otimes \alpha) &= \sigma(xs_p) \cdot (s_1 \cdots s_q) \otimes \lambda^i \alpha \\ &= \sigma(x) \cdot s_{p+1} \cdot (s_1 \cdots s_q) \otimes \lambda^i \alpha \\ &= \sigma(x) \cdot s_{p+1} \cdot (s_1 \cdots s_{p-1}) \cdot (s_p s_{p+1}) \cdot (s_{p+2} \cdots s_q) \otimes \lambda^i \alpha \\ &= \sigma(x) \cdot (s_1 \cdots s_{p-1}) \cdot (s_{p+1} s_p s_{p+1}) \cdot (s_{p+2} \cdots s_q) \otimes \lambda^i \alpha. \end{aligned}$$

Recall from Definition 2.21 that

$$s_{p+1}s_p s_{p+1} = \lambda s_p s_{p+1} + \lambda s_{p+1} s_p - \lambda^2 s_p - \lambda^2 s_{p+1} + \lambda^3.$$

Now

$$\begin{aligned} (s_1 \cdots s_{p-1}) \cdot (s_p s_{p+1}) \cdot (s_{p+2} \cdots s_q) &= (s_1 \cdots s_q) \\ (s_1 \cdots s_{p-1}) \cdot (s_{p+1} s_p) \cdot (s_{p+2} \cdots s_q) &= (s_{p+1} \cdots s_q) \cdot (s_1 \cdots s_p) \\ (s_1 \cdots s_{p-1}) \cdot s_p \cdot (s_{p+2} \cdots s_q) &= (s_{p+2} \cdots s_q) \cdot (s_1 \cdots s_p) \\ (s_1 \cdots s_{p-1}) \cdot s_{p+1} \cdot (s_{p+2} \cdots s_q) &= (s_{p+1} \cdots s_q) \cdot (s_1 \cdots s_{p-1}) \\ (s_1 \cdots s_{p-1}) \cdot 1 \cdot (s_{p+2} \cdots s_q) &= (s_{p+2} \cdots s_q) \cdot (s_1 \cdots s_{p-1}) \end{aligned}$$

so it follows that

$$\begin{aligned} \psi_i^k(x s_p \otimes \alpha) &= \sigma(x) \cdot (s_1 \cdots s_q) \otimes \lambda^{i+1} \alpha \\ &\quad + \sigma(x) \cdot (s_{p+1} \cdots s_q) \cdot (s_1 \cdots s_p) \cdot \otimes \lambda^{i+1} \alpha \\ &\quad - \sigma(x) \cdot (s_{p+2} \cdots s_q) \cdot (s_1 \cdots s_p) \otimes \lambda^{i+2} \alpha \\ &\quad - \sigma(x) \cdot (s_{p+1} \cdots s_q) \cdot (s_1 \cdots s_{p-1}) \otimes \lambda^{i+2} \alpha \\ &\quad + \sigma(x) \cdot (s_{p+2} \cdots s_q) \cdot (s_1 \cdots s_{p-1}) \otimes \lambda^{i+3} \alpha. \end{aligned}$$

Now $p < n - i - 1 + k$, which means that the final four terms above all lie in F^{k-1} , so that in F^k/F^{k-1} we have

$$\begin{aligned} \psi_i^k(x s_p \otimes \alpha) &= \sigma(x) \cdot (s_1 \cdots s_q) \otimes \lambda^{i+1} \alpha \\ &= \sigma(x) \cdot (s_1 \cdots s_{n-i-1+k}) \otimes \lambda^{i+1} \alpha \\ &= \psi_i^k(x \otimes \lambda \alpha) \end{aligned}$$

as required. \square

Lemma 8.20. *Let $k \geq 1$ and let $k \leq i \leq n - 1$. Then for j in the range $j \geq k + 1$ we have $\psi_{i-1}^k \circ d_{j-k-1}^{i-k-1} = d_j^i \circ \psi_i^k$.*

Proof. Let $x \otimes \alpha \in W(n-1)_{i-k-1} = \text{TL}_{n-1} \otimes_{\text{TL}_{n-i-1+k}} \mathbb{1}$. Then

$$\begin{aligned} d_j^i(\psi_i^k(x \otimes \alpha)) &= d_j^i(\sigma(x) \cdot (s_1 \cdots s_{n-i-1+k}) \otimes \lambda^i \alpha) \\ &= \sigma(x) \cdot (s_1 \cdots s_{n-i-1+k}) \cdot (s_{n-i+j-1} \cdots s_{n-i}) \otimes \lambda^{i-j} \alpha. \end{aligned}$$

Since $(n - i - 1) \leq (n - i - 1 + k) + 1 \leq (n - i + j - 1)$, we may apply the first part of Lemma 8.16 to obtain

$$\begin{aligned}
d_j^i(\psi_i^k(x \otimes \alpha)) &= \sigma(x) \cdot (s_1 \cdots s_{n-i-1+k}) \cdot (s_{n-i+j-1} \cdots s_{n-i}) \otimes \lambda^{i-j} \alpha \\
&= \sigma(x) \cdot (s_{n-i+j-1} \cdots s_{n-i+1}) \cdot (s_1 \cdots s_{n-i+k}) \otimes \lambda^{i-j} \alpha \\
&= \sigma(x) \cdot (s_{n-i+j-1} \cdots s_{n-i+1}) \cdot (s_1 \cdots s_{n-(i-1)-1+k}) \otimes \lambda^{(i-1)} \lambda^{1-j} \alpha \\
&= \sigma(x \cdot (s_{n-i+j-2} \cdots s_{n-i})) \cdot (s_1 \cdots s_{n-(i-1)-1+k}) \otimes \lambda^{(i-1)} \lambda^{1-j} \alpha \\
&= \psi_{i-1}^k(x \cdot (s_{n-i+j-2} \cdots s_{n-i}) \otimes \lambda^{1-j} \alpha).
\end{aligned}$$

In the last line of the above computation, $x \cdot (s_{n-i+j-2} \cdots s_{n-i}) \otimes \lambda^{1-j} \alpha$ is an element of $W(n-1)_{(i-1)-k-1} = \text{TL}_{n-1} \otimes_{\text{TL}_{n-i+k}} \mathbb{1}$, so we have

$$\begin{aligned}
x \cdot (s_{n-i+j-2} \cdots s_{n-i}) \otimes \lambda^{1-j} \alpha &= x \cdot (s_{n-i+j-2} \cdots s_{n-i+k}) \cdot (s_{n-i+k-1} \cdots s_{n-i}) \otimes \lambda^{1-j} \alpha \\
&= x \cdot (s_{n-i+j-2} \cdots s_{n-i+k}) \otimes \lambda^k \lambda^{1-j} \alpha \\
&= x \cdot (s_{n-i+j-2} \cdots s_{n-i+k}) \otimes \lambda^{-(j-k-1)} \alpha.
\end{aligned}$$

Thus

$$\begin{aligned}
d_j^i(\psi_i^k(x \otimes \alpha)) &= \psi_{i-1}^k(x \cdot (s_{n-i+j-2} \cdots s_{n-i}) \otimes \lambda^{1-j} \alpha) \\
&= \psi_{i-1}^k(x \cdot (s_{n-i+j-2} \cdots s_{n-i+k}) \otimes \lambda^{-(j-k-1)} \alpha) \\
&= \psi_{i-1}^k(x \cdot (s_{(n-1)-(i-k-1)+(j-k-1)-1} \cdots s_{(n-1)-(i-k-1)}) \otimes \lambda^{-(j-k-1)} \alpha) \\
&= \psi_{i-1}^k(d_{j-k-1}^{i-k-1}(x \otimes \alpha))
\end{aligned}$$

as required. \square

Corollary 8.21. Ψ^k is a chain map.

Proof. The boundary map of $\tau_{n-1} \Sigma^{k+1} W(n-1)$ is given in degree i by the boundary map $d^{i-k-1}: W(n-1)_{i-k-1} \rightarrow W(n-1)_{i-k-2}$, which is itself given by the formula $\sum_{j=0}^{i-k-1} (-1)^j d_j^{i-k-1}$.

The boundary map of F^k/F^{k-1} is given in degree i by the boundary map of $W(n)$ in degree i , which is the alternating sum $\sum_{j=0}^i (-1)^j d_j^i$. However, the proof of Lemma 8.17 shows that d_0^i, \dots, d_k^i all send F^k into F^{k-1} , and hence that they vanish in the quotient F^k/F^{k-1} . Thus the boundary map of F^k/F^{k-1} is $\sum_{j=k+1}^i (-1)^j d_j^i$. It follows that

$$\begin{aligned}
d^i \circ \Psi_i^k &= \sum_{j=k+1}^i (-1)^j d_j^i \circ [(-1)^{-i(k+1)} \psi_i^k] \\
&= \sum_{j=k+1}^i (-1)^{j-i(k+1)} \psi_{i-1}^k \circ d_{j-k-1}^{i-k-1} \\
&= \sum_{j=0}^{i-k-1} (-1)^{j+(k+1)-i(k+1)} \psi_{i-1}^k \circ d_j^{i-k-1} \\
&= [(-1)^{-(i-1)(k+1)} \psi_{i-1}^k] \circ \sum_{j=0}^{i-k-1} (-1)^j \psi_{i-1}^k \circ d_j^{i-k-1} \\
&= \Psi_{i-1}^k \circ d^{i-k-1}
\end{aligned}$$

as required. \square

8.4. Proof of Theorem 8.7. In this subsection we prove Theorem 8.7, which in turn completes the proof of Theorem E.

We begin by finding a basis for each part of the filtration in terms of the Jones normal form. This is done in Lemma 8.25 below, after some preliminary work.

Lemma 8.22. *Any word in the s_i not containing s_1 is a linear combination of words in the U_i , none of which involve U_1 . Conversely, any word in the U_i not containing U_1 is a linear combination of words in the s_i not containing s_1 .*

Proof. Recall from Definition 2.21 that $s_i = \lambda + \mu U_i$, where λ and μ are both units in the ground ring, so that $U_i = -\mu^{-1}\lambda + \mu^{-1}s_i$. The claim follows immediately. \square

Lemma 8.23. *For $1 \leq p \leq n-1$, the word $s_1 \dots s_p$ written in terms of the U_i generators is equal to $\mu^p U_1 \dots U_p$, plus a linear combination of scalar multiples—by units—of words w in the U_i with the following properties:*

- $i(w) \geq 2$ and $t(w) \leq p$ or
- $i(w) = 1$ and $t(w) < p$.

In particular only the summand $w = \mu^p U_1 \dots U_p$ satisfies $i(w) = 1$ and $t(w) = p$.

Proof. Using $s_i = \lambda + \mu U_i$ and multiplying out brackets gives the following:

$$\begin{aligned}
s_1 \dots s_p &= \sum_{r=0}^p \sum_{(1 \leq i_1 \leq \dots \leq i_r \leq p)} \lambda^{p-r} \mu^r U_{i_1} U_{i_2} \dots U_{i_r} \\
&= \mu^p U_1 \dots U_p + \sum_{r=0}^{p-1} \sum_{(1 \leq i_1 \leq \dots \leq i_r \leq p)} \lambda^{p-r} \mu^r U_{i_1} U_{i_2} \dots U_{i_r}
\end{aligned}$$

If $r = 0$ the term is a scalar, which has index ∞ by convention (thus the first point is satisfied). Suppose $0 < r < p$. Then if $i_1 > 1$ it follows that $i(U_{i_1} \dots U_{i_p}) \geq 2$. Otherwise $i_1 = 1$ and, since $r < p$, there is some $j \geq 2$ such that $i_j \geq i_{j-1} + 2$, so that $U_{i_1} \dots U_{i_r}$ can be written as a word with terminus i_{j-1} , and then the claim follows. Coefficients are given by powers of λ and μ , and products of these. The terms λ and μ are defined via the homomorphisms in Definition 2.20 and lie in the set $\{-1, \pm v, v^2\}$. Since v is a unit it follows that all coefficients are units. \square

Lemma 8.24. *Let $k \geq 0$ and $-1 \leq i \leq n-1$, and consider elements $x_{\underline{a}, \underline{b}} \otimes 1$, where $x_{\underline{a}, \underline{b}}$ is in Jones normal form and satisfies either:*

- $i(x_{\underline{a}, \underline{b}}) \geq 2$ and $t(x_{\underline{a}, \underline{b}}) \geq n - i - 1$, or
- $i(x_{\underline{a}, \underline{b}}) = 1$ and $n - i - 1 \leq t(x_{\underline{a}, \underline{b}}) \leq n - i - 1 + k$.

Then these elements all lie in F_i^k .

Proof. The first type of element lies in F^0 , since in this case $x_{\underline{a}, \underline{b}}$ is a word in the U_i s not containing U_1 , and by Lemma 8.22, this is a linear combination of words in the s_i s not containing s_1 (a basic element). For the second type of element, from the definition of Jones normal form, $x_{\underline{a}, \underline{b}}$ must end in a string $U_1 \dots U_{n-i-1+j}$ for $0 \leq j \leq k$. We proceed by induction on k .

Base case: We start with the base case $k = 0$, so the only option is that $j = 0$. i.e. $x_{\underline{a}, \underline{b}} = y_{\underline{a}, \underline{b}} U_1 \dots U_{n-i-1}$ for some $y_{\underline{a}, \underline{b}}$ in Jones normal form with $i(y_{\underline{a}, \underline{b}}) \geq 2$. We aim to show in this case $x_{\underline{a}, \underline{b}}$ lies in F_i^0 . Compare $x_{\underline{a}, \underline{b}}$ to $y_{\underline{a}, \underline{b}} s_1 \dots s_{n-i-1}$, which does lie in F_i^0 by Definition 8.1. From Lemma 8.23 multiplying out the string $s_1 \dots s_{n-i-1}$ will result in $y_{\underline{a}, \underline{b}} s_1 \dots s_{n-i-1}$ being written as a linear combination (up to scalar multiplication by units) of three types of elements, and we consider their image in $\text{TL}_n \otimes_{\text{TL}_{n-i-1}} \mathbb{1}$.

- (1) $y_{\underline{a}, \underline{b}} U_1 \dots U_{n-i-1}$. This is equal to $x_{\underline{a}, \underline{b}}$ and appears as a single summand of $y_{\underline{a}, \underline{b}} s_1 \dots s_{n-i-1}$.
- (2) $y_{\underline{a}, \underline{b}} w$ where $i(w) \geq 2$ and $t(w) \leq n - i - 1$. These are all basic elements since $i(y_{\underline{a}, \underline{b}}) \geq 2$, and thus lie in F^0 .
- (3) $y_{\underline{a}, \underline{b}} w$ where $i(w) = 1$ and $t(w) < n - i - 1$. Due to the terminus, these are all zero in $\text{TL}_n \otimes_{\text{TL}_{n-i-1}} \mathbb{1}$.

So it follows that in $\text{TL}_n \otimes_{\text{TL}_{n-i-1}} \mathbb{1}$, $x_{\underline{a}, \underline{b}}$ is, up to scalar multiplication by units, equal to a linear combination of $y_{\underline{a}, \underline{b}} s_1 \dots s_{n-i-1}$ and basic elements. Since this is a linear combination of elements in F_i^0 , it follows that $x_{\underline{a}, \underline{b}}$ lies in F_i^0 as required.

Inductive step: We now assume the Lemma is true for $k-1$ and prove for k . Let $x_{\underline{a}, \underline{b}} = y_{\underline{a}, \underline{b}} U_1 \dots U_{n-i-1+j}$ for some $y_{\underline{a}, \underline{b}}$ in Jones normal form with $i(y_{\underline{a}, \underline{b}}) \geq 2$, $t(y_{\underline{a}, \underline{b}}) > n - i - 1 + j$, and $0 \leq j \leq k$. When $0 \leq j < k$, by the inductive hypothesis this element lies in $F_i^{k-1} \subset F_i^k$, and so we can restrict to the case where $j = k$, i.e. $x_{\underline{a}, \underline{b}} = y_{\underline{a}, \underline{b}} U_1 \dots U_{n-i-1+k}$. We aim to show that, in this case, $x_{\underline{a}, \underline{b}}$ lies in F_i^k . As in the base case, we compare $x_{\underline{a}, \underline{b}}$ with $y_{\underline{a}, \underline{b}} s_1 \dots s_{n-i-1+k}$, which lies in F_i^k by Definition 8.1. From Lemma 8.23, $y_{\underline{a}, \underline{b}} s_1 \dots s_{n-i-1+k}$ is a linear combination (up

to scalar multiplication by a unit) of three types of elements, which we evaluate in $\mathrm{TL}_n \otimes_{\mathrm{TL}_{n-i-1}} \mathbb{1}$.

- (1) $y_{\underline{a},\underline{b}}U_1 \dots U_{n-i-1+k}$. This is equal to $x_{\underline{a},\underline{b}}$ and appears as a single summand of $y_{\underline{a},\underline{b}}s_1 \dots s_{n-i-1+k}$.
- (2) $y_{\underline{a},\underline{b}}w$ where $i(w) \geq 2$ and $t(w) \leq n - i - 1 + k$. Since $i(y_{\underline{a},\underline{b}}) \geq 2$ they are all basic elements, and thus lie in $F^0 \subset F^k$.
- (3) $y_{\underline{a},\underline{b}}w$ where $i(w) = 1$ and $t(w) < n - i - 1 + k$. Rewriting these in Jones normal form gives elements $y_{\underline{a},\underline{b}}w = z_{\underline{a},\underline{b}}$ such that $i(z_{\underline{a},\underline{b}}) = 1$ and $t(z_{\underline{a},\underline{b}}) \leq t(w) < n - i - 1 + k$ (by Lemmas 2.6 and 2.7). These are then Jones normal form elements ending in $U_1 \dots U_{n-i-1+j}$ for $0 \leq j \leq k - 1$ so by the inductive hypothesis these lie in $F_i^{k-1} \subset F_i^k$.

Again it follows that in $\mathrm{TL}_n \otimes_{\mathrm{TL}_{n-i-1}} \mathbb{1}$, $x_{\underline{a},\underline{b}}$ is, up to scalar multiplication by units, equal to a linear combination of $y_{\underline{a},\underline{b}}s_1 \dots s_{n-i-k}$, elements in F_i^k (by the inductive hypothesis), and basic elements. Since this is a linear combination of elements in F_i^k , it follows that $x_{\underline{a},\underline{b}}$ lies in F_i^k as required. \square

Lemma 8.25. *Let $k \geq 0$ and $-1 \leq i \leq n - 1$. Then F_i^k has basis consisting of elements $x_{\underline{a},\underline{b}} \otimes 1$, where $x_{\underline{a},\underline{b}}$ is in Jones normal form and satisfies either:*

- $i(x_{\underline{a},\underline{b}}) \geq 2$ and $t(x_{\underline{a},\underline{b}}) \geq n - i - 1$, or
- $i(x_{\underline{a},\underline{b}}) = 1$ and $n - i - 1 \leq t(x_{\underline{a},\underline{b}}) \leq n - i - 1 + k$.

Proof. This is a subset of the known basis for $\mathrm{TL}_n \otimes_{\mathrm{TL}_{n-i-1}} \mathbb{1} \supseteq F_i^k$, and by the previous lemma we know these elements lie in F_i^k , so it is enough to show that F_i^k is spanned by these elements. First of all, note that since $F_i^k \subseteq \mathrm{TL}_n \otimes_{\mathrm{TL}_{n-i-1}} \mathbb{1}$ any word in F_i^k written in Jones normal form will vanish if $t(x_{\underline{a},\underline{b}}) \leq n - i - 2$, therefore we will always have $t(x_{\underline{a},\underline{b}}) \geq n - i - 1$. By definition F_i^k is spanned by elements of the form

- $x \otimes 1$
- $x \cdot (s_1 \dots s_{n-i-1+k'}) \otimes 1$

where x is a word in the U_i with $i(x) \geq 2$ (i.e. containing no U_1 s) and $0 \leq k' \leq k$ (note in the case $i = n - 1$ and $k' = 0$ the two kinds coincide). The first kind is spanned by $x_{\underline{a},\underline{b}}$ such that $i(x_{\underline{a},\underline{b}}) \geq 2$, as described in the first bullet point in the statement of the lemma. From Lemma 8.23, expanding the product $(s_1 \dots s_{n-i-1+k'})$ in the second kind gives a linear combination of words $x \cdot w \otimes 1$ such that $t(w) \leq n - i - 1 + k'$. Either $i(w)$ will be ≥ 2 or $i(w) = 1$. In the first case, since $i(x) \geq 2$ it follows that $i(x \cdot w) \geq 2$ and so when written in Jones normal form this will remain the case, giving an element of the first type described in the Lemma. In the second case, when $i(w) = 1$, since $i(x) \geq 2$ then either $i(x \cdot w) \geq 2$ and as in the previous sentence we are done, or $i(x \cdot w) = 1$ and, by Lemma 2.7, when written in Jones normal form the terminus $t(x \cdot w) = t(w) \leq n - i - 1 + k' \leq n - i - 1 + k$ will either remain the same or reduce. This puts us in the setting of

the second bullet point in the statement of the lemma, and thus we have shown that the two types of elements span F_i^k . \square

Proposition 8.26. *The map $\Phi^0: C(W(n-1)) \longrightarrow F^0$ from Definition 8.11 is an isomorphism.*

Proof. Recall that for $-1 \leq i \leq (n-1)$,

$$\Phi_i^0: (\mathrm{TL}_{n-1} \otimes_{\mathrm{TL}_{n-i-2}} \mathbb{1}) \oplus (\mathrm{TL}_{n-1} \otimes_{\mathrm{TL}_{n-i-1}} \mathbb{1}) \rightarrow F_i^0$$

is given by

$$\Phi_i^0(x \otimes \alpha, y \otimes \beta) = \xi_i(x \otimes \alpha) + \eta_i(y \otimes \beta)$$

where

$$\xi_i(x \otimes \alpha) = \sigma(x) \otimes \lambda^{n-1} \alpha$$

and

$$\eta_i(y \otimes \beta) = \sigma(y) \cdot (s_1 \cdots s_{n-i-1}) \otimes \lambda^i \beta.$$

By Lemma 2.16, a basis for the left hand side is given by elements of either the form $(x_{\underline{a}, \underline{b}} \otimes 1, 0)$ such that $t(x_{\underline{a}, \underline{b}}) > n - i - 3$ or the form $(0, x_{\underline{a}', \underline{b}'} \otimes 1)$ such that $t(x_{\underline{a}', \underline{b}'}) > n - i - 2$. Under the map Φ_i^0 , $(x_{\underline{a}, \underline{b}} \otimes 1, 0)$ is taken to a scalar multiple (by a unit) of $\sigma(x_{\underline{a}, \underline{b}}) \otimes 1$, where $\sigma(x_{\underline{a}, \underline{b}})$ is a Jones basis element with $i(\sigma(x_{\underline{a}, \underline{b}})) \geq 2$ and $t(\sigma(x_{\underline{a}, \underline{b}})) > n - i - 2$. By Lemma 8.23, the element $(0, x_{\underline{a}', \underline{b}'}) \otimes 1$ is taken to a linear combination of scalar multiples (by units) of terms $\sigma(x_{\underline{a}', \underline{b}'}) \cdot w \otimes 1$ such that $t(w) \leq n - i - 1$. Since $F_i^0 \subseteq \mathrm{TL}_n \otimes_{\mathrm{TL}_{n-i-1}} \mathbb{1}$ the only non-zero terms in the image will occur when $t(w) = n - i - 1$. We consider two cases: $i(w) \geq 2$ or $i(w) = 1$. By Lemma 2.7, converting to Jones normal form in the first case gives an element with index $i(\sigma(x_{\underline{a}', \underline{b}'}) \cdot w) > 2$ and terminus $t(\sigma(x_{\underline{a}', \underline{b}'}) \cdot w) = n - i - 1$, or zero, since the terminus will either remain the same or reduce when converting. When $i(w) = 1$ and $t(w) = n - i - 1$, by Lemma 8.23 it follows that $w = U_1 \dots U_{n-i-1}$ and therefore the terms will be of the form $\sigma(x_{\underline{a}', \underline{b}'}) \cdot U_1 \dots U_{n-i-1}$. These elements are already in Jones normal form, with index 1 and terminus $n - i - 1$. Furthermore all Jones basis elements with this index and terminus arise in this way. By Lemma 8.25 a basis for F_i^0 is given by elements $y_{\underline{a}, \underline{b}} \otimes 1$ where $y_{\underline{a}, \underline{b}}$ is in Jones normal form and satisfies:

- $i(y_{\underline{a}, \underline{b}}) \geq 2$ and $t(y_{\underline{a}, \underline{b}}) \geq n - i - 1$ or
- $i(y_{\underline{a}, \underline{b}}) = 1$ and $t(y_{\underline{a}, \underline{b}}) = n - i - 1$.

By our analysis, all of these elements lie in the image of Φ_i^0 , up to scalar multiplication by units, hence Φ^0 is a bijection on bases and therefore an isomorphism. \square

Lemma 8.27. *A basis for $(F^k/F^{k-1})_i$ is given by words $x_{\underline{a}, \underline{b}}$ in Jones normal form such that $i(x_{\underline{a}, \underline{b}}) = 1$ and $t(x_{\underline{a}, \underline{b}}) = n - i - 1 + k$.*

Proof. This is a direct consequence of taking the quotient of the bases for F^k and F^{k-1} given in Lemma 8.25. \square

Proposition 8.28. *The map $\Psi^k : \tau_{n-1}\Sigma^{k+1}W(n-1) \longrightarrow F^k/F^{k-1}$ from Definition 8.18 is an isomorphism.*

Proof. Recall for i in the range $k \leq i \leq (n-1)$,

$$\Psi_i^k : \mathrm{TL}_{n-1} \otimes_{\mathrm{TL}_{n-i-1+k}} \mathbb{1} \longrightarrow (F^k/F^{k-1})_i$$

is given by

$$\Psi_i^k : x \otimes \alpha \longmapsto (-1)^{-i(k+1)} \sigma(x) \cdot (s_1 \cdots s_{n-i-1+k}) \otimes \lambda^i \alpha.$$

By Lemma 2.16 a basis for the domain is given by $x_{\underline{a}, \underline{b}}$ such that $t(x_{\underline{a}, \underline{b}}) > (n-i-1+k) - 1$. Note also that $x_{\underline{a}, \underline{b}}$ does not contain the letter U_{n-1} . By Lemma 8.23, the image $\Psi_i^k(x_{\underline{a}, \underline{b}})$ is a linear combination of scalar multiples (by units) of terms $\sigma(x_{\underline{a}, \underline{b}}) \cdot w$ such that $t(w) \leq n-i-1+k$. These terms are zero in $(F^k/F^{k-1})_i \subseteq \mathrm{TL}_n \otimes_{\mathrm{TL}_{n-i-1}} \mathbb{1}$ only when w cannot be written as a word with $t(w) < n-i-1$. Rewriting these elements in Jones normal form will maintain or decrease the terminus, and $i(\sigma(x_{\underline{a}, \underline{b}})) \geq 2$, so $i(\sigma(x_{\underline{a}, \underline{b}}) \cdot w) = 1$ only when $i(w) = 1$. Therefore by Lemma 8.25 quotienting out by F^{k-1} leaves only the term for which $i(w) = 1$ and $t(w) = n-i-1+k$. In particular by Lemma 8.23 this term is a scalar multiple (by a unit) of $\sigma(x_{\underline{a}, \underline{b}}) \cdot U_1 \cdots U_{n-i-1+k}$.

Since $\sigma(x_{\underline{a}, \underline{b}})$ has index ≥ 2 and terminus $> n-i-1+k$, it follows that $\sigma(x_{\underline{a}, \underline{b}}) \cdot U_1 \cdots U_{n-i-1+k}$ is in Jones normal form. From Lemma 8.27 this is a Jones basis element for F^k/F^{k-1} and all basis elements arise in this way. Therefore up to unit scalars, the map Ψ^k is a bijection on bases, and hence an isomorphism. \square

9. JONES-WENZL PROJECTORS AND VANISHING

This section relates our results with the existence of the *Jones-Wenzl projectors*, to strengthen our vanishing results when R is a field. This section is written such that the reader can read the introduction, the background on Temperley-Lieb algebras, and continue straight to this section. For the time being we make the substitutions $a \leftrightarrow \delta$ and $v \leftrightarrow q$, as is common in the recent literature concerning Jones-Wenzl projectors.

Throughout this section, we will consider a commutative ring R , a unit $q \in R$, the parameter $\delta = q + q^{-1}$, and we will work in $\mathrm{TL}_n(\delta)$. Recall that we show in Theorem A that, when δ is invertible, $\mathrm{Tor}_*^{\mathrm{TL}_n(\delta)}(\mathbb{1}, \mathbb{1})$ and $\mathrm{Ext}_{\mathrm{TL}_n(\delta)}^*(\mathbb{1}, \mathbb{1})$ vanish in every non-zero degree. In this section we investigate the case where $\delta = 0$ and R is a field using established results on Jones-Wenzl projectors. We prove the following theorem:

Theorem D. *Let $n = 2k + 1$, and let R be a field whose characteristic does not divide $\binom{k}{i}$ for any $0 \leq i \leq k$. Let q be a unit in R and assume that $\delta = q + q^{-1} = 0$. Then $\mathrm{Tor}_*^{\mathrm{TL}_n(0)}(\mathbb{1}, \mathbb{1})$ and $\mathrm{Ext}_{\mathrm{TL}_n(0)}^*(\mathbb{1}, \mathbb{1})$ vanish in positive degrees.*

For example, when $n = 3$, R is a field, and $\delta = q + q^{-1}$ for $q \in R^\times$, combining this theorem with Theorem A demonstrates that $\mathrm{Tor}_*^{\mathrm{TL}_3(\delta)}(\mathbb{1}, \mathbb{1})$ and $\mathrm{Ext}_{\mathrm{TL}_3(\delta)}^*(\mathbb{1}, \mathbb{1})$

vanish in positive degrees with no further condition on δ . And if one wishes to show that $\mathrm{Tor}_*^{\mathrm{TL}_5(\delta)}(\mathbb{1}, \mathbb{1})$ and $\mathrm{Ext}_{\mathrm{TL}_5(\delta)}^*(\mathbb{1}, \mathbb{1})$ can be nonzero in positive degrees, then the only chance of this happening is in characteristic 2.

The theorem is in strict contrast to the n even case, where we show in Theorem A that for a general ring R and δ not invertible, $\mathrm{Tor}_{n-1}^{\mathrm{TL}_n(\delta)}(\mathbb{1}, \mathbb{1}) = R/bR$ is non-zero for b some multiple of δ . Therefore, in the particular case where n is even, R is a field, and $\delta = 0$, there can be no vanishing in all positive degrees.

9.1. Jones-Wenzl projectors. In this subsection we introduce the Jones-Wenzl projector and relate its existence to the projectivity of the trivial module $\mathbb{1}$. The original references are [Jon83] and [Wen87], see also [KL94], [Lic92, section 4].

Definition 9.1. Recall that $I_n \subseteq \mathrm{TL}_n$ is the two sided ideal generated by the U_i for $i = 1, \dots, n-1$. Then, if it exists, the n th Jones-Wenzl projector JW_n is the element of TL_n characterised by the following two properties:

- (i) $\mathrm{JW}_n \in 1 + I_n$ and
- (ii) $I_n \cdot \mathrm{JW}_n = 0 = \mathrm{JW}_n \cdot I_n$.

Lemma 9.2. *If JW_n exists it is unique.*

Proof. Suppose a second element JW'_n in TL_n satisfies (i) and (ii) of Definition 9.1. Write $\mathrm{JW}_n = 1 + i$ and $\mathrm{JW}'_n = 1 + i'$ for $i, i' \in I_n$. Then $\mathrm{JW}_n \cdot i' = 0 = i \cdot \mathrm{JW}'_n$ by (ii). It follows that

$$\mathrm{JW}'_n = \mathrm{JW}'_n + i \cdot \mathrm{JW}'_n = (1 + i) \cdot \mathrm{JW}'_n = \mathrm{JW}_n \cdot \mathrm{JW}'_n$$

and similarly that $\mathrm{JW}_n = \mathrm{JW}_n \cdot \mathrm{JW}'_n$. \square

The Jones-Wenzl projector was first introduced by Jones [Jon83], was further studied by Wenzl [Wen87], and has since become important in representation theory, knot theory and the study of 3-manifolds. It is a key ingredient in the definition of the coloured Jones polynomial and $\mathrm{SU}(2)$ quantum invariants more generally, and is important in the study of tilting modules of (quantum) \mathfrak{sl}_2 .

9.2. JW_n and projectivity of $\mathbb{1}$. We will now show that the Jones-Wenzl projector exists if and only if the trivial module $\mathbb{1}$ is projective. Thus, existence of JW_n implies the vanishing of $\mathrm{Tor}_*^{\mathrm{TL}_n(\delta)}(\mathbb{1}, \mathbb{1})$ and $\mathrm{Ext}_{\mathrm{TL}_n(\delta)}^*(\mathbb{1}, \mathbb{1})$ in positive degrees. Our own Theorem A implies that vanishing for δ invertible, while Theorem C proves non-vanishing for n even and δ not invertible. It turns out that there is a rich interplay between these two sources of (non-)vanishing results.

Proposition 9.3. *JW_n exists if and only if $\mathbb{1}$ is a projective left $\mathrm{TL}_n(\delta)$ -module, which is if and only if $\mathbb{1}$ is a projective right $\mathrm{TL}_n(\delta)$ -module.*

Before proving the proposition we need the following.

Lemma 9.4. *In Definition 9.1, it is sufficient to replace (ii) with either*

- (ii)' $I_n \cdot \mathrm{JW}_n = 0$, or

(ii)'' $JW_n \cdot I_n = 0$.

Proof. Suppose $JW \in TL_n$ satisfies (i) and (ii)'. We have suggestively named this element, and will show it is in fact JW_n , by showing that JW also satisfies (ii)'' and hence (ii). Let $TL_n \rightarrow TL_n, d \mapsto \bar{d}$ be the anti-automorphism which reverses the order of letters in a monomial i.e. $\overline{U_{i_1} \dots U_{i_n}} = U_{i_n} \dots U_{i_1}$. In diagrammatic terms, this map flips the diagram corresponding to the monomial in the left-to-right direction. Since JW satisfies (ii)', it follows that \overline{JW} satisfies (ii)'' . Then the argument of Lemma 9.2 can be repeated to show that $JW = \overline{JW}$ so that JW satisfies (ii)' and (ii)'', hence it satisfies (ii) and $JW = JW_n$. \square

Proof of Proposition 9.3. We prove the equivalence for left-modules.

If JW_n exists, then the maps $\mathbb{1} \rightarrow TL_n, 1 \mapsto JW_n$ and $TL_n \rightarrow \mathbb{1}, d \mapsto d \cdot 1$ are maps of left TL_n -modules composing to the identity. It follows that $\mathbb{1}$ is a direct summand of TL_n , and thus is projective.

Conversely, if $\mathbb{1}$ is a projective left TL_n -module, then the surjection $TL_n \rightarrow TL_n/I_n = \mathbb{1}$, regarded as a map of left TL_n -modules, has a splitting $s: \mathbb{1} \rightarrow TL_n$, again a map of left TL_n -modules. By construction the element $s(1)$ then satisfies condition (i) of 9.1 and condition (ii)' of 9.4, so that $JW_n = s(1)$ exists as required. \square

9.3. Jones-Wenzl projectors and quantum binomial coefficients. Here we work in the Laurent polynomial ring $\mathbb{Z}[q, q^{-1}]$, and we set $\delta = q + q^{-1}$. For this section, let n and r be integers such that $n \geq r \geq 0$.

Definition 9.5. The *quantum integer* $[n]_q$ is defined to be

$$[n]_q = \frac{q^n - q^{-n}}{q - q^{-1}} = q^{n-1} + q^{n-3} + \dots + q^{-(n-3)} + q^{-(n-1)},$$

the *quantum factorial* $[n]_q!$ is defined by

$$[n]_q! = [n]_q [n-1]_q \cdots [1]_q,$$

and the *quantum binomial coefficient* $\begin{bmatrix} n \\ r \end{bmatrix}_q$ is then given by computing the normal binomial coefficient but replacing integers with quantum integers:

$$\begin{bmatrix} n \\ r \end{bmatrix}_q = \frac{[n]_q!}{[r]_q! [n-r]_q!}.$$

The quantum binomial coefficients satisfy the following recursion relations:

$$\begin{aligned} \begin{bmatrix} n \\ r \end{bmatrix}_q &= q^{n-r} \begin{bmatrix} n-1 \\ r-1 \end{bmatrix}_q + q^{-r} \begin{bmatrix} n-1 \\ r \end{bmatrix}_q \\ \begin{bmatrix} n \\ r \end{bmatrix}_q &= q^{-(n-r)} \begin{bmatrix} n-1 \\ r-1 \end{bmatrix}_q + q^r \begin{bmatrix} n-1 \\ r \end{bmatrix}_q. \end{aligned}$$

Either one of these relations gives an inductive proof that $\begin{bmatrix} n \\ r \end{bmatrix}_q$ lies in $\mathbb{Z}[q, q^{-1}]$. And taken together, these relations give an inductive proof that $\begin{bmatrix} n \\ r \end{bmatrix}_q$ is invariant under inverting q , and consequently that it lies in $\mathbb{Z}[\delta]$. (Recall that $\delta = q + q^{-1}$.) This means that we may evaluate $\begin{bmatrix} n \\ r \end{bmatrix}_q$ in any ring containing an element named δ , to obtain an element of that ring, which we continue to denote by $\begin{bmatrix} n \\ r \end{bmatrix}_q$.

The following result is proved in an Appendix to [EL17] by Webster, using Schur-Weyl duality. For a purely diagrammatic approach see recent work of Spencer [Spe21, Section 10.3].

Theorem 9.6 ([EL17, Theorem A.2], [Spe21, Section 10.3]). *Let $R = \mathbb{k}$ be a field, let $q \in \mathbb{k}$ be nonzero, and set $\delta = q + q^{-1}$. The n th Jones-Wenzl projector $\text{JW}_n \in \text{TL}_n(\delta)$ exists if and only if the quantum binomial coefficients $\begin{bmatrix} n \\ r \end{bmatrix}_q$ are non-zero in \mathbb{k} for all $0 \leq r \leq n$.*

Remark 9.7. Suppose that $R = \mathbb{k}$ is a field. Whenever \mathbb{k} , q and n satisfy the conditions of Theorem 9.6, we obtain the vanishing of $\text{Tor}_*^{\text{TL}_n(\delta)}(\mathbb{1}, \mathbb{1})$ and $\text{Ext}_{\text{TL}_n(\delta)}^*(\mathbb{1}, \mathbb{1})$ in positive degrees. (We will refer to this as simply “vanishing” for the present remark.)

- In the case of n even, Theorems A and C show that vanishing holds if and only if $\delta \neq 0$, and this is in fact stronger than the result obtained from Theorem 9.6. For example, if we take $n = 4$ then the $\begin{bmatrix} n \\ k \end{bmatrix}_q$ take values 1, $\delta(\delta^2 - 2)$ and $(\delta^2 - 1)(\delta^2 - 2)$, so that Theorem 9.6 requires δ to avoid the values $0, \pm 1, \pm\sqrt{2}$. For n even, δ is always a factor of $\begin{bmatrix} n \\ 1 \end{bmatrix}_q$, so that Theorem A will always apply more generally than Theorem 9.6 in this case.
- In the case of n odd, the situation is more interesting. Theorem A demonstrates vanishing when $\delta \neq 0$. But if we take $n = 3$, for example, then the $\begin{bmatrix} n \\ k \end{bmatrix}_q$ take values 1 and $(\delta^2 - 1)$, so that Theorem 9.6 demonstrates vanishing so long as $\delta \neq \pm 1$. Neither of these vanishing results implies the other, but taken together they demonstrate vanishing for *all* values of δ .

9.4. Identifying the quantum binomial coefficients. In this section, we identify the quantum binomial coefficients upon specialising $\delta = q + q^{-1} = 0$. The results are assembled in the following proposition.

Proposition 9.8. *When $\delta = q + q^{-1} = 0$ then the quantum binomial coefficients have the following form:*

- When n is even and r is odd, $\begin{bmatrix} n \\ r \end{bmatrix}_q = 0$.
- When n and r are both even, let $n = 2a$ and $r = 2t$. Then $\begin{bmatrix} n \\ r \end{bmatrix}_q = \binom{a}{t}$.
- When n is odd and r is even, let $n = 2a + 1$ and $r = 2t$. Then $\begin{bmatrix} n \\ r \end{bmatrix}_q = (-1)^t \binom{a}{t}$.

2. Then the cyclotomic polynomial $\Phi_2(p) = 1 + p = 1 + q^2 = 0$. Let $n = 2a + b$ and $r = 2t + s$ with $0 \leq b, s \leq 1$, then the quantum binomial coefficient satisfies

$$\begin{bmatrix} n \\ r \end{bmatrix}_q = q^{r^2 - nr} \begin{bmatrix} n \\ r \end{bmatrix}_{q^2}^G = q^{r^2 - nr} \binom{a}{t} \begin{bmatrix} b \\ s \end{bmatrix}_{q^2}^G.$$

When n is even and r is odd, $\begin{bmatrix} b \\ s \end{bmatrix}_{q^2}^G = \begin{bmatrix} 0 \\ 1 \end{bmatrix}_{q^2}^G = 0$ which gives the first case of the proposition. For all other cases, $\begin{bmatrix} b \\ s \end{bmatrix}_{q^2}^G = 1$ and so

$$\begin{bmatrix} n \\ r \end{bmatrix}_q = q^{r^2 - nr} \binom{a}{t}.$$

Computing the coefficient $q^{r^2 - nr}$, using $q^{\pm 2} = -1$, yields the result for the remaining three cases. \square

9.5. Proof of theorem. In this section we prove Theorem D.

Proof. This proof puts together three previous results. By Proposition 9.3 we know that if JW_n exists then $\mathbb{1}$ is projective and it follows that $\text{Tor}_i^{\text{TL}(0)}(\mathbb{1}, \mathbb{1})$ and $\text{Ext}_{\text{TL}_n(0)}^*(\mathbb{1}, \mathbb{1})$ vanish for all $i > 0$. So it is enough to show that JW_n exists under the hypotheses of the theorem. Proposition 9.6 tells us that JW_n exists precisely when the quantum binomial coefficients $\begin{bmatrix} n \\ r \end{bmatrix}_q$ are non-zero for all $1 \leq r \leq n$. Finally, Proposition 9.8 explicitly describes these coefficients when $\delta = 0$. We see that for n even, there is always a quantum binomial coefficient $\begin{bmatrix} n \\ r \end{bmatrix}_q = 0$ and so we learn nothing new. However when $n = 2k + 1$ is odd, the quantum binomial coefficients take values in the set

$$\left\{ \pm \binom{k}{t} \mid 0 \leq t \leq k \right\}$$

and—up to sign—all values in this set are realised as some $\begin{bmatrix} n \\ r \end{bmatrix}_q$. The hypotheses of the theorem precisely say that these numbers are non-zero in R and so the result follows. \square

REFERENCES

- [BCH20] Chris Bowman, Anton Cox, and Amit Hazi. Path isomorphisms between quiver Hecke and diagrammatic Bott-Samelson endomorphism algebras. [arXiv:2005.02825](#), 2020.
- [BH21] Rachael Boyd and Richard Hepworth. Combinatorics of injective words for Temperley-Lieb algebras. *Journal of Combinatorial Theory, Series A*, 181:105446, 2021.
- [BHP21] Rachael Boyd, Richard Hepworth, and Peter Patzt. The homology of the Brauer algebras. *Selecta Math. (N.S.)*, 27(5):Paper No. 85, 31, 2021.
- [BK09] Jonathan Brundan and Alexander Kleshchev. Blocks of cyclotomic Hecke algebras and Khovanov-Lauda algebras. *Invent. Math.*, 178(3):451–484, 2009.
- [Boyd20] Rachael Boyd. Homological stability for Artin monoids. *Proceedings of the London Mathematical Society*, 121(3):537–583, 2020.

- [BW83] Anders Björner and Michelle Wachs. On lexicographically shellable posets. *Trans. Amer. Math. Soc.*, 277(1):323–341, 1983.
- [Cha80] Ruth M. Charney. Homology stability for GL_n of a Dedekind domain. *Invent. Math.*, 56(1):1–17, 1980.
- [Chu12] Thomas Church. Homological stability for configuration spaces of manifolds. *Invent. Math.*, 188(2):465–504, 2012.
- [Dés83] Jacques Désarménien. Une autre interprétation du nombre de dérangements. *Sém. Lothar. Combin.*, 8(B08b), 1983.
- [DS01] Emeric Deutsch and Louis Shapiro. A survey of the Fine numbers. *Discrete Math.*, 241(1-3):241–265, 2001. Selected papers in honor of Helge Tverberg.
- [EL17] Ben Elias and Nicolas Libedinsky. Indecomposable Soergel bimodules for universal Coxeter groups. *Trans. Amer. Math. Soc.*, 369(6):3883–3910, 2017. With an appendix by Ben Webster.
- [Far79] Frank D. Farmer. Cellular homology for posets. *Math. Japon.*, 23(6):607–613, 1978/79.
- [FG97] C. K. Fan and Richard M. Green. Monomials and Temperley-Lieb algebras. *J. Algebra*, 190(2):498–517, 1997.
- [GKRW18] Søren Galatius, Alexander Kupers, and Oscar Randal-Williams. Cellular E_k -algebras. [arXiv:1805.07184](https://arxiv.org/abs/1805.07184), 2018.
- [GL96] J. J. Graham and G. I. Lehrer. Cellular algebras. *Invent. Math.*, 123(1):1–34, 1996.
- [GRW18] Søren Galatius and Oscar Randal-Williams. Homological stability for moduli spaces of high dimensional manifolds. I. *J. Amer. Math. Soc.*, 31(1):215–264, 2018.
- [Har85] John L. Harer. Stability of the homology of the mapping class groups of orientable surfaces. *Ann. of Math. (2)*, 121(2):215–249, 1985.
- [Hep16] Richard Hepworth. Homological stability for families of Coxeter groups. *Algebr. Geom. Topol.*, 16(5):2779–2811, 2016.
- [Hep20] Richard Hepworth. Homological stability for Iwahori-Hecke algebras. [arXiv:2006.04252](https://arxiv.org/abs/2006.04252), 2020.
- [HMR09] Tom Halverson, Manuela Mazzocco, and Arun Ram. Commuting families in Hecke and Temperley-Lieb algebras. *Nagoya Math. J.*, 195:125–152, 2009.
- [HV98] Allen Hatcher and Karen Vogtmann. Rational homology of $\text{Aut}(F_n)$. *Math. Res. Lett.*, 5(6):759–780, 1998.
- [HV04] Allen Hatcher and Karen Vogtmann. Homology stability for outer automorphism groups of free groups. *Algebr. Geom. Topol.*, 4:1253–1272, 2004.
- [HW10] Allen Hatcher and Nathalie Wahl. Stabilization for mapping class groups of 3-manifolds. *Duke Math. J.*, 155(2):205–269, 2010.
- [Jon83] Vaughan F. R. Jones. Index for subfactors. *Invent. Math.*, 72(1):1–25, 1983.
- [Jon85] Vaughan F. R. Jones. A polynomial invariant for knots via von Neumann algebras. *Bull. Amer. Math. Soc. (N.S.)*, 12(1):103–111, 1985.
- [Kau87] Louis H. Kauffman. State models and the Jones polynomial. *Topology*, 26(3):395–407, 1987.
- [Kau90] Louis H. Kauffman. An invariant of regular isotopy. *Trans. Amer. Math. Soc.*, 318(2):417–471, 1990.
- [Kau05] Louis H. Kauffman. Knot diagrammatics. In *Handbook of knot theory*, pages 233–318. Elsevier B. V., Amsterdam, 2005.
- [Ker05] Moritz C. Kerz. The complex of words and Nakaoka stability. *Homology Homotopy Appl.*, 7(1):77–85, 2005.

- [KL94] Louis H. Kauffman and Sostenes L. Lins. *Temperley-Lieb recoupling theory and invariants of 3-manifolds*, volume 134 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 1994.
- [KT08] Christian Kassel and Vladimir Turaev. *Braid groups*, volume 247 of *Graduate Texts in Mathematics*. Springer, New York, 2008. With the graphical assistance of Olivier Dodane.
- [Lic92] W. B. R. Lickorish. Calculations with the Temperley-Lieb algebra. *Comment. Math. Helv.*, 67(4):571–591, 1992.
- [Maa79] Hendrik Maazen. *Homology stability for the general linear group*. Utrecht University, 1979. Thesis (Ph.D.).
- [Nak60] Minoru Nakaoka. Decomposition theorem for homology groups of symmetric groups. *Ann. of Math. (2)*, 71:16–42, 1960.
- [OEIS] Neil J. A. Sloane. The On-Line Encyclopaedia of Integer Sequences. <https://oeis.org/A001045>.
- [Qui72] Daniel Quillen. On the cohomology and K -theory of the general linear groups over a finite field. *Ann. of Math. (2)*, 96:552–586, 1972.
- [RSA14] David Ridout and Yvan Saint-Aubin. Standard modules, induction and the structure of the Temperley-Lieb algebra. *Adv. Theor. Math. Phys.*, 18(5):957–1041, 2014.
- [RW13] Oscar Randal-Williams. Homological stability for unordered configuration spaces. *Q. J. Math.*, 64(1):303–326, 2013.
- [RW16] Oscar Randal-Williams. Resolutions of moduli spaces and homological stability. *J. Eur. Math. Soc. (JEMS)*, 18(1):1–81, 2016.
- [RW21] Oscar Randal-Williams. A remark on the homology of Temperley-Lieb algebras, 2021. <https://www.dpmms.cam.ac.uk/~or257/notes/Descent.pdf>.
- [RWW17] Oscar Randal-Williams and Nathalie Wahl. Homological stability for automorphism groups. *Adv. Math.*, 318:534–626, 2017.
- [Spe21] Robert Spencer. The modular Temperley-Lieb algebra. [arXiv:2011.01328](https://arxiv.org/abs/2011.01328), 2021.
- [Spe22] Robert Spencer. Jacobsthal Elements. <https://robertandrewspencer.com/research/jacobsthal/>.
- [Sro22] Robin Sroka. The homology of a Temperley-Lieb algebra on an odd number of strands. [arXiv:2202.08799](https://arxiv.org/abs/2202.08799), 2022.
- [SW19] Markus Szymik and Nathalie Wahl. The homology of the Higman-Thompson groups. *Invent. Math.*, 216(2):445–518, 2019.
- [TL71] H. N. V. Temperley and E. H. Lieb. Relations between the “percolation” and “colouring” problem and other graph-theoretical problems associated with regular planar lattices: some exact results for the “percolation” problem. *Proc. Roy. Soc. London Ser. A*, 322(1549):251–280, 1971.
- [vdK80] Wilberd van der Kallen. Homology stability for linear groups. *Invent. Math.*, 60(3):269–295, 1980.
- [Wei94] Charles A. Weibel. *An introduction to homological algebra*, volume 38 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1994.
- [Wen87] Hans Wenzl. On sequences of projections. *C. R. Math. Rep. Acad. Sci. Canada*, 9(1):5–9, 1987.
- [Wes95] B. W. Westbury. The representation theory of the Temperley-Lieb algebras. *Math. Z.*, 219(4):539–565, 1995.

DPMMS, UNIVERSITY OF CAMBRIDGE

Email address: rachaelboyd@dpmms.cam.ac.uk

URL: <https://www.dpmms.cam.ac.uk/~rjb226/>

INSTITUTE OF MATHEMATICS, UNIVERSITY OF ABERDEEN

Email address: r.hepworth@abdn.ac.uk

URL: <http://homepages.abdn.ac.uk/r.hepworth/pages/>