



Crystal structure of 1-butyl-2,3-dimethylimidazolium dicarba-7,8-nido-undecaborate

M. J. Klemes, L. Soderstrom, J. L. Hunting and
A. S. Larsen*

Department of Chemistry, Ithaca College, 953 Danby Road, Ithaca, NY 14850, USA.

*Correspondence e-mail: alarsen@ithaca.edu

Received 25 November 2014; accepted 9 February 2015

Edited by W. T. A. Harrison, University of Aberdeen, Scotland


In the title molecular salt, $C_9H_{17}N_2^+ \cdot C_2H_{12}B_9^-$, the carborane cage has a bridging B—H—B bond on the open B_3C_2 face. The butyl side chain of the cation adopts an extended conformation [C—C—C—C = 179.6 (1)°]. In the crystal, the imidazolium ring is almost coplanar with the open face of the carborane anion. The cations stack in the [010] direction and the dihedral angle between the imidazolium rings of adjacent cations is 68.45 (6)°. The butyl chains extend into the space between carborane anions.

Keywords: crystal structure; carborane cage anion; imidazolium cation; bridging B—H—B bond.

CCDC reference: 1048494

1. Related literature

For structural and thermodynamic properties of the title compound and similar boron cluster anion low-melting ionic compounds, see: Larsen *et al.* (2000); Dymon *et al.* (2008); Suarez *et al.* (2011). A similar bridging hydrogen atom was reported by Jones *et al.* 1997 in an analogous crystal structure.



2. Experimental

2.1. Crystal data

$C_9H_{17}N_2^+ \cdot C_2H_{12}B_9^-$
 $M_r = 286.78$
Monoclinic, $P2_1/n$
 $a = 9.5242$ (2) Å
 $b = 11.5173$ (2) Å
 $c = 16.3357$ (3) Å
 $\beta = 104.821$ (1)°

$V = 1732.30$ (6) Å³
 $Z = 4$
Mo $K\alpha$ radiation
 $\mu = 0.06$ mm⁻¹
 $T = 100$ K
0.42 × 0.32 × 0.26 mm

2.2. Data collection

Bruker SMART CCD 1K area-
detector diffractometer
Absorption correction: multi-scan
(*SADABS*; Bruker, 2012)
 $T_{min} = 0.701$, $T_{max} = 0.746$

27028 measured reflections
3964 independent reflections
3582 reflections with $I > 2\sigma(I)$
 $R_{int} = 0.021$

2.3. Refinement

$R[F^2 > 2\sigma(F^2)] = 0.043$
 $wR(F^2) = 0.120$
 $S = 1.05$
3964 reflections
225 parameters

H atoms treated by a mixture of
independent and constrained
refinement
 $\Delta\rho_{max} = 0.40$ e Å⁻³
 $\Delta\rho_{min} = -0.28$ e Å⁻³

Data collection: *APEX2* (Bruker, 2012); cell refinement: *SAINT* (Bruker, 2012); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL2013* (Sheldrick, 2015); molecular graphics: *OLEX2* (Dolomanov *et al.*, 2009); software used to prepare material for publication: *OLEX2*.

Acknowledgements

AL thanks Professor Maitland Jones for the generous donation of the starting orthocarborane stock and Dr John Holbrey for the supply of imidazolium halide reagents. The research support by ACS PRF and Cottrell College awards (44692.01-GB and CC6755) is gratefully acknowledged.

Supporting information for this paper is available from the IUCr electronic archives (Reference: HB7327).

References

- Bruker (2012). *APEX2*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Dolomanov, O. V., Bourhis, L. J., Gildea, R. J., Howard, J. A. K. & Puschmann, H. (2009). *J. Appl. Cryst.* **42**, 339–341.
- Dymon, J., Wibby, R., Kleingardner, J., Tanski, J. M., Guzei, I. A., Holbrey, J. D. & Larsen, A. S. (2008). *Dalton Trans.* pp. 2999–3006.
- Jones, P. G., Villacampa, M. D., Crespo, O., Gimeno, M. C. & Laguna, A. (1997). *Acta Cryst.* **C53**, 570–572.
- Larsen, A. S., Holbrey, J. D., Tham, F. S. & Reed, C. A. (2000). *J. Am. Chem. Soc.* **122**, 7264–7272.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Sheldrick, G. M. (2015). *Acta Cryst.* **C71**, 3–8.
- Suarez, S. A., Foi, A., Eady, S., Larsen, A. & Doctorovich, F. (2011). *Acta Cryst.* **C67**, o417–o420.

supporting information

Acta Cryst. (2015). E71, o183 [doi:10.1107/S2056989015002765]

Crystal structure of 1-butyl-2,3-dimethylimidazolium dicarba-7,8-nido-undecaborate

M. J. Klemes, L. Soderstrom, J. L. Hunting and A. S. Larsen

S1. Comment


The title compound was synthesized as part of a study of low melting ionic compounds with carborane cage anions. (Larsen *et al.* 2000, Dymon *et al.* 2008, Suarez *et al.* 2011). The formula unit consists of a carborane anion and an alkylated imidazolium cation. A similar bridging hydrogen atom was also seen, for example, by Jones *et al.* 1997 in an analogous crystal structure. In the crystal packing the imidazolium rings are almost coplanar with open face of the carborane anions. The angle between two orientations of coplanar imidazolium rings is 68.45°. The butyl chains form interlinking pattern extending into the space between carborane anions. The carborane anion possesses a bridging hydrogen at the open face of the cage (shown on Figure 1).

S2. Experimental

Caesium dicarba-7,8-nido-undecaborate was synthesized according to published procedure (Dymon *et al.* 2008). Caesium dicarba-7,8-nido-undecaborate (0.300 g, 1.14 mmol) was dissolved in acetone (2 ml). This solution was added to 1-butyl-2,3-dimethylimidazolium chloride (0.215 g, 1.14 mmol) dissolved in dichloromethane (30 ml). The turbid solution was stirred at room temperature for 30 minutes. The mixture was filtered through a plug of celite to remove the caesium chloride precipitate. The volatiles were removed *in vacuo* and the residue was dissolved in dichloromethane (30 ml) and filtered *via* celite again. The dichloromethane was removed *in vacuo*. The solid residue was dissolved in a small amount of absolute ethanol and crystals were grown by slow vapor diffusion of hexane into the absolute ethanol solution at 233 K.

S3. Refinement

Crystal data, data collection and structure refinement details are summarized in Table 1. The methyl and aromatic H atoms were constrained to an ideal geometry; the methyl H atoms were allowed to rotate freely about the C—C bonds. The H atoms attached to B atoms were placed in calculated positions.

**Figure 1**

Crystal structure of the title compound with displacement ellipsoids at the 50% probability level.





Figure 2

Packing diagram of the title compound, showing coplanar alignment of the imidazolium rings and parallel butyl chains. The angle between two orientations of coplanar imidazolium rings is 68.45° .

**Figure 3**

Packing diagram of the title compound showing the imidazolium rings nearly coplanar with the open B3C2 face. For clarity, H-atoms are removed.

1-Butyl-2,3-dimethylimidazolium dicarba-7,8-nido-undecaborate

Crystal data

$\text{C}_9\text{H}_{17}\text{N}_2^+ \cdot \text{C}_2\text{H}_{12}\text{B}_9^-$
 $M_r = 286.78$
 Monoclinic, $P2_1/n$
 $a = 9.5242(2) \text{ \AA}$
 $b = 11.5173(2) \text{ \AA}$
 $c = 16.3357(3) \text{ \AA}$
 $\beta = 104.821(1)^\circ$
 $V = 1732.30(6) \text{ \AA}^3$
 $Z = 4$

$F(000) = 616$
 $D_x = 1.099 \text{ Mg m}^{-3}$
 Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
 Cell parameters from 3582 reflections
 $\theta = 2.2\text{--}27.5^\circ$
 $\mu = 0.06 \text{ mm}^{-1}$
 $T = 100 \text{ K}$
 Prism, clear light colourless
 $0.42 \times 0.32 \times 0.26 \text{ mm}$

Data collection

Bruker SMART CCD 1K area-detector
 diffractometer
 Radiation source: sealed X-ray tube
 Graphite monochromator
 Detector resolution: $7.9 \text{ pixels mm}^{-1}$
 ω scans
 Absorption correction: multi-scan
 (*SADABS*; Bruker, 2012)
 $T_{\min} = 0.701$, $T_{\max} = 0.746$

27028 measured reflections
 3964 independent reflections
 3582 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.021$
 $\theta_{\max} = 27.5^\circ$, $\theta_{\min} = 2.2^\circ$
 $h = -12 \rightarrow 12$
 $k = -14 \rightarrow 14$
 $l = -21 \rightarrow 20$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.043$
 $wR(F^2) = 0.120$

$S = 1.05$
 3964 reflections
 225 parameters
 0 restraints

37 constraints
 Primary atom site location: structure-invariant
 direct methods
 H atoms treated by a mixture of independent
 and constrained refinement

$$w = 1/[\sigma^2(F_o^2) + (0.0633P)^2 + 0.7865P]$$

where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.004$
 $\Delta\rho_{\max} = 0.40 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.28 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
N1	0.76133 (10)	0.49244 (8)	0.20979 (6)	0.0148 (2)
N2	0.67739 (10)	0.47136 (8)	0.31974 (6)	0.0149 (2)
C3	0.85032 (13)	0.52932 (10)	0.15375 (7)	0.0201 (2)
H3a	0.8045 (5)	0.5960 (5)	0.1201 (4)	0.0301 (4)*
H3b	0.9471 (3)	0.5514 (8)	0.18769 (8)	0.0301 (4)*
H3c	0.8590 (8)	0.4653 (3)	0.1159 (4)	0.0301 (4)*
C6	0.90467 (12)	0.59313 (10)	0.34029 (7)	0.0195 (2)
H6a	0.99760 (12)	0.5562 (4)	0.3408 (5)	0.0292 (4)*
H6b	0.9000 (6)	0.6697 (3)	0.3136 (3)	0.0292 (4)*
H6c	0.8963 (6)	0.6019 (7)	0.39853 (18)	0.0292 (4)*
C5	0.78385 (12)	0.51986 (9)	0.29174 (7)	0.0145 (2)
C4	0.63722 (12)	0.42556 (9)	0.18504 (7)	0.0169 (2)
H4	0.59636 (12)	0.39499 (9)	0.13008 (7)	0.0203 (3)*
C7	0.58539 (12)	0.41206 (9)	0.25370 (7)	0.0167 (2)
H7	0.50115 (12)	0.36962 (9)	0.25636 (7)	0.0200 (3)*
C8	0.66132 (12)	0.47245 (10)	0.40748 (7)	0.0165 (2)
H8a	0.72201 (12)	0.53512 (10)	0.43998 (7)	0.0198 (3)*
H8b	0.55882 (12)	0.48834 (10)	0.40668 (7)	0.0198 (3)*
C9	0.70717 (12)	0.35634 (10)	0.45034 (7)	0.0178 (2)
H9a	0.64746 (12)	0.29405 (10)	0.41672 (7)	0.0214 (3)*
H9b	0.80982 (12)	0.34134 (10)	0.45104 (7)	0.0214 (3)*
C10	0.69123 (12)	0.35140 (10)	0.54080 (7)	0.0164 (2)
H10a	0.75173 (12)	0.41286 (10)	0.57497 (7)	0.0197 (3)*
H10b	0.58876 (12)	0.36634 (10)	0.54054 (7)	0.0197 (3)*
C11	0.73738 (13)	0.23351 (10)	0.58104 (7)	0.0205 (2)
H11a	0.7246 (10)	0.2324 (3)	0.6387 (2)	0.0308 (4)*
H11b	0.6775 (7)	0.17259 (13)	0.5473 (3)	0.0308 (4)*
H11c	0.8397 (3)	0.2197 (4)	0.5830 (5)	0.0308 (4)*
C2	0.17288 (12)	0.08417 (9)	0.42860 (7)	0.0147 (2)
C1	0.21122 (12)	0.08264 (9)	0.34140 (7)	0.0149 (2)
B5	0.33450 (13)	0.14839 (11)	0.42313 (8)	0.0167 (2)
H5	0.44690 (13)	0.11671 (11)	0.45362 (8)	0.0201 (3)*
B6	0.22452 (13)	0.21212 (11)	0.48302 (8)	0.0158 (2)
H6	0.26690 (13)	0.22484 (11)	0.55298 (8)	0.0190 (3)*

B1	0.04205 (13)	0.16869 (11)	0.43443 (8)	0.0147 (2)
B2	−0.01661 (13)	0.23783 (11)	0.33083 (8)	0.0150 (2)
B3	0.10220 (13)	0.16468 (11)	0.27283 (8)	0.0153 (2)
B4	0.28625 (13)	0.20985 (11)	0.32130 (8)	0.0161 (2)
H4a	0.36949 (13)	0.22198 (11)	0.28446 (8)	0.0193 (3)*
B9	0.28663 (13)	0.29761 (11)	0.40908 (8)	0.0161 (2)
H9	0.36735 (13)	0.36940 (11)	0.42941 (8)	0.0193 (3)*
B7	0.10328 (13)	0.30870 (10)	0.41803 (8)	0.0147 (2)
H7a	0.06430 (13)	0.38730 (10)	0.44608 (8)	0.0176 (3)*
B8	0.14080 (13)	0.30817 (11)	0.31390 (8)	0.0154 (2)
H8	0.12680 (13)	0.38547 (11)	0.27096 (8)	0.0185 (3)*
H2	0.1878 (15)	0.0099 (13)	0.4581 (9)	0.019 (3)*
H2a	−0.1222 (16)	0.2751 (13)	0.2999 (9)	0.025 (4)*
H1	0.2458 (16)	0.0102 (14)	0.3259 (9)	0.024 (4)*
H1a	−0.0255 (15)	0.1466 (12)	0.4776 (9)	0.020 (3)*
H3	0.0733 (16)	0.1437 (13)	0.2064 (9)	0.022 (4)*
H2b	−0.0116 (18)	0.1414 (15)	0.3104 (10)	0.038 (4)*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N1	0.0161 (4)	0.0128 (4)	0.0155 (4)	0.0003 (3)	0.0040 (3)	0.0013 (3)
N2	0.0157 (4)	0.0125 (4)	0.0168 (5)	0.0006 (3)	0.0049 (3)	−0.0002 (3)
C3	0.0226 (6)	0.0218 (6)	0.0174 (5)	−0.0036 (4)	0.0080 (4)	0.0018 (4)
C6	0.0191 (5)	0.0163 (5)	0.0208 (6)	−0.0031 (4)	0.0010 (4)	−0.0010 (4)
C5	0.0166 (5)	0.0112 (5)	0.0158 (5)	0.0031 (4)	0.0045 (4)	0.0019 (4)
C4	0.0164 (5)	0.0134 (5)	0.0191 (5)	0.0000 (4)	0.0013 (4)	−0.0006 (4)
C7	0.0149 (5)	0.0131 (5)	0.0208 (5)	−0.0009 (4)	0.0025 (4)	−0.0001 (4)
C8	0.0201 (5)	0.0158 (5)	0.0161 (5)	0.0014 (4)	0.0091 (4)	−0.0003 (4)
C9	0.0205 (5)	0.0179 (5)	0.0168 (5)	0.0059 (4)	0.0081 (4)	0.0017 (4)
C10	0.0174 (5)	0.0179 (5)	0.0142 (5)	0.0033 (4)	0.0045 (4)	−0.0011 (4)
C11	0.0258 (6)	0.0212 (6)	0.0150 (5)	0.0065 (4)	0.0059 (4)	0.0013 (4)
C2	0.0176 (5)	0.0120 (5)	0.0151 (5)	0.0005 (4)	0.0055 (4)	0.0016 (4)
C1	0.0166 (5)	0.0129 (5)	0.0165 (5)	0.0003 (4)	0.0067 (4)	−0.0014 (4)
B5	0.0138 (5)	0.0173 (6)	0.0187 (6)	0.0015 (4)	0.0035 (4)	0.0006 (5)
B6	0.0175 (6)	0.0153 (6)	0.0142 (5)	0.0010 (4)	0.0032 (4)	−0.0003 (4)
B1	0.0152 (5)	0.0145 (6)	0.0156 (6)	0.0002 (4)	0.0060 (4)	0.0013 (4)
B2	0.0128 (5)	0.0152 (6)	0.0167 (6)	0.0004 (4)	0.0033 (4)	0.0007 (4)
B3	0.0159 (6)	0.0161 (6)	0.0146 (5)	−0.0017 (4)	0.0052 (4)	−0.0005 (4)
B4	0.0150 (5)	0.0167 (6)	0.0180 (6)	−0.0015 (4)	0.0067 (4)	−0.0000 (5)
B9	0.0144 (5)	0.0156 (6)	0.0178 (6)	−0.0021 (4)	0.0034 (4)	−0.0005 (4)
B7	0.0161 (6)	0.0126 (5)	0.0157 (5)	0.0006 (4)	0.0048 (4)	−0.0007 (4)
B8	0.0169 (6)	0.0139 (5)	0.0157 (6)	−0.0011 (4)	0.0047 (4)	0.0018 (4)

Geometric parameters (Å, °)

N1—C3	1.4608 (14)	C1—B5	1.7125 (16)
N1—C5	1.3382 (14)	C1—B3	1.6219 (16)

N1—C4	1.3819 (14)	C1—B4	1.6986 (16)
N2—C5	1.3367 (14)	C1—H1	0.955 (16)
N2—C7	1.3836 (14)	B5—H5	1.1200
N2—C8	1.4798 (13)	B5—B6	1.7656 (17)
C3—H3a	0.9800	B5—B4	1.7574 (18)
C3—H3b	0.9800	B5—B9	1.7778 (18)
C3—H3c	0.9800	B6—H6	1.1200
C6—H6a	0.9800	B6—B1	1.7873 (17)
C6—H6b	0.9800	B6—B9	1.7732 (18)
C6—H6c	0.9800	B6—B7	1.7524 (17)
C6—C5	1.4827 (15)	B1—B2	1.8238 (17)
C4—H4	0.9500	B1—B7	1.7584 (17)
C4—C7	1.3451 (16)	B1—H1a	1.099 (14)
C7—H7	0.9500	B2—B3	1.8538 (17)
C8—H8a	0.9900	B2—B7	1.7796 (17)
C8—H8b	0.9900	B2—B8	1.7868 (17)
C8—C9	1.5208 (15)	B2—H2a	1.091 (15)
C9—H9a	0.9900	B2—H2b	1.165 (17)
C9—H9b	0.9900	B3—B4	1.8056 (17)
C9—C10	1.5246 (14)	B3—B8	1.7862 (17)
C10—H10a	0.9900	B3—H3	1.077 (14)
C10—H10b	0.9900	B3—H2b	1.401 (16)
C10—C11	1.5236 (15)	B4—H4a	1.1200
C11—H11a	0.9800	B4—B9	1.7536 (18)
C11—H11b	0.9800	B4—B8	1.7693 (18)
C11—H11c	0.9800	B9—H9	1.1200
C2—C1	1.5585 (14)	B9—B7	1.7938 (17)
C2—B5	1.7305 (16)	B9—B8	1.8046 (18)
C2—B6	1.7263 (16)	B7—H7a	1.1200
C2—B1	1.6030 (16)	B7—B8	1.8254 (17)
C2—H2	0.975 (15)	B8—H8	1.1200
C5—N1—C3	126.02 (10)	B6—B1—C2	60.93 (7)
C4—N1—C3	124.66 (10)	B2—B1—C2	105.57 (8)
C4—N1—C5	109.30 (9)	B2—B1—B6	108.80 (8)
C7—N2—C5	108.90 (9)	B7—B1—C2	104.76 (8)
C8—N2—C5	127.11 (9)	B7—B1—B6	59.23 (7)
C8—N2—C7	123.89 (9)	B7—B1—B2	59.54 (7)
H3a—C3—N1	109.5	H1a—B1—C2	119.3 (7)
H3b—C3—N1	109.5	H1a—B1—B6	116.2 (7)
H3b—C3—H3a	109.5	H1a—B1—B2	127.3 (7)
H3c—C3—N1	109.5	H1a—B1—B7	125.5 (7)
H3c—C3—H3a	109.5	B3—B2—B1	101.16 (8)
H3c—C3—H3b	109.5	B7—B2—B1	58.40 (7)
H6b—C6—H6a	109.5	B7—B2—B3	105.48 (8)
H6c—C6—H6a	109.5	B8—B2—B1	105.76 (8)
H6c—C6—H6b	109.5	B8—B2—B3	58.73 (7)
C5—C6—H6a	109.5	B8—B2—B7	61.57 (7)

C5—C6—H6b	109.5	H2a—B2—B1	128.9 (8)
C5—C6—H6c	109.5	H2a—B2—B3	123.6 (8)
N2—C5—N1	107.59 (9)	H2a—B2—B7	121.1 (8)
C6—C5—N1	124.98 (10)	H2a—B2—B8	117.4 (8)
C6—C5—N2	127.42 (10)	H2b—B2—B1	79.9 (8)
H4—C4—N1	126.59 (6)	H2b—B2—B3	49.1 (8)
C7—C4—N1	106.82 (10)	H2b—B2—B7	127.0 (8)
C7—C4—H4	126.59 (7)	H2b—B2—B8	106.9 (8)
C4—C7—N2	107.39 (9)	H2b—B2—H2a	110.1 (11)
H7—C7—N2	126.31 (6)	B2—B3—C1	106.17 (8)
H7—C7—C4	126.31 (7)	B4—B3—C1	59.13 (7)
H8a—C8—N2	109.56 (6)	B4—B3—B2	107.22 (8)
H8b—C8—N2	109.56 (6)	B8—B3—C1	104.23 (8)
H8b—C8—H8a	108.1	B8—B3—B2	58.76 (7)
C9—C8—N2	110.48 (9)	B8—B3—B4	59.02 (7)
C9—C8—H8a	109.56 (6)	H3—B3—C1	121.3 (8)
C9—C8—H8b	109.56 (6)	H3—B3—B2	125.8 (8)
H9a—C9—C8	108.98 (6)	H3—B3—B4	118.4 (8)
H9b—C9—C8	108.98 (6)	H3—B3—B8	124.1 (8)
H9b—C9—H9a	107.8	H2b—B3—C1	90.9 (7)
C10—C9—C8	113.02 (9)	H2b—B3—B2	38.9 (7)
C10—C9—H9a	108.98 (6)	H2b—B3—B4	129.5 (7)
C10—C9—H9b	108.98 (6)	H2b—B3—B8	96.9 (7)
H10a—C10—C9	109.39 (6)	H2b—B3—H3	111.8 (10)
H10b—C10—C9	109.39 (6)	B5—B4—C1	59.38 (7)
H10b—C10—H10a	108.0	B3—B4—C1	55.04 (6)
C11—C10—C9	111.23 (9)	B3—B4—B5	106.77 (8)
C11—C10—H10a	109.39 (6)	H4a—B4—C1	127.03 (5)
C11—C10—H10b	109.39 (6)	H4a—B4—B5	120.62 (6)
H11a—C11—C10	109.5	H4a—B4—B3	122.79 (5)
H11b—C11—C10	109.5	B9—B4—C1	104.60 (8)
H11b—C11—H11a	109.5	B9—B4—B5	60.84 (7)
H11c—C11—C10	109.5	B9—B4—B3	108.87 (8)
H11c—C11—H11a	109.5	B9—B4—H4a	120.92 (6)
H11c—C11—H11b	109.5	B8—B4—C1	101.80 (8)
B5—C2—C1	62.50 (7)	B8—B4—B5	109.46 (9)
B6—C2—C1	112.20 (8)	B8—B4—B3	59.94 (7)
B6—C2—B5	61.43 (7)	B8—B4—H4a	122.20 (6)
B1—C2—C1	115.37 (9)	B8—B4—B9	61.62 (7)
B1—C2—B5	117.30 (9)	B6—B9—B5	59.63 (7)
B1—C2—B6	64.82 (7)	B4—B9—B5	59.68 (7)
H2—C2—C1	113.8 (8)	B4—B9—B6	107.80 (9)
H2—C2—B5	112.5 (8)	H9—B9—B5	122.78 (5)
H2—C2—B6	120.6 (8)	H9—B9—B6	122.09 (6)
H2—C2—B1	121.4 (8)	H9—B9—B4	121.72 (6)
B5—C1—C2	63.68 (7)	B7—B9—B5	106.33 (8)
B3—C1—C2	111.49 (8)	B7—B9—B6	58.85 (7)
B3—C1—B5	118.09 (9)	B7—B9—B4	108.02 (8)

B4—C1—C2	112.05 (8)	B7—B9—H9	122.14 (6)
B4—C1—B5	62.02 (7)	B8—B9—B5	106.97 (8)
B4—C1—B3	65.83 (7)	B8—B9—B6	107.88 (8)
H1—C1—C2	115.4 (9)	B8—B9—B4	59.62 (7)
H1—C1—B5	112.3 (9)	B8—B9—H9	121.63 (5)
H1—C1—B3	122.0 (9)	B8—B9—B7	60.97 (7)
H1—C1—B4	120.8 (9)	B1—B7—B6	61.21 (7)
C1—B5—C2	53.82 (6)	B2—B7—B6	112.47 (8)
H5—B5—C2	126.83 (5)	B2—B7—B1	62.06 (7)
H5—B5—C1	126.11 (5)	B9—B7—B6	59.99 (7)
B6—B5—C2	59.17 (7)	B9—B7—B1	108.35 (8)
B6—B5—C1	103.31 (8)	B9—B7—B2	109.93 (8)
B6—B5—H5	121.61 (6)	H7a—B7—B6	120.08 (6)
B4—B5—C2	101.56 (8)	H7a—B7—B1	121.43 (5)
B4—B5—C1	58.60 (7)	H7a—B7—B2	119.10 (5)
B4—B5—H5	122.74 (6)	H7a—B7—B9	121.49 (6)
B4—B5—B6	107.97 (8)	B8—B7—B6	107.87 (8)
B9—B5—C2	102.58 (8)	B8—B7—B1	106.89 (8)
B9—B5—C1	102.99 (8)	B8—B7—B2	59.41 (7)
B9—B5—H5	123.83 (5)	B8—B7—B9	59.81 (7)
B9—B5—B6	60.05 (7)	B8—B7—H7a	123.25 (5)
B9—B5—B4	59.47 (7)	B3—B8—B2	62.51 (7)
B5—B6—C2	59.40 (7)	B4—B8—B2	111.87 (8)
H6—B6—C2	128.17 (5)	B4—B8—B3	61.04 (7)
H6—B6—B5	120.64 (6)	B9—B8—B2	109.11 (8)
B1—B6—C2	54.26 (6)	B9—B8—B3	107.48 (8)
B1—B6—B5	106.55 (8)	B9—B8—B4	58.76 (7)
B1—B6—H6	123.06 (5)	B7—B8—B2	59.02 (7)
B9—B6—C2	102.94 (8)	B7—B8—B3	106.41 (8)
B9—B6—B5	60.31 (7)	B7—B8—B4	105.96 (8)
B9—B6—H6	121.77 (6)	B7—B8—B9	59.23 (7)
B9—B6—B1	107.98 (8)	H8—B8—B2	119.15 (5)
B7—B6—C2	99.97 (8)	H8—B8—B3	121.34 (5)
B7—B6—B5	108.69 (8)	H8—B8—B4	121.15 (6)
B7—B6—H6	123.01 (6)	H8—B8—B9	122.36 (5)
B7—B6—B1	59.56 (7)	H8—B8—B7	124.10 (5)
B7—B6—B9	61.16 (7)	B3—H2b—B2	92.0 (11)
